

139151

VOLUME II

Part 2

GEODYN PROGRAMMER'S GUIDE

Contract No.: NAS 5-11735 - MOD 65
PCN 550-W-72416

(NASA-CR-139151) GEODYN PROGRAMMER'S GUIDE, VOLUME 2, PART 2 (Wolf Research and Development Corp.) 276 p HC \$8.75
CSCI 08E
G3/43
N75-11416
Unclas
03463

Prepared by


N.E. Mullins
N.C. Dao

T.V. Martin
C.C. Goad

N.L. Boulware
M.M. Chin

Wolf Research and Development Corporation
Riverdale, Maryland

For
Goddard Space Flight Center
Greenbelt, Maryland



October 1972

TABLE OF CONTENTS

		<u>Page</u>
	INTRODUCTION	i
1.0	INTRODUCTION TO THE GEODYN PROGRAM	1.0-1
2.0	GEODYN ENVIRONMENTAL REQUIREMENTS	2.0-1
3.0	DIAGRAM OF OVERLAY STRUCTURE	3.0-1
4.0	DIAGRAMS OF SUBROUTINE STRUCTURE	4.0-1
5.0	SUMMARY OF SUBPROGRAMS USED BY GEODYN	5.0-1
6.0	SUBROUTINE CROSS REFERENCE CHART	6.0-1
7.0	COMMON BLOCK CROSS REFERENCE CHART	7.0-1
8.0	PROGRAM DESCRIPTIONS	8.0-1
9.0	COMMON BLOCK DESCRIPTIONS	9.0-1

ORBIT

DESCRIPTION

ORBIT is the executive control routine for the orbit integration.

At the beginning of each arc it initializes required program constants as well as the variational partials at epoch. If epoch needs to be reset to a previous time, ORBIT negates the stepsize, and calls for COWELL integration backwards to the desired time. After backward integration is completed, ORBIT resets the stepsize to the proper positive quantity.

For each time point for which it is called, ORBIT performs the following tasks:

- If necessary, calls subroutine COWELL to integrate the orbit further.
- Calls INTRP to obtain values for the position, velocity. In the data reduction mode, variational partials associated with the orbit parameters are also calculated.
- Converts position and velocity from true equator and equinox of reference day to true equator and equinox of date using subroutine REFCOR.

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

NAME ORBIT
ENTRY POINT PURPOSE
ORBIT1 TO INITIALIZE
ORBIT TO RETURN SATELLITE STATE (POSITION & VELOCITY)
AND FORCE MODEL PARTIALS AT THE CALLED TIME (DAY)

CALLING SEQUENCE CALL CRBIT1(FCT,SUM,XPART)

SYMBOL	TYPE	DESCRIPTION
FCT (3,1)	DP	INPUT - ACCELERATION ARRAY (ORBIT AND FORCE MODEL PARTIALS)
SUM (2,3,1)	DP	INPUT - SUM ARRAY USED BY SUMMED-COWELL INTEGRATOR & INTERPOLATOR
XPART (6,NCONMX,1)	DP	OUTPUT - ORBIT AND FORCE MODEL PARTIAL MATRIX

CALLING SEQUENCE CALL CRBIT(DAY)

SYMBOL	TYPE	DESCRIPTION
DAY	DP	INPUT - DESIRED OUTPUT TIME

SUBROUTINES USED COWELL INTFP SLEN DATES REFCOR

COMMON BLOCKS INTBLK INITBK INTERP PFIORI XYZOUT
CTIME CELEM XYZ APARAM CONOUT

INPLT FILES NONE

OUTPUT FILES PRINTER

REFERENCES 'GEODYN SYSTEMS DESCRIPTION'
VOLUME 1 - GEODYN DOCUMENTATION

SUBROUTINE CRBIT1(FCT,SUM,XPART)	CRBI	44
REAL*8 DAY2(DAY,SECT,XPART(6,NCONMX,1),SUM(2,3,1),STEPSZ,FCT(3,1))	CRBI	45
REAL*8 LLENST,GV,THET25,AE,ALSC,DAYK,M(4),TIM(4),BDUT,GRBE	CRBI	46
REAL*8 MSUN,PMOON,MSAT,GNE,G,EO,APCM,APLY,PPRESS,ASAT	CRBI	47
REAL*8 EPS(C,BODY,CC,CLD,PMISS,LLMER,FAC,LATAID,XYZENT,	CRBI	48
• ERGPAR,CRBELA,INT1,INT2,PP(20,2),P(20,2),CC(20,2),C(20,2),	CRBI	49
• VCC(20,2),VC(20,2)	CRBI	50
INTEGER SPAL,ADLR,ORDER,ICOR(4)	CRBI	51
LOGICAL TORSET,INITAL,VAI,STP,PLVES,INITC	CRBI	52
COMMON/INTBLK/THOUT(4),THET25,CH,AE,ALSC,PLAT(5),GNE(6),P(2),	CRBI	53
• BDUT(2),P(2),ARGM(2),APLM(2),PP(20,2),INITAL,NORPAT,THETG(2),	CRBI	54
• MBODY(6),STEPSZ(4),P(20,2),ICOR(4)	CRBI	55

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

```

    • ASAT(2),MSAT(2),VARSTP(2),HLVCSK(2),NECN(2),ACER(4),
    • SRAC(2),ISPAD(6),TORCFT,NOCY
  COMMON/INITOR/ISPYMD,ILPHM,EFSLC,ICUMI(4),OREEL(6,2),ICUMK(25)
  COMMON/INTERP/IDUM(882),M(2,2)
  COMMON/PPIDRI/LEMIN(6,2),ICLE(156),CC(2),CDD(2),EMISS(2)
  COMMON/XYZOLT/XYZLND(6,2),DRGPAR(6,2)
  COMMON/CTIME/DATARP,DAYREF(2),DAYC,DAYSTP(17)
  COMMON/CELEM/LEFMS(6,2),ORRELA(6,2),TORREL(5)
  COMMON/XYZ/XYZDUM(16),ISAT,IFORCF(2)
  COMMON/APAR/AM/KKS(4),NSAT,KD(5)
  COMMON/CCNBLT/MINOUT(8),NECNMX,IVAR(10)
  RETURN
  ENTRY ORBIT(DAY)
  INITO=INITAL
  IF(.NOT.INITAL)GO TO 9
  IBAC=0
  C SET STEP SIZE AND INTEGRATION ORDER
  M(1)=STEPSZ(1)
  M(2)=STEPSZ(3)
  M(3)=STEPSZ(2)
  M(4)=STEPSZ(4)
  IORDER(1)=ORDER(1)
  IORDER(2)=ORDER(3)
  IORDER(3)=ORDER(2)
  IORDER(4)=ORDER(4)
  AFSQ=AF**2
  DO 50 I=1,6
  50 GM2(I)=GM*MODY(I)
  9 ISAT=0
  IPT1=1
  IDISP=1
  10 ISAT=ISAT+1
  ISATL1=ISAT-1
  IF(ISAT.GT.NSAT)RETURN
  C DETERMINE STARTING LOCATION OF ARRAYS IF MORE THAN 1 SATELLITE
  IF(ISAT.GT.1)IDISP=IDISP+M(1,ISATL1)+(NECN(ISATL1)-1)*M(2,ISATL1)
  IPT=1+(ISATL1)*2
  IF(ISAT.GT.1)IPT1=IPT1+NECN(ISATL1)
  IF(.NOT.INITAL)GO TO 200
  C INITIALIZE
  M(1,ISAT)=0
  M(2,ISAT)=0
  TIM(IPT)=E.64E4*DAYC
  TIM(IPT+1)=TIM(IPT)
  BC(ISAT)=C.DC
  IF(MSAT(ISAT).GT.0.D0)BC(ISAT)=.5E3*ASAT(ISAT)/MSAT(ISAT)
  C DRAG
  B(ISAT)=B.C(ISAT)*CC(ISAT)
  EDCT(ISAT)=F.C(ISAT)*CDD(ISAT)
  APLV(ISAT)=C.DC
  IF(MSAT(ISAT).GT.C.DC)AFLM(ISAT)=/SAT(ISAT)*FPPSS/MSAT(ISAT)
  C SOLAR REFLECTIVITIES
  APMV(ISAT)=APLV(ISAT)*EMISS(ISAT)
  IF(NECN(ISAT).E.O.1)SRAC(ISAT)=0
  1 DO 100 I=1,6
  C INITIALIZE ORBIT
  ORBI 56
  ORBI 57
  ORBI 58
  ORBI 59
  ORBI 60
  ORBI 61
  ORBI 62
  ORBI 63
  ORBI 64
  ORBI 65
  ORBI 66
  ORBI 67
  ORBI 68
  ORBI 69
  ORBI 69
  ORBI 70
  ORBI 71
  ORBI 72
  ORBI 73
  ORBI 74
  ORBI 75
  ORBI 76
  ORBI 77
  ORBI 78
  ORBI 79
  ORBI 80
  ORBI 81
  ORBI 82
  ORBI 83
  ORBI 84
  ORBI 85
  ORBI 86
  ORBI 87
  ORBI 88
  ORBI 89
  ORBI 90
  ORBI 91
  ORBI 92
  ORBI 93
  ORBI 94
  ORBI 95
  ORBI 96
  ORBI 97
  ORBI 98
  ORBI 99
  ORBI 100
  ORBI 101
  ORBI 102
  ORBI 103
  ORBI 104
  ORBI 105
  ORBI 106
  ORBI 107
  ORBI 108
  ORBI 109
  ORBI 110
  ORBI 111
  
```

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR




```



      XPART(1,1,ISAT)=ELEMST(1,ISAT)
150 CONTINUE
      IF(NEQN(ISAT).EQ.1)GOTO190
C INITIALIZE PARTIALS
      NNG=NEQN(ISAT)
      DO 195 I=1,6
      DO 170 J=2,NNG
170 XPART(1,J,ISAT)=0.00
160 XPART(1,I+1,ISAT)=1.00
190 IF(1BAC.EQ.ISAT)GOTO650
      IF(DAYC.LG.DATARP)GOTO650
C INTEGRATE BACKWARDS IN TIME IF TIME FCINT DESIRED PRECEDES EPOCH
      H(IPT)=-H(IPT)
      CALL COMELL(DATARP,XPART(1,1,ISAT),FCT(1,DISP),IOFCOR(IPT),
      • H(IPT),TIM(IPT),SUM(1,1,IPT1),VARSTP(ISAT),-1.00,M(1,ISAT),
      • 1,ISAT,PP(1,ISAT),P(1,ISAT),CC(1,ISAT),C(1,ISAT),VCC(1,ISAT),
      • VC(1,ISAT))
      SEC1=(8.6404*DATARP-TIM(IPT))/H(IPT)
      CALL INTER(SEC1,H(IPT),IOFCOR(IPT),1,ELEMST(1,ISAT),FCT(1,DISP),
      • H(1,ISAT),SUM(1,1,IPT1))
      CALL ELEM(ELEMST(1,ISAT),ORBELA(1,ISAT),1,TFUE,ORBELA(1,ISAT))
      DO 240 I=1,6
      ORBEL(I,ISAT)=ORBELA(I,ISAT)
      ELFMIN(I,ISAT)=ELEMST(I,ISAT)
240 CONTINUE
      1BAC=1BAC+1
      INITAL=.TRUE.
C HAVING FINISHED BACKWARD INTEGRATION, REINITIALIZE TO BEGIN FORWARD
C INTEGRATION
      H(IPT)=-H(IPT)
      TIM(IPT)=8.6404*DATARP
      TIM(IPT+1)=TIM(IPT)
      H(IPT)=DMIN(H(IPT),STEPSZ(ISAT))
      FAC=H(IPT)/STEPSZ(ISAT)
      IF(FAC.NE.1.00)H(IPT+1)=STEPSZ(ISAT+2)*FAC
      IF(1BAC.LT.NSAT)GOTO 1
      DAYC=DATARP
C RESET EPOCH TIME
      CALL DATES(DAYC,IEPYMD,IEPHM,OFSEC)
      GO TO 1
650 CONTINUE
      CALL COMELL(DAY,XPART(1,1,ISAT),FCT(1,DISP),IOFCOR(IPT),H(IPT),
      • TIM(IPT),SUM(1,1,IPT1),VARSTP(ISAT),1.00,M(1,ISAT),NEQN(ISAT),
      • ISAT,PP(1,ISAT),P(1,ISAT),CC(1,ISAT),C(1,ISAT),VCC(1,ISAT),
      • VC(1,ISAT))
      IF(ISAT.LT.NSAT)INITAL=INITC
      GO TO 225
C DETERMINE IF INTEGRATION IS NEEDED
200 DINT1=8.6404*DAY
      DINT2=DINT1+2.00*H(IPT+1)
      DINT3=DINT1+H(IPT)+H(IPT)
      IF(DINT1.GE.TIM(IPT).OR.(NEQN(ISAT).GT.1.AND.DINT2.GE.
      • TIM(IPT+1)))CALL COMELL(DAY,XPART(1,1,ISAT),FCT(1,DISP),
      • IOFCOR(IPT),H(IPT),TIM(IPT),SUM(1,1,IPT1),VARSTP(ISAT),
      • 1.00,M(1,ISAT),NEQN(ISAT),ISAT,PP(1,ISAT),P(1,ISAT),CC(1,ISAT),
      • C(1,ISAT),VCC(1,ISAT),VC(1,ISAT))

```


ORBI 112
ORBI 113
ORBI 114
ORBI 115
ORBI 116
ORBI 117
ORBI 118
ORBI 119
ORBI 120
ORBI 121
ORBI 122
ORBI 123
ORBI 124
ORBI 125
ORBI 126
ORBI 127
ORBI 128
ORBI 129
ORBI 130
ORBI 131
ORBI 132
ORBI 133
ORBI 134
ORBI 135
ORBI 136
ORBI 137
ORBI 138
ORBI 139
ORBI 140
ORBI 141
ORBI 142
ORBI 143
ORBI 144
ORBI 145
ORBI 146
ORBI 147
ORBI 148
ORBI 149
ORBI 150
ORBI 151
ORBI 152
ORBI 153
ORBI 154
ORBI 155
ORBI 156
ORBI 157
ORBI 158
ORBI 159
ORBI 160
ORBI 161
ORBI 162
ORBI 163
ORBI 164
ORBI 165
ORBI 166
ORBI 167

C COMPUTE INTEGRATION TIME IN FRACTIANS OF STEPSIZES FROM INTEGRATOR	ORBI 168
C TIME	ORBI 169
225 SEC1=(3.6404*DAY-TIM(IPT))/H(IPT)	ORBI 170
CALL INTRP(SEC1,H(IPT),ICOR(IFT),1,XPART(1,1,ISAT),FCT(1,DISP),	ORBI 171
• M(1,ISAT),SUM(1,1,IPT))	ORBI 172
250 IF(NCON(ISAT).LT.2)GO TO 300	ORBI 173
SEC1=(3.6404*DAY-TIM(IPT+1))/H(IPT+1)	ORBI 174
CALL INTRP(SEC1,H(IPT+1),ICOR(IFT+1),NCON(ISAT)-1,	ORBI 175
• XPART(1,2,ISAT),FCT(1,DISP+M(1,ISAT)),M(2,ISAT),	ORBI 176
• SUM(1,1,IPT+1))	ORBI 177
300 IF(TDREFT)GO TO 700	ORBI 178
C CORRECT ORBIT FROM TRUE OF REFERENCE TO TRUE OF DATE	ORBI 179
CALL RLFCOR(DAY,.FALSE.,XPART(1,1,ISAT))	ORBI 180
CALL REFCOR(DAY,.FALSE.,XPART(4,1,ISAT))	ORBI 181
C STORE ORBIT IN COMMON	ORBI 182
750 DO 800 I=1,6	ORBI 183
800 XYZEND(I,ISAT)=XPART(I,1,ISAT)	ORBI 184
IF(ADJR(ISAT).EQ.0)GOTO10	ORBI 185
C STORE DRAG PARTIALS IN COMMON	ORBI 186
DO 900 I=1,6	ORBI 187
900 DRGPAR(I,ISAT)=XPART(I,8,ISAT)	ORBI 188
GO TO 10	ORBI 189
END	ORBI 190





REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR



NAME OUTRAD

PURPOSE CONVERTS INPUT IN RADIAN TO EITHER DEGREES,
 MINUTES AND SECONDS OR HOURS, MINUTES AND SECONDS

CALLING SEQUENCE CALL CUTRAD(RAD, IH, IM, S, K)

SYMBOL	TYPE	DESCRIPTION
RAD	DP	INPUT - ANGLE TO BE CONVERTED IN RADIAN
IH	I	OUTPUT - SIGNED DEGREES OF HOURS
IM	I	OUTPUT - UNSIGNED MINUTES OF ARC OR TIME
S	DP	OUTPUT - UNSIGNED SECONDS OF ARC OR TIME
K	I	INPUT - SWITCH FOR OUTPUT ... K=1 OUTPUT WILL BE IN TIME UNITS K=2 OUTPUT WILL BE IN ARC UNITS

SUBROUTINES USED NONE

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE



SUBROUTINE OUTRAD(RAD, IH, IM, S, K)	JUTP	33
REAL*8 RAD, S, RADIAN, SMIN	JUTR	34
C CONVERT TO DECIMAL DEGREES	OUTR	35
RADIAN=RAD*57.295779500	JUTP	36
C IF OUTPUT IS TO BE IN HOURS, DIVIDE BY 15	OUTR	37
IF(K.EQ.2) RADIAN=RADIAN/15.00	JUTR	38
C EXTRACT HOURS OR DEGREES	OUTR	39
IH=RADIAN	JUTP	40
H=IH	JUTP	41
C EXTRACT MINUTES	OUTP	42
SMIN=(RADIAN-H)*60.00	JUTR	43
IM=SMIN	JUTR	44
C EXTRACT SECONDS	JUTR	45
S=(SMIN-IM)*60.000	JUTR	46
RETURN	JUTR	47
END	JUTP	48

NAME PCERD
 PURPOSE TO READ PCE FORMAT DATA
 CALLING SEQUENCE CALL PCERD
 SUBROUTINES USED RANDWR YMDAY TDIF
 COMMON BLOCKS APARAM CGE05 CONSTS CTIME INTBLK
 PREBLK SIGBLK
 INPUT FILES NONE
 OUTPUT FILES NONE
 REFERENCES 'GECODYN SYSTEMS DESCRIPTION' - APPENDIX C
 VOLUME 3 - GECODYN DOCUMENTATION

```

SUBROUTINE PCERD
  IMPLICIT REAL*8 (A-H,C-Z)
  LOGICAL*4 OKSAT,VHFCHN,PREPRC
  LOGICAL NORATE
  INTEGER*2 CULL,CHANEL,NMEAS,MTYPE,PRETYP,IMTYPE,ISTNO,ISATNO,ITYPE
  INTEGER PCNO
  REAL TDIF,SIGSTC,ISATNO,SIGCHG,SGPNT
  COMMON/CGE05/ISATNO,IPREPR(251),NPRE,NSIG,NCULL,SIGCHG(50)
  * INTYPE(50),ISTNO(50),CULL(2,100)
  COMMON/CONSTS/PI,D2R,S2R
  COMMON/CTIME/DATARP,DAYREF,DAYE,DAYSTP,DAYINT(15)
  COMMON/INTBLK/INTBK1(53),NORATE,INTBK2(78)
  COMMON/PREBLK/DAY,CBS1,CBS2,SIG1,SIG2,SRFNOX,ISTA,MTYPE,NMEAS,
  * ISATNO,PRETYP,CHANEL,VHFCHN,PREPRC,PCNO
  COMMON/SIGBLK/SIGSTC(30),SGPNT(30),ICBS,ICTAPE(3)
  ATIME(DAY)=TDIF(4,3,DAY)/8.64E4
  IF(ICBS.LL.C) ICBS=20
C INITIALIZE
  NUMBER=C
  NORATE=.FALSE.
  SIGC=C.CC
  ISTA=C
  ISATNO=1
  NMEAS=1
  PRETYP=C
  CHANEL=C
  VHFCHN=.FALSE.
  PREPRC=.FALSE.
  SRFNOX=C.CC
  CBS2=C.CC
C READ OBSERVATIONS
  DO FLAG(ICBS,100,ERR)=10,END=200) MTYPE,IMNO,IFM,SEC,CBS1,SIG1
  MTYPE=MTYPE-1
  IF(IMNO.LT.C) GO TO 200
  PCER 22
  PCER 23
  PCER 24
  PCER 25
  PCER 26
  PCER 27
  PCER 28
  PCER 29
  PCER 30
  PCER 31
  PCER 32
  PCER 33
  PCER 34
  PCER 35
  PCER 36
  PCER 37
  PCER 38
  PCER 39
  PCER 40
  PCER 41
  PCER 42
  PCER 43
  PCER 44
  PCER 45
  PCER 46
  PCER 47
  PCER 48
  PCER 49
  PCER 50
  PCER 51
  PCER 52
  PCER 53
  PCER 54
  PCER 55
  
```

IF(MTYPE.LT.15.OR.MTYPE.GT.26) GO TO 10	PCER	56
C CONVERT TIME TO A.1 DAYS FROM JAN 0.0 OF REFERENCE YEAR	PCER	57
DAY=YMDAY(IYMD,IHM,SEC)	PCER	58
DAY=DAY+ATIME(DAY)	PCER	59
IF(DAY.LT.DATSTP) GO TO 10	PCER	60
IF(DAY.GT.DAYSTP) GO TO 200	PCER	61
IF(SIG1.LE.C.CC) SIG1=SIGSTO(MTYPE)	PCER	62
IF(NSIG.LE.C) GO TO 50	PCER	63
NNI=0	PCER	64
C CHECK FOR SIGMA CHANGES	PCER	65
DO 20 I=1,NSIG	PCER	66
IF(ISTNC(I).EQ.C.AND.I(MTYPE(I).EQ.MTYPE) NNI=I	PCER	67
20 CONTINUE	PCER	68
IF(NNI.GT.0) SIG1=SIGCHG(NNI)	PCER	69
50 IF(MTYPE.LT.21.AND.MTYPE.GT.17) SIG1=SIG1*1.0E-2	PCER	70
IF(MTYPE.EQ.22) SIG1=SIG1*1.0E-6	PCER	71
100 IF(MTYPE.LE.22) GO TO 110	PCER	72
OR S1=JOB S1+L2F	PCER	73
SIG1=SIG1*S2F	PCER	74
110 NUMBER=NUMBER+1	PCER	75
IF(NCULL.LE.C) GO TO 260	PCER	76
DO 240 I=1,NCULL	PCER	77
IF(NUMBER-CULL(1,I)) 240,230,220	PCER	78
220 IF(NUMBER.GT.CULL(2,I)) GO TO 240	PCER	79
230 SIG1=0.CC	PCER	80
240 CONTINUE	PCER	81
260 RECNO=RECNO+1	PCER	82
C STORE OBSERVATIONS	PCER	83
CALL RANDWP	PCER	84
GO TO 10	PCER	85
C STORE LAST OBSERVATION AS ZERO	PCER	86
200 MTYPE=0	PCER	87
RECNO=RECNO+1	PCER	88
CALL RANDWP	PCER	89
PRINT 2000,NUMBER,JOBS	PCER	90
DAYSTP=DAY	PCER	91
RETURN	PCER	92
1000 FORMAT(1X,I2,I6,I4,F8.4,D24.16,E10.4)	PCER	93
2000 FORMAT(1H07/SIX,I6,' OBSERVATIONS SELECTED FROM MASTER PCE DATA ',	PCER	94
' TAPE NUMBER',I3)	PCER	95
END	PCER	96



NAME	PDEN
ENTRY POINT	PURPOSE
PDLN2	INITIALIZATION
PDEN	TO PRINT ADJUSTED SURFACE DENSITIES
CALLING SEQUENCE	CALL PDEN2(BCENTR, AREA, DEN, ADJDEN, APSIG, SUM1, DENCON, SAVSIG)

SYMBOL	TYPE	DESCRIPTION
BCENTR (2,2)	DP	INPUT - THE LATITUDE AND LONGITUDE OF THE ADJUSTED SURFACE DENSITY BLOCKS
AREA (4,1)	DP	INPUT - SURFACE DENSITY SUB-BLOCK AREAS
DEN (1)	DP	INPUT - SURFACE DENSITY VALUES
ADJDEN (1)	DP	INPUT & OUTPUT - ADJUSTED SURFACE DENSITY VALUES
APSIG (1)	DP	INPUT - PRIORI SIGMAS FOR ADJUSTED DENSITIES
SUM1 (1)	DP	INPUT - NORMAL MATRIX
DENCON (NCONST,1)	DP	INPUT - MATRIX OF CONSTRAINT EQUATION
SAVSIG (1)	DP	SCRATCH

CALLING SEQUENCE CALL PDEN(ICUTER)

SYMBOL	TYPE	DESCRIPTION
ICUTER	I	INPUT - CUTER ITERATION
SUBROUTINES USED		NONE
COMMON BLOCKS		CPAFAN TPEBLK
INPUT FILES		NONE
OUTPUT FILES		IOUT - PRINTER

SUBROUTINE PDEN2(BCENTR, AREA, DEN, ADJDEN, APSIG, SUM1, DENCON, SAVSIG) PDEN 55



REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

IMPLICIT REAL*8(A-H,O-Z)	PDEN	56
LOGICAL CMPGPR	PDEN	57
DIMENSION BCENTR(2,1), AREA(4,1), CON(1), ADJEN(1), APSIG(1), SUM1(1)	PDEN	58
DIMENSION DFNCON(NCCNST,1), SAVSIG(1), INCS(4)	PDEN	59
COMMON/CPARAM/NSTA, NMAST, NSTEST, NDIRM, MFIAS, NGPC1, NGPC2, NGPC04,	PDEN	60
• NCSFST, CMPGPR, LIM1, LIM2, NCON, NABLOCK, NTIDEN, INNRSW,	PDEN	61
• NCONST, NCCNS	PDEN	62
COMMON/TPRBLK/INTP, ICUT, ITAPES(10)	PDEN	63
INDXNO(I) = NDIRM*(I-1) - (I*(I-1))/2	PDEN	64
NCCNZ = NABLOCK - NCCNST	PDEN	65
INDS(1) = 1	PDEN	66
INDS(2) = NCON2	PDEN	67
INDS(3) = NABLOCK + 1	PDEN	68
INDS(4) = NCON	PDEN	69
LM = 2	PDEN	70
IF(NCON.GT.NABLOCK) LM = 4	PDEN	71
RETURN	PDEN	72
ENTRY PDEN(OUTER)	PDEN	73
I2 = NDIRM - 3 + NMAST - NABLOCK + NCCNST	PDEN	74
DO 100 I = 1, NABLOCK	PDEN	75
I2 = I2 + 1	PDEN	76
IF(MOD(I,45).EQ.1) WRITE(ICUT,1000) ICUT, I	PDEN	77
IF(MOD(I,5).EQ.1) WRITE(ICUT,1002)	PDEN	78
DEG = BCENTR(1,1)	PDEN	79
IDP = DEG	PDEN	80
SP = (DEG - IDP) * 60.000	PDEN	81
IMP = SP	PDEN	82
SP = (SP - IMP) * 60.00	PDEN	83
DEG = BCENTR(2,1)	PDEN	84
IDL = DEG	PDEN	85
SL = (DEG - IDL) * 60.000	PDEN	86
IML = SL	PDEN	87
SL = (SL - IML) * 60.00	PDEN	88
C SUM SUB-BLOCK APLAS AND COMPUTE ADJUSTED SIGMAS	PDEN	89
ASUM = AREA(1,1) + AREA(2,1) + AREA(3,1) + AREA(4,1)	PDEN	90
IF(I.GT.NCCNZ) GO TO 50	PDEN	91
I1 = INDXNO(I2) + I2	PDEN	92
ADJSIG = DSORT(SUM1(I1))	PDEN	93
SAVSIG(1) = 40JSIG	PDEN	94
GO TO 100	PDEN	95
50 SUM = 0.000	PDEN	96
SLMSG2 = 0.000	PDEN	97
I1 = 1 - NCCNZ	PDEN	98
DO 60 L = 1, LV, 2	PDEN	99
J1 = INDS(L)	PDEN	100
J2 = INDS(L+1)	PDEN	101
DO 60 J = J1, J2	PDEN	102
POCDA = DFNCON(I1, J)	PDEN	103
SLM = SUM + POCDA * ADJEN(J)	PDEN	104
IF(L.GT.1) GO TO 60	PDEN	105
SLMSG2 = SLMSG2 + (POCDA * SAVSIG(J)) ** 2	PDEN	106
60 CONTINUE	PDEN	107
ADJSIG = DSORT(SLMSG2)	PDEN	108
ADJTN(I) = SUM	PDEN	109
C PRINT DENSITIES	PDEN	110
100 WRITE(ICUT,1001) I, IDP, IMP, SP, IDL, IML, SL, /SUM, CON(I)	PDEN	111

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

	• ADJDEN(I), APSIG(I), ADJSIG	PDEN 112
	RETURN	PDEN 113
1000	FORMAT(1H1,40X,46HSURFACE DENSITIES ADJUSTED FOR OUTER ITERATION,	PDEN 114
	• I2/ 30X,15HCENTER OF BLOCK,14X,5HBLOCK,15X,5HVALUE,24X,	PDEN 115
	• 5HSIGMA/21X,35HGEOMETRIC LATITUDE EAST LONGITUDE,4X,	PDEN 116
	• 4HAREA,7X,2HA-PRICR1,7X,2HADJUST1,7X,2HA-PRICR2,4X,2HADJUST2/PDEN 117	
	• 13X,(HBLOCK *2(4X,13HDEC MI SECOND),4X,7H(KM**2),	PDEN 118
	• 3(6X,9H(KG/M**2)),3X,9H(KG/M**2))	PDEN 119
1001	FORMAT(12X,15,2X,2(17,13,F7.3),-CP12.0,1X,1P2015.6,1X,2E12.3)	PDEN 120
1002	FORMAT(1X)	PDEN 121
	END	PDEN 122

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR



NAME PDEN1
PURPOSE TO PRINT INPUT SURFACE DENSITIES
CALLING SEQUENCE CALL PDEN1
SUBROUTINES USED NONE
COMMON BLOCKS SRFBLK TPEBLK
INPUT FILES NCNE
OUTPUT FILES IOUT - PRINTER

```

SUBROUTINE PDEN1                                PDEN 18
  IMPLICIT REAL*8(A-H,O-Z)                      PDEN 19
  REAL DLAT,DLON,DEL,APSIG                      PDEN 20
  INTEGER*2 NLAT,NLON                          PDEN 21
  COMMON/SRFBLK/SLAT(675),SLON(675),DLAT(675),DLON(675),DEL(675),
  * APSIG(675),NLAT(675),NLON(675),NSRBLK      PDEN 22
  COMMON/TPEBLK/INTF,ICUT,ITAFES(10)          PDEN 23
  NLINES=0                                       PDEN 24
  IDEN2=0                                       PDEN 25
  DO 100 K=1,2                                   PDEN 26
  DO 100 I=1,NSRBLK                             PDEN 27
    IF (APSIG(I).LT.0.0) GO TO (100,20),K      PDEN 28
    IF (K.EQ.2) GO TO 100                       PDEN 29
  20 NLINES=NLINES+1                             PDEN 30
    IF (MOD(NLINES,45).EQ.1) WRITE (ICUT,1000) PDEN 31
    IF (MOD(NLINES,5).EQ.1) WRITE (IOUT,1003) PDEN 32
    IDEN1=IDEN2+1                               PDEN 33
    IDEN2=IDEN1+NLAT(I)+NLON(I)-1             PDEN 34
    ICP=SLAT(I)                                 PDEN 35
    ICL=SLON(I)                                 PDEN 36
    SP=(SLAT(I)-ICP)*60.00                     PDEN 37
    SL=(SLON(I)-ICL)*60.00                     PDEN 38
    IMP=SP                                       PDEN 39
    IML=SL                                       PDEN 40
    SP=(SP-IMP)*60.00                          PDEN 41
    SL=(SL-IML)*60.00                          PDEN 42
    EN=NLAT(I)+DLAT(I)                         PDEN 43
    EL=NLON(I)+DLON(I)                         PDEN 44
    BS=SLAT(I)-EN*0.500                       PDEN 45
    BE=SLON(I)-EL*0.500                       PDEN 46
    ES=BS+EN                                     PDEN 47
    EL=BE+EL                                     PDEN 48
    IF (K.EQ.1) WRITE (ICUT,1001) IDEN1, IDEN2, ICP, IMP, SP, ICL, IML, SL,
    * DLAT(I), DLON(I), NLAT(I), NLON(I), DEL(I), APSIG(I), EN, BS, SE, BW PDEN 49
    IF (K.EQ.2) WRITE (ICUT,1002) IDEN1, IDEN2, ICP, IMP, SP, ICL, IML, SL,
    * DLAT(I), DLON(I), NLAT(I), NLON(I), DEL(I), EN, ES, BE, BW PDEN 50
  100 CONTINUE                                  PDEN 51
  RETURN                                        PDEN 52

```

1000	FORMAT(1H1,53X,27HSURFACE DENSITY BLOCKS USED/	PDEN	55
	• 17X,22HCENTER OF MASTER BLOCK,7X,18HSIZE OF INCREMENTS.	PDEN	56
	• 3X,9HNUMBER OF,8X,15HSURFACE DENSITY.	PDEN	57
	• 8X,23HMASTER BLOCK BOUNCAPLES/	PDEN	58
	• 11X,25HGEODESIC LATITUDE EAST LONGITUDE,3X,3HLAT,6X,3HLON,	PDEN	59
	• 5X,1CHINCREMENTS,7X,5HVALUE,9X,5HSIGMA,4X,12HWORTH SOUTH.	PDEN	60
	• 3X,4HEAST,3X,4HWEST/	PDEN	61
	• 4X,6HEBLOCKS,4X,2(1X,14HDEG MN SECONDS,2X),2(2X,3HELG,4X).	PDEN	62
	• 2X,3HLAT,2X,3HLON,1X,2(5X,9H(KG/M**2)),4(3X,3HELG,1X))	PDEN	63
1001	FORMAT(2X,13,3H TC,14,2(2X,213,F7.3,1X),2(F8.2,1X),1X,215,1PE18.6,	PDEN	64
	• 512.3,CP4F7.1)	PDEN	65
1002	FORMAT(2X,13,3H TC,14,2(3X,213,F7.3,1X),2(F8.2,1X),1X,215,1PE18.6,	PDEN	66
	• 2X,1CHLNADJUSTED,CP4F7.1)	PDEN	67
1003	FORMAT(1X)	PDEN	68
	END	PDEN	69
		PDEN	70

PLHOUT

DESCRIPTION

Subroutine PLHOUT converts an input Earth-fixed Cartesian station position and its associated Cartesian covariance matrix to the spherical geodetic coordinate system.

The order of computation is:

- Compute the spheroid height, h , using an iterative procedure.
- Compute the geodetic latitude ϕ , and the east longitude λ .
- Compute the partial derivatives of ϕ , λ , and h with respect to the rectangular coordinates.
- Invoke subroutine VCONV to convert the input covariance matrix.

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

NAME PLHOUT

PURPOSE 1) TO COMPUTE GEODETIC PHI, LAMBDA, H FROM
 GEOCENTRIC X, Y, Z
 2) TO COMPUTE PARTIAL DERIVATIVES OF GEODETIC
 COORDINATES WITH RESPECT TO GEOCENTRIC
 COORDINATES

CALLING SEQUENCE CALL PLHOUT(STAP,XYZSIG,PLHSIG,PHI,LAMBDA,H)

SYMBOL	TYPE	DESCRIPTION
STAP (3)	DP	INPUT - GEOCENTRIC X,Y,Z
XYZSIG (3,3)	R	INPUT - SIGMAS ON X,Y,Z
PLHSIG (3,3)	R	OUTPUT - SIGMAS ON PHI, LAMBDA, H
PHI	DP	OUTPUT - GEODETIC LATITUDE
LAMBDA	DP	OUTPUT - GEODETIC EAST LONGITUDE
H	R	OUTPUT - GEODETIC HEIGHT

SUBROUTINES USED DARCTN VCCNV

COMMON BLOCKS INTBLK

INPUT FILES NONE

OUTPUT FILES NONE



REFERENCES 'GEOGYN SYSTEMS DESCRIPTION'
 VOLUME 1 - GEOGYN DOCUMENTATION

SUBROUTINE PLHOUT(STAP,XYZSIG,PLHSIG,PHI,LAMBDA,H)	PLHO	42
DIMENSION PLHSIG(3,3),XYZSIG(3,3),PARTL(3,3)	PLHO	43
REAL*8 PHI,LAMBDA,SINPHI,ZT,H1,XYSO,T,ESQ,ESQ1,T1,H2,ESQP,DARCTN,	PLHO	44
AE,STAP(3),SPSC,CCNH,PTXYSC	PLHO	45
COMMON/INTBLK/G1(8),AE,ALSC(2),FLAT,G2(118)	PLHO	46
REAL*8 FLAT	PLHO	47
DATA DELTA/.001/	PLHO	48
C CALCULATE CONSTANTS	PLHO	49
ESQ=FLAT	PLHO	50
ESQ1=(1.000-ESQ)**2	PLHO	51
ESQ=1.000-ESQ1	PLHO	52
T=ESQ*STAP(3)	PLHO	53
XYSO=STAP(1)**2+STAP(2)**2	PLHO	54
C ITERATIVE PROCEDURE FOR HEIGHT	PLHO	55

```

DO 10 J=1,25
Z1=STAP(3)+T
H1=SQRT(XYSQ+Z1**2)
SINPHI=Z1/H1
ESCSP=ESQ*SINPHI
H2=AE/ SQRT(1.000-ESCSP*SINPHI)
T1=H2*ESQ
IF(DABS(T1-T).LT.DELTA) GO TO 20
10 T=T1
C HEIGHT
20 H=H1-H2
C GEODETIC LATITUDE
RTXYSQ=DSQRT(XYSQ)
PHI=ATAN2(ZT,RTXYSQ)
C EAST LONGITUDE
LAMBDA=DAFC IN(STAP(2),STAP(1))
C CALCULATE CONSTANTS FOR PARTIALS
ZSQ=STAP(3)**2
SPSQ=SINPHI**2
COSPHI=SQRT(1.000-SPSQ)
CONPHI=ESQ/(ESQ**2*XYSQ+ZSQ)*RTXYSQ
CONH=1.000-TSQ*SPSQ
CONH=CONH*SQRT(CONH)
CONH=-ESQ*AE*ESQ*SINPHI*CONPHI/(CONH-STAP(3)*COSPHI/SPSQ)
C PARTIALS OF PHI WITH RESPECT TO X,Y,Z
PARTL(1,1)=-STAP(1)*STAP(3)*CONPHI
PARTL(2,1)=-STAP(2)*STAP(3)*CONPHI
PARTL(3,1)=XYSQ*CONPHI
C PARTIALS OF LAMBDA WITH RESPECT TO X,Y,Z
PARTL(1,2)=-STAP(2)/XYSQ
PARTL(2,2)=STAP(1)/XYSQ
PARTL(3,2)=0.
C PARTIALS OF H WITH RESPECT TO X,Y,Z
PARTL(1,3)=CONH*PARTL(1,1)
PARTL(2,3)=CONH*PARTL(2,1)
PARTL(3,3)=CONH*PARTL(3,1)+1.000/SINPHI
C CONVERT SIGMAS ON X,Y,Z TO SIGMAS ON PHI, LAMBDA, H
CALL VCCNV(XYZSIG,PLHSIG,PARTL)
RETURN
END
  
```

PLHO 56
 PLHO 57
 PLHO 58
 PLHO 59
 PLHO 60
 PLHO 61
 PLHO 62
 PLHO 63
 PLHO 64
 PLHO 65
 PLHO 66
 PLHO 67
 PLHO 68
 PLHO 69
 PLHO 70
 PLHO 71
 PLHO 72
 PLHO 73
 PLHO 74
 PLHO 75
 PLHO 76
 PLHO 77
 PLHO 78
 PLHO 79
 PLHO 80
 PLHO 81
 PLHO 82
 PLHO 83
 PLHO 84
 PLHO 85
 PLHO 86
 PLHO 87
 PLHO 88
 PLHO 89
 PLHO 90
 PLHO 91
 PLHO 92
 PLHO 93
 PLHO 94
 PLHO 95



NAME POLE

PURPOSE TO SELECT FROM A TABLE FOR A GIVEN INPUT DATE THE COORDINATES OF THE TRUE POLE

CALLING SEQUENCE. CALL POLE(XP,YP,DAY)

SYMBOL	TYPE	DESCRIPTION
XP	DP	OUTPUT - X COORDINATE OF THE POLE
YP	DP	OUTPUT - Y COORDINATE OF THE POLE
DAY	DP	INPUT - TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR FOR THE ARC

SUBROUTINES USED DJUL

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES *GEDDYN SYSTEMS DESCRIPTION*
VOLUME 1 - GEDDYN DOCUMENTATION

SUBROUTINE POLE(XP,YP,DAY)	POLE	31
DOUBLE PRECISION XP,YP,DAY,D,DJUL,DFIRST,DLAST	POLE	32
DIMENSION X(687),Y(687),X1(117),X2(180),X3(181),X4(209),Y1(117),	POLE	33
Y2(180),Y3(181),Y4(209)	POLE	34
EQUIVALENCE (X1(1),X(1)),(Y1(1),Y(1)),(X2(1),X(118)),(Y2(1),Y(118))	POLE	35
.(X3(1),X(298)),(Y3(1),Y(298)),(X4(1),X(479)),(Y4(1),Y(479))	POLE	36
C FIRST POINT IS ON 17 SEPT 1957, JULIAN DATE 2436099.5	POLE	37
C POSITIONS ARE RELATIVE TO THE MEAN POLE OF 1900 0.5	POLE	38
C THE INCREMENT BETWEEN POINTS IS 10 DAYS	POLE	39
DATA X1/	POLE	40
X0.340,0.296,0.248,0.199,0.151,0.103,0.047,-.015,-.085,-.140,-.178,POLE	POLE	41
X-.203,-.220,-.227,-.232,-.232,-.224,-.212,-.193,-.166,-.136,-.104,POLE	POLE	42
X-.071,-.036,0.002,0.042,0.082,0.119,0.153,0.194,0.215,0.245,0.273,POLE	POLE	43
X0.300,0.327,0.353,0.374,0.391,0.403,0.389,0.360,0.323,0.281,0.234,POLE	POLE	44
X0.186,0.133,0.080,0.027,-.026,-.058,-.102,-.124,-.143,-.157,-.168,POLE	POLE	45
X-.175,-.174,-.169,-.161,-.151,-.138,-.123,-.104,-.083,-.060,-.033,POLE	POLE	46
X-.002,0.033,0.074,0.117,0.169,0.221,0.266,0.299,0.318,0.323,0.323,POLE	POLE	47
X0.318,0.309,0.296,0.277,0.255,0.227,0.196,0.163,0.134,0.105,0.076,POLE	POLE	48
X0.048,0.020,-.008,-.024,-.055,-.077,-.096,-.095,-.101,-.101,-.100,POLE	POLE	49
X-.099,-.096,-.091,-.081,-.068,-.050,-.030,-.008,0.012,0.032,0.050,POLE	POLE	50
X0.069,0.086,0.102,0.118,0.132,0.145,0.149/	POLE	51
DATA X2/	POLE	52
X .148, .147, .147, .147, .142, .142, .141, .140, .138, .135, .131,POLE	POLE	53
X .127, .123, .118, .111, .101, .088, .075, .064, .055, .046, .037,POLE	POLE	54
X .028, .019, .011, .003,-.003,-.008,-.012,-.015,-.018,-.021,-.024,POLE	POLE	55

X-.027,-.029,-.031,-.033,-.031,-.025,-.015,-.010, .002, .014, .026, POLE	56
X .038, .050, .062, .076, .091, .105, .119, .131, .142, .152, .161, POLE	57
X .169, .176, .181, .183, .179, .171, .161, .150, .138, .124, .108, POLE	58
X .091, .073, .054, .032, .009,-.014,-.036,-.056,-.068,-.098,-.098, POLE	59
X-.112,-.112,-.108,-.100,-.088,-.073,-.055,-.032,-.004, .027, .058, POLE	60
X .088, .117, .143, .167, .189, .208, .225, .238, .246, .249, .247, POLE	61
X .240, .227, .207, .180, .147, .109, .057, .023,-.019,-.057,-.002, POLE	62
X-.122,-.146,-.164,-.223,-.247,-.262,-.272,-.273,-.265,-.248,-.228, POLE	63
X-.202,-.171,-.137,-.105,-.075,-.046,-.018, .012, .049, .099, .129, POLE	64
X .163, .199, .209, .224, .235, .240, .236, .229, .217, .202, .183, POLE	65
X .157, .123, .085, .049, .018,-.010,-.039,-.070,-.105,-.146,-.176, POLE	66
X-.197,-.209,-.217,-.225,-.228,-.227,-.219,-.204,-.186,-.165,-.142, POLE	67
X-.117,-.099,-.052,-.014, .024, .062, .099, .132, .159, .176, .185, POLE	68
X .192, .197, .200, .199/ POLE	69
DATA X3/ POLE	70
X .195, .196, .167, .147, .126, .102, .075, .049, .020,-.012,-.043, POLE	71
X-.070,-.092,-.107,-.122,-.138,-.154,-.169,-.175,-.173,-.155,-.153, POLE	72
X-.141,-.126,-.108,-.089,-.068,-.048,-.027,-.007, .013, .032, .049, POLE	73
X .065, .081, .098, .113, .124, .127, .125, .121, .112, .097, .081, POLE	74
X .067, .056, .046, .034, .022, .010, .002,-.010,-.030,-.027,-.027, POLE	75
X-.021,-.022,-.022,-.023,-.023,-.021,-.019,-.017,-.013,-.009,-.001, POLE	76
X .008, .004,-.003,-.007,-.009,-.011,-.015,-.021,-.022,-.015,-.015, POLE	77
X-.014,-.009,-.004,-.006,-.006,-.006,-.004, .000, .004, .007, .009, POLE	78
X .011, .014, .018, .022, .028, .037, .047, .055, .062, .068, .074, POLE	79
X (.79, .082, .080, .071, .059, .046, .037, .031, .024, .009,-.014, POLE	80
X-.037,-.058,-.101,-.134,-.146,-.149,-.149,-.157,-.119,-.098,-.078, POLE	81
X-.062,-.045,-.034,-.025,-.014,-.001, .014, .029, .046, .062, .080, POLE	82
X .099, .170, .134, .147, .152, .151, .137, .120, .098, .079, .065, POLE	83
X .050, .039, .029, .021, .004,-.020,-.044,-.074,-.103,-.129,-.151, POLE	84
X-.165,-.174,-.173,-.161,-.139,-.119,-.109,-.106,-.101,-.089,-.071, POLE	85
X-.051,-.029, .017, .055, .088, .119, .145, .168, .186, .199, .207, POLE	86
X .209, .205, .195, .191, .183/ POLE	87
DATA X4/ POLE	88
X .160, .120, .078, .049, .024, .000,-.029,-.057,-.085,-.111,-.133, POLE	89
X-.152,-.174,-.190,-.209,-.216,-.212,-.200,-.192,-.183,-.170,-.150, POLE	90
X-.129,-.104,-.061, .002, .060, .103, .140, .170, .198, .224, .245, POLE	91
X .253, .263, .264, .252, .229, .203, .179, .157, .135, .109, .083, POLE	92
X .055, .022,-.014,-.052,-.083,-.107,-.127/ POLE	93
DATA Y1/ POLE	94
Y0.057,0.029,0.001,-.025,-.045,-.058,-.068,-.075,-.077,-.073,-.061, POLE	95
Y-.044,-.012,0.029,0.074,0.127,0.194,0.239,0.284,0.330,0.372,0.403, POLE	96
Y0.424,0.441,0.457,0.467,0.473,0.471,0.463,0.451,0.430,0.398,0.365, POLE	97
Y0.325,0.281,0.251,0.219,0.167,0.133,0.082,0.047,-.002,-.030,-.065, POLE	98
Y-.076,-.093,-.082,-.079,-.071,-.060,-.047,-.030,-.007,0.019,0.051, POLE	99
Y0.083,0.119,0.157,0.193,0.229,0.262,0.293,0.321,0.347,0.367,0.381, POLE	100
Y0.384,0.391,0.376,0.366,0.349,0.325,0.286,0.271,0.251,0.233,0.215, POLE	101
Y0.197,0.179,0.159,0.137,0.115,0.095,0.076,0.070,0.056,0.044,0.034, POLE	102
Y0.023,0.028,0.033,0.044,0.059,0.079,0.098,0.116,0.133,0.149,0.165, POLE	103
Y0.179,0.192,0.204,0.216,0.229,0.242,0.257,0.271,0.283,0.293,0.301, POLE	104
Y0.305,0.305,0.303,0.299,0.294,0.284,0.271/ POLE	105
DATA Y2/ POLE	106
X .252, .235, .221, .209, .210, .199, .188, .177, .164, .149, .134, POLE	107
X .121, .110, .101, .090, .099, .109, .104, .112, .118, .128, .140, POLE	108
X .154, .170, .187, .203, .207, .210, .213, .215, .217, .219, .220, POLE	109
X .221, .222, .222, .223, .225, .230, .238, .258, .267, .273, .277, POLE	110
X .281, .295, .299, .289, .287, .283, .276, .267, .257, .245, .231, POLE	111

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

X	.217.	.202.	.185.	.165.	.142.	.121.	.105.	.093.	.083.	.075.	.070.	POLE	112
X	.067.	.068.	.071.	.073.	.090.	.106.	.127.	.153.	.181.	.207.	.231.	POLE	113
X	.259.	.279.	.29	.307.	.318.	.328.	.337.	.344.	.349.	.351.	.350.	POLE	114
X	.344.					.333.	.317.	.297.	.272.	.247.	.222.	POLE	115
X	.197.	.172.	.147.	.122.	.096.	.070.	.044.	.020.	.001.	.017.	.025.	POLE	116
X	.025.	.017.	.003.	.016.	.040.	.069.	.101.	.134.	.162.	.190.	.220.	POLE	117
X	.253.	.239.	.325.	.359.	.370.	.417.	.443.	.465.	.470.	.482.	.478.	POLE	118
X	.472.	.467.	.460.	.445.	.424.	.394.	.350.	.320.	.301.	.270.	.235.	POLE	119
X	.199.	.155.	.136.	.111.	.092.	.074.	.058.	.040.	.052.	.054.	.055.	POLE	120
X	.057.	.062.	.075.	.094.	.120.	.150.	.180.	.211.	.245.	.291.	.315.	POLE	121
X	.349.	.333.	.410.	.425.	.441.	.454.	.465.	.458.	.460.	.448.	.425.	POLE	122
X	.401.	.377.	.350.	.323.	.299.	.275.	.249.	.222/				POLE	123
												POLE	124
												POLE	125
DATA Y3/													
Y	.192.	.153.	.137.	.117.	.101.	.087.	.073.	.061.	.054.	.052.	.054.	POLE	125
Y	.063.	.074.	.087.	.103.	.124.	.149.	.177.	.202.	.223.	.243.	.263.	POLE	126
Y	.295.	.304.	.322.	.340.	.356.	.361.	.361.	.361.	.358.	.354.	.350.	POLE	127
Y	.346.	.341.	.332.	.316.	.294.	.273.	.254.	.235.	.220.	.206.	.190.	POLE	128
Y	.175.	.169.	.167.	.165.	.164.	.164.	.166.	.169.	.173.	.177.	.183.	POLE	129
Y	.193.	.200.	.202.	.204.	.206.	.210.	.216.	.220.	.222.	.216.	.210.	POLE	130
Y	.210.	.207.	.203.	.204.	.207.	.209.	.212.	.219.	.232.	.245.	.253.	POLE	131
Y	.254.	.265.	.270.	.272.	.274.	.276.	.273.	.269.	.263.	.261.	.258.	POLE	132
Y	.254.	.254.	.256.	.258.	.250.	.236.	.220.	.208.	.198.	.188.	.180.	POLE	133
Y	.174.	.160.	.165.	.162.	.160.	.158.	.151.	.145.	.141.	.133.	.130.	POLE	134
Y	.130.	.139.	.156.	.186.	.215.	.234.	.251.	.262.	.267.	.307.	.316.	POLE	135
Y	.323.	.337.	.350.	.360.	.368.	.372.	.373.	.371.	.365.	.357.	.344.	POLE	136
Y	.332.	.317.	.300.	.273.	.241.	.205.	.173.	.147.	.125.	.115.	.111.	POLE	137
Y	.103.	.106.	.104.	.102.	.101.	.106.	.114.	.125.	.135.	.146.	.175.	POLE	138
Y	.210.	.201.	.200.	.202.	.200.	.201.	.200.	.200.	.210.	.221.	.234.	POLE	139
Y	.444.	.448.	.448.	.442.	.426.	.402.	.373.	.342.	.309.	.275.	.244.	POLE	140
Y	.213.	.183.	.155.	.125.	.101/							POLE	141
												POLE	142
DATA Y4/													
Y	.078.	.063.	.053.	.038.	.023.	.013.	.013.	.022.	.035.	.057.	.085.	POLE	143
Y	.117.	.148.	.179.	.211.	.253.	.301.	.340.	.370.	.397.	.424.	.452.	POLE	144
Y	.480.	.505.	.519.	.524.	.520.	.500.	.476.	.450.	.421.	.387.	.351.	POLE	145
Y	.317.	.293.	.249.	.211.	.175.	.139.	.118.	.103.	.092.	.078.	.062.	POLE	146
Y	.048.	.035.	.026.	.027.	.034.	.050.	.074/					POLE	147
												POLE	148
C FIRST POINT 570917 *** LAST POINT 711204												POLE	148
DATA DFIRST/2436099.500/												POLE	149
CLAST/2441379.500/												POLE	150
LOGICAL FRSTME/.TRUE./												POLE	151
IF(.NOT.FRSTME) GO TO 20												POLE	152
FRSTME = .FALSE.												POLE	153
IN=(CLAST-DFIRST)/10.00+1.00												POLE	154
DO 10 I=1,IN												POLE	155
X(I)=X(I)*0.4848137E-5												POLE	156
Y(I)=Y(I)*0.4848137E-5												POLE	157
10 CONTINUE												POLE	158
20 CONTINUE												POLE	159
D = DJUL(DAY)												POLE	160
IF(D .GE. DFIRST) GO TO 30												POLE	161
XP = X(1)												POLE	162
YP = Y(1)												POLE	163
RETURN												POLE	164
30 CONTINUE												POLE	165
IF(D .LT. CLAST) GO TO 40												POLE	166
XP = X(IN)												POLE	167

COPIED BY THE
ORIGINAL PAGE IS POOR

```
YP = Y(IN)
RETURN
40 CONTINUE
D=(D-DFIRST)/10.CD0+1.0D0
INTERPLATE FOR COORDINATES OF THE POLE
ID = D
IDP1 = ID + 1
D1=D-DFLOAT(IDP1)
D2=D-DFLOAT(ID)
YP = -D1*Y(ID) + D2*Y(IDP1)
XP = -D1*X(ID) + D2*X(IDP1)
RETURN
END
```

```
POLE 168
POLE 169
POLE 170
POLE 171
POLE 172
POLE 173
POLE 174
POLE 175
POLE 176
POLE 177
POLE 178
POLE 179
POLE 180
```

Circular D69

1 - UNIVERSAL TIME AND COORDINATES OF THE POLE

Date (Oh UT) 1972	J.D. 2400000.5 +	smoothed values				raw values			UT1 -10.
		x 0'001	y 0'000	UT2-UTC 0.0001s	UT1-UTC 0.0001s	x 0'001	y 0'001	UT1-UTC 0.0001s	
June 1	41 469	-145	+356	-5252	-5553	-151	+341	-5541	
6	474	-134	+366	-5416	-5710	-129	+371	-5743	
11	479	-120	+376	-5579	-5861	-114	+359	-5872	
16	484	-105	+385	-5740	-6006	-92	+374	-6019	
21	489	-89	+394	-5899	-6145	-92	+421	-6165	
26	494	-72	+402	-6057	-6279	-51	+382	-6263	
July 1	499	-54	+409	+3786	+3591	-58	+407	+3616	

IAT-UTC is exactly 10s in June 1972
IAT-UTC is exactly 11s since 1972 July 1st, Oh UTC.

2 - EMISSION TIME OF TIME SIGNALS, for June 1972 (E = UTC-Signal in 0.0001s)

Signal	E	Signal	E	Signal	E
CHU	0	FTM42, FTK77, FTM87	0	NSS (o.c.)	+ 9
DAM, DAN, DAO	0	HBG	0	OLBS	(2)
DCF77	0	IAM	0	OMA	(2)
DGI	0	IBF	+ 3	PPE	- 5
DIZ	0	JJY	0	RWM (1)	0
FFH	0	LOL	- 5	VNG	0
FTA91	0	MSF	+ 1	WWV, WWVB, WWVH	0
		GBZ (3)	- 3	ZUO	(2)

(1) and other signals from USSR (2) no data available
(3) corrected values : April 1972, E = - 3 ; May 1972, E = - 2

3 - COORDINATED UNIVERSAL TIME (approximation UTC(i) of UTC, kept by the laboratory Ref. CCIR Recommendation 458, 1970)

a - From LORAN-C and Television pulses receptions

Date 1972 J.D. 2400000.5 +	June 11 41 479	June 21 41 489	July 1 41 499	
Laboratory i	UTC-UTC(i)		(unit : 1 µs)	
PTB (Braunschweig)	+ 2.9	+ 3.0	+ 2.9	
USNO (Washington)	- 6.6	- 6.5	- 6.2	
GP (Paris)	+ 1.6	+ 1.6	+ 1.6	
NBS (Boulder)	- 2.4	- 2.5	- 2.7	
RCO (Herstmonceux)	+ 3.2	+ 3.8	+ 4.4	
NRC (Ottawa)	+ 0.8	+ 0.9	+ 1.2	
FOA (Stockholm)	+ 23.9	+ 26.3	+ 28.6	
DHI (Hamburg)	- 16.9	- 15.3	- 13.9	
CH (Geneve)	+ 20.6	+ 20.7	+ 20.6	P. T.

NAME POSVEL

PURPOSE TO CONVERT OSCULATING ORBITAL ELEMENTS TO INERTIAL POSITION AND VELOCITY VECTORS

CALLING SEQUENCE. POSVEL(XYZ,AEI,IDRAD)

SYMBOL	TYPE	DESCRIPTION
XYZ (6)	DP	OUTPUT - CARTESIAN ELEMENTS
AEI (6)	DP	INPUT - KEPLER ELEMENTS
IDRAD	I	INPUT - =2 MEANS INPUT IN RADIANS =1 MEANS INPUT IN DEGREES

SUBROUTINES USED NONE

COMMON BLOCKS CONSTS INTRLK



INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES *GEDDYN SYSTEMS DESCRIPTION*
VOLUME 1 - GEDDYN DOCUMENTATION

SUBROUTINE POSVEL(XYZ,AEI,IDRAD)	POSV	33
IMPLICIT REAL*8 (A-H,C-Z)	POSV	34
REAL RMSTOT	POSV	35
DOUBLE PRECISION INCL,MEAN,NODE	POSV	36
DIMENSION AEI(6),XYZ(6),AEINPM(6),XYZXYZ(6)	POSV	37
COMMON/CONSTS/PI,TWOPI,RAD,RSEC	POSV	38
COMMON/INTRLK/THODTS(3),GM,AE(62)	POSV	39
EQUIVALENCE (A,AEINPM(1)),(E,AEINPM(2)),(INCL,AEINPM(3)),	POSV	40
(NODE,AEINPM(4)),(P,AEINPM(5)),(MEAN,AEINPM(6)),	POSV	41
(X,XYZXYZ(1)),(Y,XYZXYZ(2)),(Z,XYZXYZ(3)),	POSV	42
(XDOT,XYZXYZ(4)),(YDOT,XYZXYZ(5)),(ZDOT,XYZXYZ(6))	POSV	43
C DEFINE CONVERSION CRITERIA	POSV	44
DATA DELTA/0.1D-10/	POSV	45
SCALE=1.0D0	POSV	46
IF(IDRAD.EQ.1) SCALE=RAD	POSV	47
DO 10 I=1,6	POSV	48
AEINPM(I)=AEI(I)	POSV	49
IF(I.LE.2) GO TO 10	POSV	50
AEINPM(I)=AEI(I)*SCALE	POSV	51
10 CONTINUE	POSV	52
SOVUA=DSQRT(GM/DABS(A))*3	POSV	53
E2=E**2	POSV	54
ONEE2=DSQRT(DABS(1.0D0-E2))	POSV	55

C SINES AND COSINES OF THE ELEMENTS	POSV 56
COSI=DCOS(INCL)	POSV 57
SINI=DSIN(INCL)	POSV 58
SINK=DSIN(NODE)	POSV 59
COSN=DCOS(NODE)	POSV 60
SINP=DSIN(P)	POSV 61
COSP=DCOS(P)	POSV 62
C SET ECC. ANOM. EQUAL TO MEAN ANOM. FOR FIRST APRDX.	POSV 63
ECC=MEAN	POSV 64
C ITERATE	POSV 65
IF (E2.GE.1.000) GO TO 150	POSV 66
C ...FOR ELLIPTIC ORBITS	POSV 67
DO 100 J=1,50	POSV 68
E00=ECC	POSV 69
SINECC=DSIN(E00)	POSV 70
COSECC=DCOS(E00)	POSV 71
ECOS=1.000-E*COSECC	POSV 72
ECC=E00-(E00-E*SINECC-MEAN)/ECOS	POSV 73
IF (DABS(E00-ECC).LT.DELTA) GO TO 200	POSV 74
100 CONTINUE	POSV 75
PRINT 1000	POSV 76
GO TO 200	POSV 77
C ...FOR HYPERBOLIC ORBITS	POSV 78
150 DO 160 J=1,100	POSV 79
E00=ECC	POSV 80
SINECC=DSINH(E00)	POSV 81
COSECC=DCOSH(E00)	POSV 82
ECOS=E*COSECC-1.000	POSV 83
ECC=E00-(E*SINECC-E00-MEAN)/ECOS	POSV 84
IF (DABS(E00-ECC).LT.DELTA) GO TO 200	POSV 85
160 CONTINUE	POSV 86
PRINT 1000	POSV 87
200 SPCN=SINP*COSN	POSV 88
CPSN=COSP*SINN	POSV 89
CPCN=COSP*COSN	POSV 90
SPSN=SINP*SINN	POSV 91
A2=DABS(A)*ONEME2	POSV 92
AX=A*(CPCN-SPSN*COSI)	POSV 93
AY=A*(SPCN+CPSN*COSI)	POSV 94
AZ=A*SINP*SINI	POSV 95
BX=-AZ*(SPCN+CPSN*COSI)	POSV 96
BY=A2*(CPCN*COSI-SPSN)	POSV 97
BZ=A2*COSP*SINI	POSV 98
C=COSECC-E	POSV 99
EDOT=SOMUA/ECOS	POSV 100
C ...FOR X,Y,Z	POSV 101
X=AX*C+BX*SINECC	POSV 102
Y=AY*C+BY*SINECC	POSV 103
Z=AZ*C+BZ*SINECC	POSV 104
C ...FOR XDOT,YDOT,ZDOT	POSV 105
IF (E2.GE.1.000) SINECC=-SINECC	POSV 106
XDOT=EDOT*(BX*COSECC-AX*SINECC)	POSV 107
YDOT=EDOT*(BY*COSECC-AY*SINECC)	POSV 108
ZDOT=EDOT*(BZ*COSECC-AZ*SINECC)	POSV 109
DO 300 I=1,6	POSV 110
300 XYZ(I)=XYZYZ(I)	POSV 111
RETURN	POSV 112
1000 FORMAT(1H1,3HECCENTRIC ANOMALY NOT CONVERGED)	POSV 113
END	POSV 114



PRECES

DESCRIPTION

Subroutine PRECES generates the rotation matrix to precess a vector from the mean equator and equinox of an input date to the mean equator and equinox of 1950.0.

The precession angles are evaluated using polynomials derived by Simon Newcomb. The rotation matrices are evaluated by ROTMAT; the output rotation matrix is computed as the product of the three input matrices by MULMAT.

NAME PRECES

PURPOSE TO GENERATE THE MATRIX FOR PRECESSION FROM MEAN EQUATOR AND EQUINOX OF AN EPOCH TO MEAN EQUATOR AND EQUINOX OF 1950

CALLING SEQUENCE CALL PRECES(DAY,X)

SYMBOL	TYPE	DESCRIPTION
DAY	DP	INPUT - TIME IN DAYS FROM JAN 0.0 OF REFERENCE YEAR
X (3,3)	DP	OUTPUT - PRECESSION MATRIX

SUBROUTINES USED ROTMAT MULMAT YMDAY

COMMON BLOCKS INITBK

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES 'GEOGYN SYSTEMS DESCRIPTION'
 VOLUME 1 - GEOGYN DOCUMENTATION

```

SUBROUTINE PRECES(DAY,X)
REAL*8 DAY, DBASE, D, X(3,3), Z(3,3,3), ANGLE, YMDAY
REAL*8 COEF(3,3) / +.30595327465D-6, +.3972049D-14, +.191031D-20,
    -.26603999754D-6, +.1548118D-14, +.4139C2D-20,
    +.30595320465D-6, +.1077492D-14, +.178097D-20/
INTEGER AXIS(3) / 3, 2, 3/
COMMON/INITBK/IG1(53), NOTIST, IG2(3)
LOGICAL NOTIST
IF(NOTIST) GO TO 10
NOTIST=.TRUE.
DBASE=YMDAY(500100.0, 0.0, CD0) - .07200
10 D=DAY-DBASE
DO 30 I=1,3
  ANGLE=0.00
  DO 20 J=1,3
    ANGLE=(ANGLE+COEF(4-J,I))*D
  20 CALL ROTMAT(ANGLE, AXIS(I), Z(1,1,I))
  30 CALL MULMAT(X, Z(1,1,3), Z(1,1,2), Z(1,1,1))
RETURN
END
  
```

PREC 31
 PREC 32
 PREC 33
 PREC 34
 PREC 35
 PREC 36
 PREC 37
 PREC 38
 PREC 39
 PREC 40
 PREC 41
 PREC 42
 PREC 43
 PREC 44
 PREC 45
 PREC 46
 PREC 47
 PREC 48
 PREC 49
 PREC 50

PREDCT

DESCRIPTION

Subroutine PREDCT computes the residuals and partial derivatives for observations involving fewer than two tracking stations for the parameter estimation. In addition, PREDCT predicts the measurement values from a priori data and the geodetic spherical coordinates of the satellite.

The observation types by program index number are:

- 1) right ascension and declination
- 2) range (including sat-sat summed range)
- 3) range-rate (including sat-sat summed range rate)
- 4) altimeter height and height rate
- 5) l and m direction cosines
- 6) X and Y angles
- 7) azimuth and elevation

The order of computation is as follows:

- Call ORBIT to obtain the satellite position, velocity, and variational partials.
- Call GRHRAN to obtain the right ascension of Greenwich and the station-satellite vector for observing station.
- Compute the equivalent for each measurement and the associated residual.
- Compute the Earth-fixed geometric partial derivatives of the calculated equivalents.

- Convert the partial derivatives to inertial coordinates and chain them back to epoch,
- If necessary compute the spheroid height, geodetic latitude, and east longitude of the satellite.

NAME PREDCT
ENTRY POINT PURPOSE
PREDCI INITIAL IZATION
PREDCT TO COMPUTE MEASUREMENTS, RESIDUALS (O-C), AND MEASUREMENT PARTIALS
CALLING SEQUENCE CALL PREDCI(EHAT,NHAT,ZHAT,PMPX0,PPPX0,NPARG,NEONMX)

SYMBOL	TYPE	DESCRIPTION
EHAT (3,1)	DP	INPUT - STATION UNIT EAST VECTOR
NHAT (3,1)	DP	INPUT - STATION UNIT NORTH VECTOR
ZHAT (3,1)	DP	INPUT - STATION UNIT VERTICAL VECTOR
PMPX0 (NPARG,1)	DP	OUTPUT - MEASUREMENT PARTIALS
PPPX0 (6,NEONMX,2)	DP	INPUT - SATELLITE STATE PARTIAL WRT EPOCH PARAMETERS
NPARG	I	INPUT - MAXIMUM NUMBER OF PARAMETERS PER MEASUREMENTS
NEONMX	I	INPUT - NPARG+8

CALLING SEQUENCE CALL PREDCT(ISTA,DAY,RESID1,RESID2,DATASW)

SYMBOL	TYPE	DESCRIPTION
ISTA	I	INPUT - STATION INDEX
DAY	DP	INPUT & OUTPUT - MEASUREMENT TIME
RESID1	R	OUTPUT - FIRST MEASUREMENT RESIDUAL (O-C)
RESID2	R	OUTPUT - SECOND MEASUREMENT RESIDUAL
DATASW	L	INPUT - .TRUE. WHEN POSITION OF SATELLITE WANTED .FALSE. WHEN MEASUREMENT PARTIALS WANTED

SUBROUTINES USED ORBIT GRHRAN PROCES GLEM CLFAR
DOTPRD DARCTN XINERT XEFIX YINERT
YEFIX

COMMON BLOCKS APARAM CONSTS CORR1 CUVECT GNDRK
INITRK INTOLK PREPLK XYZOUT

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES *GEDDYN SYSTEMS DESCRIPTION*
VOLUME 1 - GEDDYN DOCUMENTATION

```

SUBROUTINE PREDCT(EHAT,NHAT,ZHAT,PMPXO,PXPXO,NPARM,NEQNMX)      PRED 67
  IMPLICIT REAL*8 (A-H,C-Z)                                       PRED 68
  LOGICAL*1 VHFCHN,PREPRO,NOT1ST,TWOSTA                          PRED 69
  LOGICAL NOEST,SATSW,SUBSAT,DATASW,TRKSW,SATSAT,ELEVSW        PRED 70
  INTEGER*2 MTYPE,NMEAS,PRETYP,CHANNEL,ISAT                     PRED 71
  INTEGER RECNO,ADDR                                             PRED 72
  DOUBLE PRECISION NHAT,LOVE                                     PRED 73
  DIMENSION PMSTA1(3),PMSTA2(3),ORSC(2),XYZDOT(3),PM(6,4),G(3),  PRED 74
  • G2(3,3),R2(3,3),PARH2(3,3),PMSTA3(3,2),PMPXC(NPARM,1),    PRED 75
  • PXPX(6,NEQNMX,2),PMSTA(3,2),EHAT(3,1),NHAT(3,1),ZHAT(3,1), PRED 76
  • URHOO(3),CR(3),HI(3),HI(3),AEIXYZ(6,6),P12(3),PS2(3),PS1(3), PRED 77
  • PS3(3),P23(3),V12(3),VS2(3),VS1(3),VS3(3),V23(3)         PRED 78
  COMMON/APARAM/NPAR,INPAR1(3),NSAT,NGPARC(5)                   PRED 79
  COMMON/CONSTS/DPI,DTWOPI,DRAD,DRSEC                           PRED 80
  COMMON/CORB1/T,W,U,THETG,PERHT(2),APHT(2),PRD(2)              PRED 81
  COMMON/CVJECT/UNAT(3,2),XYZ(3,2),RXYZ(3,2),RNMV(3,2),P(2),PSO(2), PRED 82
  • XYSO(2)                                                       PRED 83
  COMMON/GNDRK/SATLAT(2),SATLON(2),SATH(2),FLEV(2),SATSW        PRED 84
  COMMON/INITBK/IG1(42),SUBSAT,IG2(5),MISLOG(9)                 PRED 85
  COMMON/INTBLK/THDOT1(2),THDOT2S,GM,AE,AFSO,FLAT,FSQ32,FFSQ32, PRED 86
  • GM3(49),NEONS(2),ADDR(2,3),LOVE(4)                           PRED 87
  COMMON/PREBLK/DAYSTA,ORSO(2),SIG(2),SRFNDX,ISN,MTYPE,NMEAS,   PRED 88
  • ISAT,PRETYP,CHANNEL,VHFCHN,PREPRO,RECNO                      PRED 89
  COMMON/XYZOUT/XYZI(6,4)                                        PRED 90
  DATA NOT1ST/.FALSE./                                          PRED 91
  DATA C/2.99792508/,DTDL/1.GD-9/                               PRED 92
  EQUIVALENCE (F,FLAT),(PMSTA1(1),PMSTA(1,1)),(PMSTA2(1),PMSTA(1,2)) PRED 93
  EQUIVALENCE (PRD(1),TRKSW),(AEIXYZ(1,1),URHOO(1)),           PRED 94
  • (AEIXYZ(1,2),CR(1)),(AEIXYZ(1,3),HI(1)),(AEIXYZ(1,4),HI(1)) PRED 95
  EQUIVALENCE (DTRANS,ORSO(2)),(KKSAT,SIG(2)),(TIME2,SRFNDX)   PRED 96
  RETURN                                                         PRED 97
  ENTRY PREDCT(ISTA,DAY,RESID1,RESID2,DATASW)                   PRED 98
  IF(NOT1ST) GO TO 10                                           PRED 99
C INITIALIZE                                                    PRED 100
  C1=1.5D0*AE*F*F                                               PRED 101
  C2=AE*F+C1                                                    PRED 102
  FLAT21=(1.0D0-FLAT)**2                                        PRED 103
  NOT1ST=.TRUE.                                                PRED 104
10 CONTINUE                                                    PRED 105
  SATSAT=(.NOT.DATASW).AND.(MTYPE.EQ.2.OR.MTYPE.EQ.3).AND.KKSAT.GT. PRED 106
  • .AND.KKSAT.LE.NSAT                                         PRED 107
  ISAT1=ISAT                                                    PRED 108
  ISAT2=ISAT                                                    PRED 109
  INCR=1                                                         PRED 110
  K2=0                                                           PRED 111

```

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

ELEVSW=PREPRO	PRED 112
C IF SAT-SAT TRACKING THEN GO TO 300	PRED 113
IF(SATSAT) GO TO 300	PRED 114
TWOSTA=MTYPE.GT.26.AND..NOT.DATASW	PRED 115
NOEST=.FALSE.	PRED 116
C FOR VLBI & AVERAGE RANGE RATE DATA SKIP INTEGRATOR CALL	PRED 117
IF(TWOSTA) GO TO 50	PRED 118
C OBTAIN SATELLITE ORBIT	PRED 119
CALL ORBIT(DAY)	PRED 120
C OBTAIN R.A. GREENWICH & STATION - SATELLITE VECTORS	PRED 121
THETG=GRHRAN(DAY,ISTA)	PRED 122
C SKIP MEASUREMENTS & PARTIALS IF ONLY GROUND TRACK REQUESTED	PRED 123
IF(DATASW) GO TO 2020	PRED 124
IF(ISTA.EQ.0) GO TO 200	PRED 125
NOEST=NPAR.EQ.0.OR.(SIG(1).EQ.0.ODC.AND.SIG(2).EQ.0.CDD)	PRED 126
50 NP=3	PRED 127
C COMPUTE ELEVATION	PRED 128
ENSG=1.000-RENV(3,ISAT)**2	PRED 129
EN=DSQRT(ENSG)	PRED 130
ELEV(ISAT)=DATAN(RENV(3,ISAT)/EN)	PRED 131
C IF VLBI OR AVERAGE RANGE RATE THEN GO TO 230	PRED 132
IF(TWOSTA) GO TO 230	PRED 133
C PROCESS DATA IF REQUESTED	PRED 134
IF(PREPRO) CALL PROCES(ISTA,DAY,THETG)	PRED 135
C IF TRANSIT TIME CORRECTION WAS MADE THEN RECOMPUTE ELEVATION	PRED 136
IF(.NOT.ELEVSW) GO TO 190	PRED 137
ENSG=1.000-RENV(3,ISAT)**2	PRED 138
EN=DSQRT(ENSG)	PRED 139
ELFV(ISAT)=DATAN(RENV(3,ISAT)/EN)	PRED 140
190 GO TO (400,450,500,550,600,700,800),MTYPE	PRED 141
C PCE DATA	PRED 142
200 DO 210 I=1,6	PRED 143
210 PM(I,1)=0.000	PRED 144
MT=MOD(MTYPE-15,6)+1	PRED 145
PM(MT,1)=1.000	PRED 146
RESID1=OBSO(1)-XYZI(MT,ISAT)	PRED 147
ELEV(ISAT)=0.500*DPI	PRED 148
NP=6	PRED 149
IF(MTYPE.LT.21) GO TO 1900	PRED 150
C CALL ELEM FOR KEPLER DATA	PRED 151
CALL ELEM(XYZI(1,ISAT),PMSTA1,3,.FALSE.,AEIXYZ)	PRED 152
RESID1=OBSO(1)-PMSTA1(MT)	PRED 153
IF(DABS(RESID1).GT.DPI) RESID1=RESID1-DSIGN(DTWOPI,RESID1)	PRED 154
DO 220 I=1,6	PRED 155
220 PM(I,1)=AEIXYZ(MT,I)	PRED 156
GO TO 1900	PRED 157
230 KTYPE=MTYPE-26	PRED 158
GO TO (240,500,450,450),KTYPE	PRED 159
C TIME DELAY	PRED 160
240 RESID1=R(ISAT)/C	PRED 161
DO 250 I=1,3	PRED 162
250 PMSTA1(I)=-UPAT(I,ISAT)/C	PRED 163
GO TO 1900	PRED 164
C IF SAT-SAT DATA THEN COMPUTE UPLINK & DOWNLINK TRANSIT TIMES	PRED 165
300 NOEST=NPAR.EQ.0.OR.SIG(1).EQ.0.CDD	PRED 166
NP=3	PRED 167

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

J=1SAT	PRED 168
ISAT1=MIN0(J,KKSAT)	PRED 169
ISAT2=MAX0(J,KKSAT)	PRED 170
301 IPRE=0	PRED 171
IF(.NOT.PREPRO) GO TO 380	PRED 172
IPRE=PRETYP/10	PRED 173
PRETYP=PRETYP-IPRE*10	PRED 174
IF(IPRE.GT.0) GO TO 310	PRED 175
CALL ORBIT(DAY)	PRED 176
THETG=GRHRAN(DAY,ISTA)	PRED 177
DO 302 I=1,3	PRED 178
302 PS2(I)=RXYZ(I,ISAT)	PRED 179
KK=0	PRED 180
DAYP=DAY	PRED 181
304 R12=0.000	PRED 182
DO 306 I=1,3	PRED 183
P12(I)=PS2(I)-RXYZ(I,KKSAT)	PRED 184
306 R12=R12+P12(I)**2	PRED 185
R12=DSQRT(R12)	PRED 186
DTRANS=DAY+R12/(C*8.6404)	PRED 187
DT=DABS(DTRANS-DAYP)	PRED 188
IF(DT.LE.DTOL) GO TO 380	PRED 189
KK=KK+1	PRED 190
IF(KK.GT.5) GO TO 308	PRED 191
DAYP=DTRANS	PRED 192
CALL ORBIT(DTRANS)	PRED 193
THETG=GRHRAN(DAY,ISTA)	PRED 194
GO TO 304	PRED 195
308 DT=DT*8.6404	PRED 196
PRINT 3000,DT,DTOL	PRED 197
380 KK=0	PRED 198
IF(.ELEVSW) TIME2=DAY	PRED 199
382 DAYP=TIME2	PRED 200
CALL ORBIT(TIME2)	PRED 201
THETG=GRHRAN(DAY,ISTA)	PRED 202
DO 384 I=1,3	PRED 203
384 PS3(I)=RXYZ(I,KKSAT)	PRED 204
IF(.NOT.ELEVSW) GO TO 390	PRED 205
R23=0.000	PRED 206
DO 386 I=1,3	PRED 207
P23(I)=PS3(I)-PS2(I)	PRED 208
386 R23=R23+P23(I)**2	PRED 209
R23=DSQRT(R23)	PRED 210
TIME2=DAY-R23/(C*8.6404)	PRED 211
DT=DABS(TIME2-DAYP)	PRED 212
IF(DT.LE.DTOL) GO TO 390	PRED 213
KK=KK+1	PRED 214
IF(KK.LE.5) GO TO 382	PRED 215
DT=DT*8.6404	PRED 216
PRINT 3000,DT,DTOL	PRED 217
390 IF(MTYPE.NE.3) GO TO 310	PRED 218
VS3(1)=XFFIX(XYZI(4,KKSAT),XYZI(5,KKSAT))+THDT25*XY7(2,KKSAT)	PRED 219
VS3(2)=YFFIX(XYZI(4,KKSAT),XYZI(5,KKSAT))-THDT25*XY7(1,KKSAT)	PRED 220
VS3(3)=XYZI(6,KKSAT)	PRED 221
310 KK=0	PRED 222
318 DAYP=DTRANS	PRED 223

REPRODUCIBILITY OF THIS
ORIGINAL PAGE IS POOR

CALL ORBIT(DTRANS)	PRED 224
THETG=GRHRAN(DAY,ISTA)	PRED 225
RS1=0.000	PRED 226
DO 320 I=1,3	PRED 227
PS1(I)=RXYZ(I,KKSAT)	PRED 228
320 RS1=RS1+PS1(I)**2	PRED 229
RS1=DSORT(RS1)	PRED 230
IF(IPRE.LE.0) GO TO 324	PRED 231
KK=KK+1	PRED 232
IF(KK.GT.5) GO TO 322	PRED 233
DTRANS=DAY-RS1/(C*8.64D4)	PRED 234
DT=DABS(DAYP-DTRANS)	PRED 235
IF(DT.GT.DTOL) GO TO 318	PRED 236
GO TO 323	PRED 237
322 DT=DT*8.64D4	PRED 238
PRINT 3000,DT,DTOL	PRED 239
323 DAY=DTRANS	PRED 240
324 ENS0=1.000-RENV(3,KKSAT)**2	PRED 241
EN=DSORT(ENS0)	PRED 242
ELEV(ISAT)=ATAN(RENV(3,KKSAT)/EN)	PRED 243
ELEV(KKSAT)=ELEV(ISAT)	PRED 244
PREPRO=PREPRC.AND.PRETYP.GT.0	PRED 245
IF(.NOT.PREPRO) GO TO 325	PRED 245
J=ISAT	PRED 247
ISAT=KKSAT	PRED 248
CALL PROCES(ISTA,DTRANS,THETG)	PRED 249
THETG=GRHRAN(DAY,ISTA)	PRED 250
ISAT=J	PRED 251
325 IF(MTYPE.NE.3) GO TO 326	PRED 252
NP=6	PRED 253
VS1(1)=XFIX(XYZI(4,KKSAT),XYZI(5,KKSAT))+THDT2S*XYZ(2,KKSAT)	PRED 254
VS1(2)=YFIX(XYZI(4,KKSAT),XYZI(5,KKSAT))-THDT2S*XYZ(1,KKSAT)	PRED 255
VS1(3)=XYZI(6,KKSAT)	PRED 256
326 KK=0	PRED 257
328 DAYP=DAY	PRED 258
CALL ORBIT(DAY)	PRED 259
THETG=GRHRAN(DAY,ISTA)	PRED 260
R12=0.000	PRED 261
DO 330 I=1,3	PRED 262
PS2(I)=RXYZ(I,ISAT)	PRED 263
P12(I)=PS2(I)-PS1(I)	PRED 264
330 R12=R12+P12(I)**2	PRED 265
R12=DSORT(R12)	PRED 266
IF(IPRE.LE.0) GO TO 334	PRED 267
KK=KK+1	PRED 268
IF(KK.GT.5) GO TO 332	PRED 269
DAY=DTRANS-R12/(C*8.64D4)	PRED 270
DT=CABS(DAYP-DAY)	PRED 271
IF(DT.GT.DTOL) GO TO 328	PRED 272
GO TO 331	PRED 273
332 DT=DT*8.64D4	PRED 274
PRINT 3000,DT,DTOL	PRED 275
GO TO 331	PRED 276
334 RS1I=1.000/RS1	PRED 277
R12I=1.000/R12	PRED 278
IF(MTYPE.NE.3) GO TO 338	PRED 279

VS2(1)=XEFIX(XYZI(4, ISAT), XYZI(5, ISAT))+THDT2S*XYZ(2, ISAT)	PRED 280
VS2(2)=YEFIX(XYZI(4, ISAT), XYZI(5, ISAT))-THDT2S*XYZ(1, ISAT)	PRED 281
VS2(3)=XYZI(6, ISAT)	PPED 282
DO 336 I=1,3	PPFD 283
V12(I)=(VS2(I)-VS1(I))*R12I	PPED 284
336 VS1(I)=VS1(I)*RS1I	PPFD 285
RRS12=DOTPRD(P12,V12)+DOTPRD(PS1,VS1)	PRED 286
338 DO 340 I=1,3	PPED 287
P12(I)=P12(I)*R12I	PPFD 288
340 PS1(I)=PS1(I)*RS1I	PPFD 289
IF(NPAR4.GT.0) CALL CLEAR(PMPXC,NPARM,4)	PRED 290
KK=1	PPFD 291
JJ=3	PPFD 292
IF(ISAT.EQ.ISAT1) GO TO 345	PPED 293
KK=3	PPED 294
JJ=1	PRED 295
345 R23=0.000	PRED 296
RS3=0.000	PPED 297
DO 351 I=1,3	PPFD 298
P23(I)=PS3(I)-PS2(I)	PPFD 299
R23=R23+P23(I)**2	PRED 300
351 RS3=RS3+PS3(I)**2	PRED 301
R23=DSORT(R23)	PPED 302
RS3=DSORT(RS3)	PRED 303
IF(MTYPE-2) 350,350,360	PPED 304
C COMPUTE RESIDUAL & PARTIALS FOR SUMMED RANGE	PRED 305
350 RESID1=ORSO(1)-0.500*(RS1+R12+R23+RS3)	PPED 306
IF(NDEST) GO TO 3010	PPED 307
DO 352 I=1,3	PPED 308
352 PMSTA1(I)=-PS1(I)	PPFD 309
PM(1, KK)=XINERT(P12(1), P12(2))	PRED 310
PM(2, KK)=YINERT(P12(1), P12(2))	PRED 311
PM(1, JJ)=XINERT(PS1(1), PS1(2))-PM(1, KK)	PRED 312
PM(2, JJ)=YINERT(PS1(1), PS1(2))-PM(2, KK)	PRED 313
PM(3, KK)=P12(3)	PRED 314
PM(3, JJ)=PS1(3)-P12(3)	PPED 315
GO TO 1930	PPED 316
C COMPUTE RESIDUAL & PARTIALS FOR SUMMED RANGE RATE	PRED 317
360 RS3I=1.000/RS3	PPFD 318
R23I=1.000/R23	PRED 319
DO 361 I=1,3	PPED 320
V23(I)=(VS3(I)-VS2(I))*R23I	PRED 321
361 VS3(I)=VS3(I)*RS3I	PPED 322
RESID1=ORSO(1)-0.500*(RRS12+DOTPRD(P23,V23)+DOTPRD(PS3,VS3I))	PRED 323
IF(NDEST) GO TO 2010	PPED 324
DP=DOTPRD(PS1,VS1)	PRED 325
DO 362 I=1,3	PPED 326
362 PMSTA1(I)=-VS1(I)+DP*PS1(I)	PPFD 327
PM(4, KK)=XINERT(P12(1), P12(2))	PPFD 328
PM(5, KK)=YINERT(P12(1), P12(2))	PRED 329
PM(6, KK)=P12(3)	PPED 330
PM(4, JJ)=XINERT(PS1(1), PS1(2))-PM(4, KK)	PPED 331
PM(5, JJ)=YINERT(PS1(1), PS1(2))-PM(5, KK)	PPFD 332
PM(6, JJ)=PS1(3)-P12(3)	PPFD 333
DP=DOTPRD(P12,V12)	PPFD 334
DO 364 I=1,3	PPED 335

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

364	PMSTA2(1)=V12(1)-DP*P12(1)	PRED 336
	PM(1, KK)=XINERT(PMSTA2(1), PMSTA2(2))	PRED 337
	PM(2, KK)=YINERT(PMSTA2(1), PMSTA2(2))	PRED 338
	PM(3, KK)=PMSTA2(3)	PRED 339
	PM(1, JJ)=-XINERT(PMSTA1(1), PMSTA1(2))-PM(1, KK)	PPED 340
	PM(2, JJ)=YINERT(PMSTA1(1), PMSTA1(2))-PM(2, KK)	PRED 341
	PM(3, JJ)=-PMSTA1(3)-PMSTA2(3)	PRED 342
	PM(1, KK)=PM(1, KK)-THDT2S*PM(5, KK)	PRED 343
	PM(2, KK)=PM(2, KK)+THDT2S*PM(4, KK)	PPED 344
	PM(1, JJ)=PM(1, JJ)-THDT2S*PM(5, JJ)	PRED 345
	PM(2, JJ)=PM(2, JJ)+THDT2S*PM(4, JJ)	PPED 346
	GO TO 1930	PRED 347
C RIGHT ASCENSION AND DECLINATION		
400	XY=DSQRT(XYSQ(ISAT))	PRED 348
	OBSC(1)=ATAN2(UHAT(2, ISAT), UHAT(1, ISAT))+THETG	PRED 349
	OBSC(1)=OBSC(1)-DMOD(OBSC(1), DTWOP1)	PPED 350
	IF(DABS(OBSC(1)).GT.CPI) OBSC(1)=OBSC(1)-DSIGN(DTWOP1, OBSC(1))	PRED 351
	RESID1=OBSC(1)	PPED 352
	OBSC(2)=OBSC(2)-ATAN(RXYZ(3, ISAT)/XY)	PPED 353
	IF(DABS(OBSC(2)).GT.OPI) OBSC(2)=OBSC(2)-DSIGN(DTWOP1, OBSC(2))	PPED 354
	RESID2=OBSC(2)	PPED 355
	IF(.NOT.TRKSW) GO TO 425	PRED 356
	URHO(1)=DCOS(OBSC(1)-THETG)*DCOS(OBSC(2))	PPED 357
	URHO(2)=DSIN(OBSC(1)-THETG)*DCOS(OBSC(2))	PPED 358
	URHO(3)=DSIN(OBSC(2))	PRED 359
	XYZDOT(1)=XEFIX(XYZI(4, ISAT), XYZI(5, ISAT))+THDT2S*XYZ(2, ISAT)	PPED 360
	XYZDOT(2)=YEFIX(XYZI(4, ISAT), XYZI(5, ISAT))-THDT2S*XYZ(1, ISAT)	PRED 361
	XYZDOT(3)=XYZI(6, ISAT)	PPED 362
	DOTPS=DOTPRD(UHAT(1, ISAT), XYZ(1, ISAT))/DOTPRD(URHO, XYZ(1, ISAT))	PRED 363
	DO 405 I=1,3	PRED 364
	DR(I)=R(ISAT)*DOTPS*URHO(I)-RXYZ(I, ISAT)	PPED 365
	I2=MOD(I, 3)+1	PRED 366
	I3=MOD(I2, 3)+1	PPED 367
	H(I)=XYZ(I2, ISAT)*XYZDOT(I3)-XYZ(I3, ISAT)*XYZDOT(I2)	PRED 368
405	HI(1)=XYZI(12, ISAT)*XYZI(13+3, ISAT)-XYZI(13, ISAT)*XYZI(12+3, ISAT)	PPED 369
	T=DOTPRD(DR, XYZDOT)	PRED 370
	W=DCTPRD(DR, H)	PRED 371
	SINU=- (XYZI(2, ISAT)*HI(2)+XYZI(1, ISAT)*HI(1))*HI(3)+XYZI(3, ISAT)*	PPED 372
	(HI(1)**2+HI(2)**2)	PRED 373
	COSU=(-XYZI(1, ISAT)*HI(2)+XYZI(2, ISAT)*HI(1))*DSQRT(HI(1)**2+	PRED 374
	HI(2)**2+HI(3)**2)	PRED 375
	U=DARCTN(SINU, COSU)	PPED 376
425	IF(NDEST) GO TO 2010	PRED 377
C CALCULATE PARTIALS FOR ESTIMATION		
	RSQXY=RXYZ(3, ISAT)/(RSQ(ISAT)*YY)	PRED 378
	PMSTA1(1)=RXYZ(2, ISAT)/XYSQ(ISAT)	PPED 379
	PMSTA1(2)=-RXYZ(1, ISAT)/XYSQ(ISAT)	PRED 380
	PMSTA1(3)=0.000	PPED 381
	PMSTA2(1)=RXYZ(1, ISAT)+RSQXY	PRED 382
	PMSTA2(2)=RXYZ(2, ISAT)*RSQXY	PPED 383
	PMSTA2(3)=-XY/RSQ(ISAT)	PRED 384
	GO TO 1930	PPED 385
C RANGE		
450	RESID1=OBSC(1)-R(ISAT)	PRED 386
	IF(NDEST) GO TO 2010	PPED 387
	DO 455 I=1,3	PRED 388
		PPED 389
		PRED 390
		PPED 391

```

455 PMSTA1(I)=-UHAT(I, ISAT)
      GO TO 1900
C RANGE RATE
500 XYZDOT(1)=XEFIX(XYZI(4, ISAT), XYZI(5, ISAT))+THDT2S*XYZ(2, ISAT)
      XYZDOT(2)=YEFIX(XYZI(4, ISAT), XYZI(5, ISAT))-THDT2S*XYZ(1, ISAT)
      XYZDOT(3)=XYZI(6, ISAT)
      OBSC(1)=DOTPRD(XYZDOT, UHAT(1, ISAT))
      RESID1=OBSO(1)-OBSC(1)
      IF(NDEST) GO TO 2010
      NP=6
      DO 505 I=1,3
      PMSTA1(I)=- (XYZDOT(I)-OBSC(I)*UHAT(I, ISAT))/R(ISAT)
505 PMSTA3(I,1)=-UHAT(I, ISAT)
      IF(.NOT.TWGSTA) GO TO 1900
      RESID1=OBSO(1)*SRFNDX/C
      DO 520 I=1,3
      PMSTA1(I)=PMSTA1(I)*SRFNDX/C
520 PMSTA3(I,1)=PMSTA3(I,1)*SRFNDX/C
      GO TO 1900
550 CONTINUE
C ALTIMETER MEASUREMENTS
      NP=6
      ELFV(ISAT)=0.500*DPI
      C3=(2.000*C2-4.000*C1*UHAT(3, ISAT)**2)*UHAT(3, ISAT)
      RR=1.000/R(ISAT)
      OBSC(1)=R(ISAT)-AE-(C1*UHAT(3, ISAT)**2-C2)*UHAT(3, ISAT)**2
      RESID1=OBSO(1)-OBSC(1)
      DO 552 I=1,3
552 G(I)=-UHAT(3, ISAT)*UHAT(I, ISAT)*RR
      G(3)=G(3)+RR
C ALTIMETER PARTIALS
      DO 555 J=1,3
555 PMSTA1(J)=- (UHAT(J, ISAT)+C3+G(J))
      XYZDOT(1)=XEFIX(XYZI(4, ISAT), XYZI(5, ISAT))+THDT2S*XYZ(2, ISAT)
      XYZDOT(2)=YEFIX(XYZI(4, ISAT), XYZI(5, ISAT))-THDT2S*XYZ(1, ISAT)
      XYZDOT(3)=XYZI(6, ISAT)
C ALTIMETER RATE
      OBSC(2)=-DOTPRD(PMSTA1, XYZDOT)
      RESID2=OBSO(2)-OBSC(2)
      IF(NDEST) GO TO 2010
      C4=-RR*RR
      DO 558 J=1,3
558 G2(J, J)=(1.000-3.000*UHAT(J, ISAT)**2)*UHAT(3, ISAT)+C4
      G2(3,3)=G2(3,3)+2.000*UHAT(3, ISAT)*C4
      G2(1,2)=-3.000*UHAT(1, ISAT)*UHAT(2, ISAT)*UHAT(3, ISAT)+C4
      G2(2,1)=G2(1,2)
      DO 6667 I=1,2
6667 G2(I,3)=(1.000-3.000*UHAT(3, ISAT)**2)*UHAT(I, ISAT)+C4
      G2(3,1)=G2(1,3)
      DO 553 I=1,3
      DO 554 J=1,3
554 R2(I, J)=-UHAT(I, ISAT)*UHAT(J, ISAT)*RR
557 R2(1,1)=R2(1,1)+5R
      C5=-12.000*C1*UHAT(3, ISAT)**2+2.000*C2
      DO 556 I=1,3
      FMSTA3(I,1)=C.C0

```

```

PRED 392
PRED 393
PRFD 394
PRFD 395
PRFD 396
PRFD 397
PRED 398
PRFD 399
PRED 400
PRFD 401
PRFD 402
PRED 403
PRED 404
PRED 405
PRED 406
PRED 407
PRED 408
PRED 409
PRED 410
PRED 411
PRED 412
PRED 413
PRED 414
PRFD 415
PRED 416
PRED 417
PRED 418
PRED 419
PRED 420
PRED 421
PRED 422
PRED 423
PRED 424
PRED 425
PRED 426
PRED 427
PRFD 428
PRED 429
PRED 430
PRED 431
PRED 432
PRFD 433
PRED 434
PRED 435
PRED 436
PRFD 437
PRFD 438
PRED 439
PRED 440
PRED 441
PRED 442
PRFD 443
PRED 444
PRFD 445
PRED 446
PRED 447

```



REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR



DO 556 J=1,3	PPED 448
C SECOND PARTIALS OF ALTIMETER WRT X,Y,Z	PRED 449
556 PARH2(I,J)=R2(I,J)+C3*G2(I,J)+G(I)*C5*G(J)	PRED 450
DO 557 I=1,3	PRED 451
C PARTIAL HDOT,WRT X,Y,Z,XDOT,YDOT,ZDOT	PRED 452
PMSTA2(I)=0.00	PPED 453
PMSTA3(I,2)=PMSTA1(I)	PRED 454
DO 557 J=1,3	PRED 455
557 PMSTA2(I)=PMSTA2(I)-XYZDOT(J)*PARH2(I,J)	PPED 456
GO TO 1930	PRED 457
C DIRECTION COSINES	PRED 458
600 RESID1=OBSO(1)-RENV(1,ISAT)	PPED 459
RESID2=OBSO(2)-RENV(2,ISAT)	PRED 460
IF(NDEST) GO TO 2010	PRED 461
DO 605 I=1,3	PRED 462
PMSTA1(I)=-((HAT(I,ISTA)-RENV(1,ISAT)*UHAT(I,ISAT))/R(ISAT))	PPED 463
605 PMSTA2(I)=-((NHAT(I,ISTA)-RENV(2,ISAT)*UHAT(I,ISAT))/R(ISAT))	PRED 464
GO TO 1900	PPED 465
C X-Y ANGLES	PPED 466
700 EZSQ=1.000-RENV(2,ISAT)**2	PPED 467
EZ=DSCR(EZSQ)	PRED 468
RESID1=OBSO(1)-DATAN(RENV(1,ISAT)/RENV(3,ISAT))	PPED 469
RESID2=OBSO(2)-DATAN(RENV(2,ISAT)/EZ)	PRED 470
IF(NDEST) GO TO 2010	PRED 471
REZ=R(ISAT)*EZ	PRED 472
REZSQ=R(ISAT)*EZSQ	PRED 473
DO 705 I=1,3	PRED 474
PMSTA1(I)=(ZHAT(I,ISTA)+RENV(1,ISAT)-EHAT(I,ISTA)+RENV(3,ISAT))/REZSQ	PPED 475
REZSQ	PPED 476
705 PMSTA2(I)=-((NHAT(I,ISTA)-RENV(2,ISAT)*UHAT(I,ISAT))/REZ)	PRED 477
GO TO 1900	PRED 478
C AZIMUTH & ELEVATION ANGLES	PRED 479
800 RESID1=OBSO(1)-DARCTN(RENV(1,ISAT),RENV(2,ISAT))	PRED 480
IF(DABS(RESID1).GT.DP(1)) RESID1=RESID1-DSIGN(DTWOPI,RESID1)	PPED 481
RESID2=OBSO(2)-ELEV(ISAT)	PRED 482
IF(NDEST) GO TO 2010	PRED 483
REN=R(ISAT)*EN	PPED 484
RENSQ=R(ISAT)*EN SQ	PRED 485
DO 805 I=1,3	PRED 486
PMSTA1(I)=-((RENV(2,ISAT)*EHAT(I,ISTA)-RENV(1,ISAT)*NHAT(I,ISTA))/RENSQ)	PPED 487
RENSQ	PRED 488
805 PMSTA2(I)=-((ZHAT(I,ISTA)-RENV(3,ISAT)*UHAT(I,ISAT))/REN)	PRED 489
C CONVERT EARTH FIXED PARTIALS TO INERTIAL	PPED 490
1900 CALL CLEAR(PMPX0,NPARM,2*NMEAS)	PRED 491
IF(MTYPE.GT.14.AND.MTYOF.LT.27) GO TO 1940	PRED 492
DO 1925 K=1,NMEAS	PPED 493
PM(1,K)=-XINERT(PMSTA(1,K),PMSTA(2,K))	PRED 494
PM(2,K)=-YINERT(PMSTA(1,K),PMSTA(2,K))	PPED 495
PM(3,K)=-PMSTA(3,K)	PRED 496
PMPX0(NPARM-5,K)=PMSTA(1,K)	PPED 497
PMPX0(NPARM-4,K)=PMSTA(2,K)	PRED 498
PMPX0(NPARM-3,K)=PMSTA(3,K)	PPED 499
PMPX0(NPARM-7,K)=1.000	PRED 500
IF(NP.LT.4) GO TO 1925	PPED 501
PM(4,K)=-XINERT(PMSTA3(1,K),PMSTA3(2,K))	PRED 502
PM(5,K)=-YINERT(PMSTA3(1,K),PMSTA3(2,K))	PPED 503



PM(6,K)=-PMSTA3(3,K)	PRED 504
IF(MTYPE.NE.3.AND.MTYPE.NE.4.AND.MTYPE.NE.28) GO TO 1925	PRED 505
IF(MTYPE.EQ.4.AND.K.EQ.1) GO TO 1925	PRED 506
PM(1,K)=PM(1,K)-THDT25*PM(5,K)	PRED 507
PM(2,K)=PM(2,K)+THDT25*PM(4,K)	PRED 508
1925 CONTINUE	PRED 509
GO TO 1940	PRED 510
1930 INCR=ISAT2-ISAT1	PRED 511
PMPX0(NPARM-5,1)=PMSTA1(1)	PRED 512
FMPX0(NPARM-4,1)=PMSTA1(2)	PRED 513
PMPX0(NPARM-3,1)=PMSTA1(3)	PRED 514
PMPX0(NPARM-7,1)=1.000	PRED 515
C CHAIN INSTANTANEOUS PARTIALS BACK TO EPOCH	PRED 516
1940 ISATNO=ISAT1	PRED 517
1945 L1=(ISATNO-1)*6	PRED 518
DO 1950 I=1,6	PRED 519
L1=L1+I	PRED 520
DO 1950 K=1,NMEAS	PRED 521
K1=K+K2	PRED 522
DO 1950 J=1,NP	PRED 523
1950 PMPX0(L1,K)=FMPX0(L1,K)+PM(J,K1)*PXPX0(J,I,ISATNO)	PRED 524
I1=6	PRED 525
J1=NSAT*5	PRED 526
DO 1970 I=1,3	PRED 527
L1=ADDR(ISATNO,I)	PRED 528
IF(L1.LE.0) GO TO 1970	PRED 529
L1=J1*L1	PRED 530
I1=I1+1	PRED 531
DO 1960 K=1,NMEAS	PRED 532
K1=K+K2	PRED 533
DO 1960 J=1,NP	PRED 534
1960 PMPX0(L1,K)=PMPX0(L1,K)+PM(J,K1)*PXPX0(J,I1,ISATNO)	PRED 535
1970 CONTINUE	PRED 536
I1=I1+1	PRED 537
I2=NECNS(ISATNO)-1	PRED 538
IF(I1.GT.I2) GO TO 2000	PRED 539
L1=MAX0(ADDR(1,1),ADDR(2,1),ADDR(1,2),ADDR(2,2),ADDR(1,3),	PRED 540
ADDR(2,3))+J1	PRED 541
DO 1980 I=I1,I2	PRED 542
L1=L1+1	PRED 543
DO 1980 K=1,NMEAS	PRED 544
K1=K+K2	PRED 545
DO 1980 J=1,NP	PRED 546
1980 PMPX0(L1,K)=FMPX0(L1,K)+PM(J,K1)*PXPX0(J,I,ISATNO)	PRED 547
2000 K2=K2+2	PRED 548
ISATNO=ISATNO+INCR	PRED 549
IF(ISATNO.EQ.ISAT2) GO TO 1945	PRED 550
2010 ELEV(ISAT)=ELEV(ISAT)/DRAD	PRED 551
C TEST FOR GROUND TRACK WANTED	PRED 552
IF(.NOT.SUESAT) RETURN	PRED 553
C EAST LONGITUDE OF SATELLITE IN DEGREES	PRED 554
2020 DO 2050 J=1,NSAT	PRED 555
SATLON(J)=(ATAN2(XYZI(2,J),XYZI(1,J))-THETS)/DRAD	PRED 556
SATLON(J)=OMOD(SATLON(J)+7.202,3.602)	PRED 557
C GEODETIC LONGITUDE OF SATELLITE IN DEGREES	PRED 558
XYSO(J)=XYZI(1,J)**2+XYZI(2,J)**2	PRED 559



```
SATLAT(J)=XYZI(3,J)/(DSQRT(XYSQ(J))*FLAT21)
SATLAT(J)=DATAN(SATLAT(J))/DRAD
C SATELLITE HEIGHT IN METERS
RSAT=DSQRT(XYSQ(J)+XYZI(3,J)**2)
SPSISO=(XYZI(3,J)/RSAT)**2
2050 SATH(J)=(RSAT-AE)-(FSQ32*SPSISO**2-FFSQ32*SPSISO)
RETURN
3000 FORMAT(' ***** ACCEPTED TRANSIT TIME ERROR AFTER SIX '
• ' ITERATIONS =',E12.5,' SECONDS. GREATER THAN',E12.5,
• ' DAYS *****')
END
```



PRED 560
PRED 561
PRED 562
PRED 563
PRED 564
PRED 565
PRED 566
PRED 567
PRED 568
PRED 569
PRED 570























NAME PRNTPR

PURPOSE TO PRINT THE OBSERVATION PREPROCESSING REQUESTED

CALLING SEQUENCE CALL PRNTPR(OUTP,ATYPE)

SYMBCL	TYPE	DESCRIPTION
OUTP	I	OUTPUT - PRINTER
ATYPE	DP	OUTPUT - MEASUREMENT TYPE NAMES
(31)		

SUBROUTINES USED NONE



COMMON BLOCKS CGEOS

INPUT FILES NONE

OUTPUT FILES OUTP - PRINTER

SUBROUTINE PRNTPR(OUTP,ATYPE)	PRNT	25
REAL*8 ATYPE(31),NAME1,NAME2,ALL,TYPES,BLANK	PRNT	26
INTEGER OUTP	PRNT	27
DIMENSION DUM(2)	PRNT	28
COMMON/CGEOS/DUM,IPREPR(4,50),RF INDX(2,50),INDPRE(2,50),NCPRR,	PRNT	29
G1(202)	PRNT	30
INTEGER*2 IPREPR,INDPRE	PRNT	31
DATA MAXTPE/31/,ALL/5H ALL/,TYPES/5HTYPES/,BLANK/1H /	PRNT	32
C PRINT HEADING	PRNT	33
WRITE(OUTP,10700)	PRNT	34
DO 320 I=1,NCPRR	PRNT	35
WRITE(OUTP,10710)	PRNT	36
L=INDPRE(2,I)	PRNT	37
IF(L.EQ.0.OR.(L.GT.14.AND.L.LT.27)) L=MAXTPE	PRNT	38
C PRINT STATION NUMBER & MEASUREMENT TYPE	PRNT	39
NAME1=ALL	PRNT	40
NAME2=TYPES	PRNT	41
IF(L.NE.MAXTPE) NAME1=ATYPE(L)	PRNT	42
IF(L.NE.MAXTPE) NAME2=BLANK	PRNT	43
IF(L.LT.8) NAME2=ATYPE(L+7)	PRNT	44
IF(INDPRE(1,I)) 320,250,225	PRNT	45
225 WRITE(OUTP,10701) INDPRE(1,I),NAME1,NAME2	PRNT	46
GO TO 275	PRNT	47
250 WRITE(OUTP,10702) NAME1,NAME2	PRNT	48
275 CONTINUE	PRNT	49
DO 305 J=1,4	PRNT	50
IF(IPREPR(J,I).GT.0) GO TO (207,308,309,310),J	PRNT	51
305 CONTINUE	PRNT	52
GO TO 320	PRNT	53
C TRANSIT TIME	PRNT	54
307 WRITE(OUTP,10703)	PRNT	55

GO TO 335	PRNT	56
308 IF(IPREPR(1,1).NE.0) WRITE(OUTP,10710)	PRNT	57
C REFRACTION	PRNT	58
A=RFINDX(1,1)	PRNT	59
IF(A.LE.0.) A=328.5	PRNT	60
WRITE(OUTP,10704) A	PRNT	61
GO TO 335	PRNT	62
C CONSTANT TIMING	PRNT	63
309 IF(IPREPR(1,1)+IPREPR(2,1).NE.0) WRITE(OUTP,10710)	PRNT	64
WRITE(OUTP,10705) RFINDX(2,1)	PRNT	65
GO TO 335	PRNT	66
C OPTICAL	PRNT	67
310 WRITE(OUTP,10708)	PRNT	68
320 CONTINUE	PRNT	69
RETURN	PRNT	70
10700 FORMAT(1H1,50X,25HPREPROCESSING CORRECTIONS/1H0,44X,7HSTATION,	PRNT	71
1 21X,10HCORRECTION/1H ,36X,6HNUMBER,5X,7HTYPE(S),12X,	PRNT	72
2 7HTYPE(S),11X,5HVALUE)	PRNT	73
10701 FORMAT(37X,15,3X,A6,2X,A6)	PRNT	74
10702 FOFMAT(32X,3HALL,4X,A6,2X,A6)	PRNT	75
10703 FOFMAT(1H+,63X,12HTRANSIT TIME)	PRNT	76
10704 FOFMAT(1H+,63X,10HREFRACTION,F12.1,3H N UNITS)	PRNT	77
10705 FOFMAT(1H+,63X,11HTIMING BIAS,3PF11.1,13H MILLISECONDS)	PRNT	78
10708 FOFMAT(1H ,63X,7HOPTICAL)	PRNT	79
10710 FOFMAT(1H)	PRNT	80
END	PRNT	81



PROCES

DESCRIPTION

PROCES is a subroutine designed specifically to complete the GEODYN preprocessing.

PROCES makes the following preprocessing corrections:

- Transit Time,
- Annual Aberration,
- Diurnal Aberration,
- Parallax Refraction,
- Tropospheric Refraction,
- Range Transponder Corrections.

The specific corrections applied by PROCES are dependent upon indicators input to PROCES through COMMON storage.

NAME PROCES
 ENTRY PCINT PURPOSE
 PROCS1 INITIALIZATION
 PROCES TO COMPLETE PREPROCESSING OF SATELLITE MEASUREMENTS
 CALLING SEQUENCE CALL PROCS1(ISTANO,STAXYZ,RLAT,RLON)

SYMBOL	TYPE	DESCRIPTION
ISTANO (1)	I*2	INPUT - TRACKING STATION NUMBERS
STAXYZ (3,1)	DP	INPUT - TRACKING STATION CARTESIAN COORDINATES
RLAT (1)	DP	INPUT - TRACKING STATION LATITUDES
RLON (1)	DP	INPUT - TRACKING STATION LONGITUDES

CALLING SEQUENCE CALL PROCES(ISTA,DAY,THETG)

SYMBOL	TYPE	DESCRIPTION
ISTA	I	INPUT - TRACKING STATION INDEX
DAY	DP	INPUT & OUTPUT - MEASUREMENT TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR
THETG	DP	INPUT & OUTPUT - RIGHT ASCENSION OF GREENWICH

SUBROUTINES USED	ORBIT	GRHRAN	REFION	DOTPRO	NUMBRA
	ORBIT	GRHRAN	REFION	DOTPRO	NUMBRA
	DBSDOT	EPHEM	ERN	DJUL	

COMMON BLOCKS	CGEOS	CUVECT	CEPHEM	GNDTRK	CONSTS
	CGEOS	CUVECT	CEPHEM	GNDTRK	CONSTS
	PREBLK				

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES *GEDDYN SYSTEMS DESCRIPTION*
 VOLUME 1 - GEODYN DOCUMENTATION

SUBROUTINE PROCS1(ISTANO,STAXYZ,RLAT,RLON)
 IMPLICIT REAL*8 (A-H,C-Z)
 LOGICAL*1 VHFCHN,PREPRO
 LOGICAL SATSW

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

INTEGER RECNO	PRNC 56
INTEGER*2 MTYPE,NMEAS,PRETYP,CHANEL,ISAT1,IPRE,ISTAND	PRNC 57
DOUBLE PRECISION L	PRNC 58
REAL TRANSP,REFION,EON,CBSOBT	PRNC 59
DIMENSION PSAT(3),STASAT(3),ISTAND(1),TRANSP(3,2,9),KSATNO(27),	PRNC 60
STXYZ(3,1),RLAT(1),RLON(1),GATE(3),PRETYP(2),STPSAT(3)	PRNC 61
COMMON/CGEOS/ISATID(2),IPREPR(453)	PRNC 62
COMMON/CJVECT/UHAT(3,2),XYZ(3,2),RXYZ(3,2),PENV(3,2),R(2),RSQ(2),	PRNC 63
XYSO(2)	PRNC 64
COMMON/CEPHEM/AO(4),SUNX,SUNY,IG2(1324)	PRNC 65
COMMON/GNDRK/SATLAT(2),SATLON(2),SATH(2),ELEV(2),SATSW	PRNC 66
COMMON/CONSTS/DPI,DTWOP1,D2R,S2R	PRNC 67
COMMON/PREBLK/CAYSTA,OBSD1,OBSD2,SIG1,SIG2,SRFNDX,ISN,MTYPE,	PRNC 68
NMEAS,ISAT1,IPRE,CHANEL,VHFCHN,PREPRO,RECNO	PRNC 69
DATA NSATNO/27/,GATE/18737031,300,4684257,800,935851,600/,	PRNC 70
VLIGHT/2,99792509/,STDNDX/328,SDO/,PCONST/C,84323350-2/	PRNC 71
DATA KSATNO/65891,650891,6508901,	PRNC 72
68021,680021,6800201,	PRNC 73
64541,640541,6405401,	PRNC 74
65811,650811,6508101,	PRNC 75
66491,660491,6604901,	PRNC 76
67731,670731,6707301,	PRNC 77
68141,680141,6801401,	PRNC 78
69511,690511,6905101,	PRNC 79
65911,65911,690511/	PRNC 80
DATA TRANSP/	PRNC 81
C GEOS A	PRNC 82
• 0.3577E4,2*0.,0.306E4,2*0.,	PRNC 83
C GEOS B	PRNC 84
• 0.3648E4,2*0.,0.3684E4,2*0.,	PRNC 85
C OGO 1	PRNC 86
• 0.3636E4,2*0.,0.3710E4,2*0.,	PRNC 87
C OGO 2	PRNC 88
• 0.352E4,2*0.,0.373E4,2*0.,	PRNC 89
C OGO 3	PRNC 90
• 0.3519E4,0.8508E-2,0.2376E-6,0.3731E4,-.5597E-2,0.3535E-6,	PRNC 91
C OGO 4	PRNC 92
• 0.3599E4,0.6014E-2,0.4384E-6,0.3740E4,0.9844E-2,0.4587E-6,	PRNC 93
C OGO 5	PRNC 94
• 0.3797E4,2*0.,0.3880E4,2*0.,	PRNC 95
C OGO 6	PRNC 96
• 0.3720E4,0.6738E-2,0.2410E-6,0.3732E4,-.6299E-2,0.2439E-6,	PRNC 97
• 0.3027E4,-.6929E-2,0.4664E-6,0.7242E4,-.2003E-2,-.4585E-6/	PRNC 98
RETURN	PRNC 99
ENTRY PROCES(ISTA,DAY,THETG)	PRNC 100
PRETYP(1)=IPRE/10	PRNC 101
PRETYP(2)=IPRE-PRETYP(1)*10	PRNC 102
PREPRO=.FALSE.	PRNC 103
DO 1000 I=1,2	PRNC 104
IF(PRETYP(1).EQ.0) GO TO 1000	PRNC 105
IF(I.EQ.2) GO TO 200	PRNC 106
IF(MTYPE.NE.2) GO TO 100	PRNC 107
I1=PRETYP(1)	PRNC 108
C TRANSIT TIME	PRNC 109
I2=(R(ISAT1)-OBSD1)/GATE(I1)+0.500	PRNC 110
IF(I2.EQ.0) GO TO 1000	PRNC 111

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

```

OBSO1=OBSO1+GATE(I1)*CFLOAT(I2)
TRANTM=DFLOAT(I2)*GATE(I1)/VLIGHT
GO TO 150
100 TRANTM=? (ISAT1)/VLIGHT
150 DAY=DAY-TRANTM/86400.00
C RECALCULATE ORBIT & GREENWICH HOUR ANGLE
  CALL ORBIT(DAY)
  THETG=GRHRAN(DAY,ISTA)
  GO TO 1000
200 IF(MTYPE.NE.1) GO TO 210
  IF(ISTANJ(ISTA).LT.9000.AND.ISTANQ(ISTA).NE.8009) GO TO 209
C DIURNAL ABERRATION
  T2= IDINT(DAY)
  UT=DAY-T2
  DCOSLT=DCOS(FLAT(ISTA))
  H=UT*24.00+9.856500*IDINT(UT*24.00)/3.603
  H=H*DTWOP/24.00+RLON(ISTA)+THETG-OBSO1
  OBSO2=OBSO2+C.021300*S2R*15.00*DCOSLT*DSIN(H)*DSIN(OBSO2)
  OBSO1=OBSO1+C.32000*S2R*DCOSLT*DCOS(H)/DCOS(OBSO2)
C PARALLACTIC REFRACTION
  PSAT(1)=UHAT(2,ISAT1)
  PSAT(2)=-UHAT(1,ISAT1)
  PSAT(3)=0.000
  STASAT(1)=STAXYZ(2,ISTA)*XYZ(3,ISAT1)
  STASAT(2)=-STAXYZ(3,ISTA)*XYZ(2,ISAT1)
  STASAT(3)=STAXYZ(1,ISTA)*XYZ(3,ISAT1)
  STASAT(1)=STAXYZ(3,ISTA)*XYZ(1,ISAT1)
  STASAT(2)=STAXYZ(1,ISTA)*XYZ(2,ISAT1)
  STASAT(3)=-STAXYZ(2,ISTA)*XYZ(1,ISAT1)
  T2=PSAT(1)**2+PSAT(2)**2
  UT=STASAT(1)**2+STASAT(2)**2+STASAT(3)**2
  T2=DSORT(T2)
  UT=DSORT(UT)
  DO 205 J=1,3
  PSAT(J)=PSAT(J)/T2
205 STASAT(J)=STASAT(J)/UT
  CQ=DOTPRD(PSAT,STASAT)
  STPSAT(1)=UHAT(3,ISAT1)*PSAT(2)-UHAT(2,ISAT1)*PSAT(3)
  STPSAT(2)=-UHAT(3,ISAT1)*PSAT(1)+UHAT(1,ISAT1)*PSAT(3)
  STPSAT(3)=UHAT(2,ISAT1)*PSAT(1)-UHAT(1,ISAT1)*PSAT(2)
  UT=DSORT(STPSAT(1)**2+STPSAT(2)**2+STPSAT(3)**2)
  S0=DOTPRD(STPSAT,STASAT)/UT
  Z=C.2500*DTWOP/1-ELEV(ISAT1)
  IF(ELEV(ISAT1).GT.0.000) GO TO 206
  Z=0.000
  PRINT 10000
10000 FORMAT('0*****$ ELEVATION NEGATIVE. MAXIMUM CORRECTION ',
1 ' $FOR PARALLACTIC REFRACTION USED. *****')
206 RCOSZ=R(ISAT1)*DCOS(Z)
  IF(RCOSZ.GT.1.100) GO TO 207
  H=1.000-DEXP(-1.3850-4*RCOSZ)
  GO TO 203
207 H=1.000
208 DR=0.43500*4.8481300*CTAN(Z)*H/RCOSZ
  OBSO2=OBSO2+CR*CQ
  OBSO1=OBSO1+CR*S0/T2
PROC 112
PROC 113
PROC 114
PROC 115
PROC 116
PROC 117
PROC 118
PROC 119
PROC 120
PROC 121
PROC 122
PROC 123
PROC 124
PROC 125
PROC 126
PROC 127
PROC 128
PROC 129
PROC 130
PROC 131
PROC 132
PROC 133
PROC 134
PROC 135
PROC 136
PROC 137
PROC 138
PROC 139
PROC 140
PROC 141
PROC 142
PROC 143
PROC 144
PROC 145
PROC 146
PROC 147
PROC 148
PROC 149
PROC 150
PROC 151
PROC 152
PROC 153
PROC 154
PROC 155
PROC 156
PROC 157
PROC 158
PROC 159
PROC 160
PROC 161
PROC 162
PROC 163
PROC 164
PROC 165
PROC 166
PROC 167

```

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

C ANNUAL ADAPTATION

209 J=ISTAND(ISTA)
IF(.NOT.(J.EQ.6009.OR.J.EQ.8015.OR.J.EQ.8019.OR.J.EQ.8030.OR.
J.EQ.8014.OR.J.EQ.8018.OR.J.EQ.8023.OR.J.EQ.8027.OR.
J.EQ.8035.OR.J.EQ.8042.OR.J.EQ.8051.OR.J.EQ.8052.OR.
J.EQ.9431.OR.J.EQ.9432)) GO TO 1000

DJ=DJUL(DAY)
CO=CON(DJ,SQ,DR,E)
CALL EPHEM(DAY,.FALSE.)
CE=DCOS(E)
L=DATAN2(SUNY,CE*SUNX)
CL=DCOS(L)
SL=CL*SUNY/(SUNX*CE)
CA=DCOS(OBSO1)
SA=DSIN(OBSO1)
CO=DCOS(OBSO2)
SO=DSIN(OBSO2)
TE=DSIN(E)/CE
OBSO1=OBSO1-20.5D3*S2R*(CA*CL*CE+SA*SL)/CD
OBSO2=OBSO2-20.5D3*S2R*(CL*CE*(TE*CD-SA*SO)+CA*SO*SL)
GO TO 1000

210 SINE=RENV(3,ISAT1)
SRINDX=STQNDX
IF(SRFNDX.NE.0.D0) SRINDX=SRFNDX
IF(MTYPE.GT.3) GO TO 260
VCORR=RCNST*SRINDX
IF(VHFCN) VCORR=VCORR+REFION(MTYPE,ISTA,DAY)
IF(MTYPE.EQ.3) GO TO 220
IF(PRETY(2).GT.2) GO TO 215

C RANGE REFRACTION
OBSO1=OBSO1-VCORR/(0.026D0+SINE)
GO TO 1000

215 COTE=DSORT(1.0D0-SINE**2)/SINE

C RANGE REFRACTION FOR FRENCH LASER
OBSO1=OBSO1-SRFNDX/(SINE+COTE*1.0D-3)
GO TO 1000

C APPLY REFRACTION CORRECTION TO RANGE RATE DATA

220 RATE1=OBSO1(7,ISTA,ECOT)
OBSO1=OBSO1+VCORR*EDOT*DSORT(1.0D0-SINE**2)/(0.026D0+SINE)**2
GO TO 1000

260 CCSF2=1.0D-SINE**2
COSE=DSORT(COSE2)
VCORR=1.0-6*SRINDX
IF(VHFCN.OR.MTYPE.EQ.5) VCORR=VCORR+REFION(MTYPE,ISTA,DAY)
DELTAE=-VCORR/(.01644D0+.93D0*SINE/COSE)
IF(MTYPE.EQ.7) GO TO 270
SINA=RENV(3,ISAT1)/COSE
COSA=RENV(2,ISAT1)/COSE
IF(MTYPE.EQ.6) GO TO 270

C APPLY REFRACTION CORRECTION TO MINI TRACK DATA

OBSO1=OBSO1-SINA*SINE*DELTAE
OBSO2=OBSO2-COSA*SINE*DELTAE
GO TO 1000

C APPLY REFRACTION CORRECTION TO X-Y ANGLE DATA

270 OBSO1=OBSO1-SINA*DELTAE/(SINE**2+SINA**2*COSE2)
OBSO2=OBSO2-COSA*SINE*DELTAE/DSORT(1.0D0-COSE2*COSA**2)





PROC 168
PROC 169
PROC 170
PROC 171
PROC 172
PROC 173
PROC 174
PROC 175
PROC 176
PROC 177
PROC 178
PROC 179
PROC 180
PROC 181
PROC 182
PROC 183
PROC 184
PROC 185
PROC 186
PROC 187
PROC 188
PROC 189
PROC 190
PROC 191
PROC 192
PROC 193
PROC 194
PROC 195
PROC 196
PROC 197
PROC 198
PROC 199
PROC 200
PROC 201
PROC 202
PROC 203
PROC 204
PROC 205
PROC 206
PROC 207
PROC 208
PROC 209
PROC 210
PROC 211
PROC 212
PROC 213
PROC 214
PROC 215
PROC 216
PROC 217
PROC 218
PROC 219
PROC 220
PROC 221
PROC 222
PROC 223



REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

```
GO TO 1000
C APPLY REFRACTION CORRECTION TO ELEVATION ANGLE DATA
280 OBS02=OBS02+DELTA E
1000 CONTINUE
IF(MTYPE.NE.2.OR.CHANEL.EQ.0) RETURN
C APPLY TRANSPONDER DELAY CORRECTION TO RANGE DATA
ISAT=NUMBR4(1SATID(1SAT1),KSATNO,NSATNO)
IF(1SAT.EQ.0) RETURN
ISAT=(1SAT+2)/3
IF(1SAT.EQ.8.AND.CHANEL.GT.3) GO TO 265
GO TO 290
265 ISAT=ISAT+1
CHANEL=CHANEL-3
IF(CHANEL.GT.1) RETURN
290 CHANEL=(CHANEL+2)/2
RDCT=CRSDOT(2,1STA,RDOT2)
DEL R=0.00
DO 300 I=1,3
300 DELR=DEL R+RDCT**((I-1)*TRANSP(I,CHANEL,1SAT))
DEL R=-DEL R*.5D-9*VLIGHT
OBS01=OBS01+DEL R
RETURN
END
```



PROC 224
PROC 225
PROC 226
PROC 227
PROC 228
PROC 229
PROC 230
PROC 231
PROC 232
PROC 233
PROC 234
PROC 235
PROC 236
PROC 237
PROC 238
PROC 239
PROC 240
PROC 241
PROC 242
PROC 243
PROC 244
PROC 245
PROC 246



REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR





REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR





RANDOM

DESCRIPTION

RANDOM is a subroutine used to read, write and read/rewrite satellite observation data.

Since each measurement must be rewritten after it has been preprocessed, direct access methods of I/O are required and since large quantities of data are processed, it is necessary to use a large data buffer.

RANDOM buffers the observation data and calls the direct access reading and writing system subroutines DREAD and DWRITE to perform the I/O operations.

```

NAME          RANDOM
ENTRY POINT   PURPOSE
RANDRD        TO UNBLOCK & READ OBSERVATIONS FROM RANDOM ACCESS
              FILE
RANDWR        TO BLOCK & WRITE OBSERVATIONS ON RANDOM ACCESS
              FILE
CALLING SEQUENCE CALL RANDRD
CALLING SEQUENCE CALL RANDWR
SUBROUTINES USED ERROR
COMMON BLOCKS PREBLK
INPUT FILES   IDISK - RANDOM ACCESS DATA FILE
OUTPUT FILES  IDISK - RANDOM ACCESS DATA FILE
  
```

```

SUBROUTINE RANDRD
IMPLICIT REAL*8 (A-H,C-Z)
COMMON/PREBLK/A(16),IREC
INTEGER A,BUF(16,113),NREC/113/,KBUF/1/,NWORDS/16/
LOGICAL WSWTCH/.FALSE./
EQUIVALENCE (DISK, IDISK)
DATA IDISK/12/
K=1
GO TO 10
ENTRY RANDWR
WSWTCH=.TRUE.
K=2
10  IRUF=(IREC-1)/NREC+1
   IF(IRUF.NE.KBUF)GO TO (50,90),K
20  JREC=IREC-(IRUF-1)*NREC
   GO TO (30,70),K
C UNBLOCK DATA
30  DO 40 I=1,NWORDS
40  A(I)=BUF(I,JREC)
   RETURN
50  IF(WSWTCH)CALL DWRITE(IDISK,KBUF,BUF)
   KBUF=IRUF
   WSWTCH=.FALSE.
C READ BLOCK
CALL DREAD(IDISK,KBUF,BUF,260)
GO TO 30
60 CALL ERROR(11, IDISK)
RETURN
C BLOCK DATA
70  DO 80 I=1,NWORDS
  
```

```

RAND 26
RAND 27
RAND 28
RAND 29
RAND 30
RAND 31
RAND 32
RAND 33
RAND 34
RAND 35
RAND 36
RAND 37
RAND 38
RAND 39
RAND 40
RAND 41
RAND 42
RAND 43
RAND 44
RAND 45
RAND 46
RAND 47
RAND 48
RAND 49
RAND 50
RAND 51
RAND 52
RAND 53
RAND 54
RAND 55
  
```

```
80   BUF(I,JREC)=A(I)  
    RETURN  
C WRITE BLOCK  
90   CALL DWRITE(IDISK,KBUF,BUF)  
    KBUF=IBUF  
    GO TO 23  
*1000 FORMAT(1X,Z8,I6,20A4,4(1X,Z8))  
    END
```

```
RAND 55  
RAND 57  
RAND 58  
RAND 59  
RAND 60  
RAND 61  
RAND 62  
RAND 63
```

4.

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

REARG

DESCRIPTION

REARG recomputes back values of acceleration and the sum array for given changes in stepsize. If the step is to be increased, first REARG will double the stepsize and then reduce to the desired stepsize. When the step is decreased, interpolation is used to obtain midpoint values for the integration values, followed by calls to F and VEVAL to evaluate the accelerations at each midpoint.

NAME REARG

PURPOSE TO RECOMPUTE BACK VALUES OF ACCELERATION AND THE SUM ARRAY FOR GIVEN CHANGES IN STEPSIZE

CALLING SEQUENCE CALL REARG(M,M1,IORDR1,IORDER,H1,H2,T1,T2,FCT,NN,
SUM,SUM1,COEFP,COEFV,Y,Y1,VAR,FCT1,P)




SYMBOL	TYPE	DESCRIPTION
M	I	INPUT - DISPLACEMENT VALUES USED BY INTEGRATOR
M1	I	INPUT - DISPLACEMENT VALUES USED BY INTEGRATOR
IORDR1	I	INPUT - ORDER
IORDER	I	INPUT - ORDER
H1	DP	INPUT - STEPSIZE
H2	DP	INPUT - STEPSIZE
T1	DP	INPUT - TIME
T2	DP	INPUT - TIME
FCT (3,1)	DP	INPUT - BACK VALUE ARRAY OF ACCELERATION
NN	I	INPUT - NUMBER OF EQUATIONS
SUM (2,3,1)	DP	OUTPUT - SUM ARRAY USED BY INTEGRATOR AND INTERPOLATOR
SUM1 (2,3)	DP	OUTPUT - SUM ARRAY USED BY INTEGRATOR AND INTERPOLATOR
COEFP (1)	DP	INPUT - POSITION COEFFICIENTS
COEFV (1)	DP	INPUT - VELOCITY COEFFICIENTS
Y (6,1)	DP	INPUT - ARRAY OF STATE VARIABLES
Y1 (6)	DP	INPUT - ARRAY OF STATE VARIABLES
VAR	L	INPUT - VARIATIONAL EQUATION SWITCH
FCT1 (3)	DP	INPUT - BACK VALUES OF ACCELERATION
P	DP	INPUT - FRACTIONAL STEPSIZE CHANGE




REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

SUBROUTINES USED	INTRP	F	VEVAL	
COMMON BLOCKS	NONE			
INPUT FILES	NONE			
OUTPUT FILES	PRINTER			
SUBROUTINE REARG(M,M1,IORDR1,IORDER,H1,H2,T1,T2,FCT,NN,SUM,				REAR 68
SUM1,COEFP,COEFV,Y,Y1, VAR,FCT1,P)				REAR 69
IMPLICIT REAL*8 (A-H,C-Z)				REAR 70
LOGICAL SWITCH,VAR,BACK				REAR 71
DIMENSION FCT(3,1),ACC(3),SUM(2,3,1),SUM1(2,3) ,COEFP(1),COEFV(1)				REAR 72
Y(6,1),Y1(6) ,FCT1(3)				REAR 73
C TEMPORARY PRINT STATEMENT				REAR 74
H=P*H2				REAR 75
WRITE(6,1) VAR,P,H				REAR 76
1 FORMAT(1X,3G20.8)				REAR 77
H=H2				REAR 78
FACT=P				REAR 79
NECN=1				REAR 80
IF(VAR)NECN=NN-1				REAR 81
IOL2=IORDER-2				REAR 82
IOL1=IOL2+1				REAR 83
IF(FACT,LT.1.D0)GO TO 100				REAR 84
C COMPUTE POSITIONS & VELOCITY AT INTEGRATOR TIME				REAR 85
5 CALL INTRP(0.D0,H2,IORDER,NECN,Y,FCT,M,SUM)				REAR 86
9 DO 10 I=1,IOL2				REAR 87
K1=M-I				REAR 88
K2=M-2*I				REAR 89
DO 10 L=1,NECN				REAR 90
LL=(L-1)*M				REAR 91
FCT(1,LL+K1)=FCT(1,LL+K2)				REAR 92
FCT(2,LL+K1)=FCT(2,LL+K2)				REAR 93
FCT(3,LL+K1)=FCT(3,LL+K2)				REAR 94
10 CONTINUE				REAR 95
11 H=H+H				REAR 96
H2=H2+H2				REAR 97
IF(.NOT.VAR)H1=H1+H1				REAR 98
J3=1				REAR 99
C RECOMPUTE SUMS				REAR 100
GO TO 1000				REAR 101
20 CONTINUE				REAR 102
IF(FACT,GE.2.D0) RETURN				REAR 103
FACT=FACT*.500				REAR 104
100 SWITCH = .FALSE.				REAR 105
DELTAH=FACT*H				REAR 106
IF((H-DELTAH-DELTAH).EQ.0.D0)SWITCH=.TRUE.				REAR 107
I=0				REAR 108
C INTERPOLATE FOR DESIRED BACK VALUE TIMES				REAR 109
101 I=I+1				REAR 110
S1=-DFLOAT(I)*DELTAH				REAR 111

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

TIM=T2+S1	REAR 112
S=(TIM-T1)/H1	REAR 113
CALL INTRP(S,H1,IORDR1,1,Y1,FCT1,M1,SUM1)	REAR 114
IF(.NOT.VAR)CALL F(TIM,Y1,ACC.,TRUE.)	REAR 115
IF(.NOT.VAR)CALL F(TIM,Y1,FCT(1,M-IOL2-1),.FALSE.)	REAR 116
IF(.NOT.VAR)GO TO 105	REAR 117
S=(TIM-T2)/H2	REAR 118
CALL INTRP(S,H2,IORDER,NEON,Y,FCT,M,SUM)	REAR 119
102 CALL VEVAL(Y,FCT(1,M-IOL2-1),6.,.TRUE.,M)	REAR 120
C IF RATIO EQUALS 1/2, JUST RECOMPUTE EVERY OTHER POINT	REAR 121
105 IF(.NOT.SWITCH)GO TO 110	REAR 122
I=I+1	REAR 123
IF(I.GT.IOL2)GO TO 110	REAR 124
LL=M-IOL2-I	REAR 125
LLL=M-(I/2)	REAR 126
FCT(1,LL)=FCT(1,LLL)	REAR 127
FCT(2,LL)=FCT(2,LLL)	REAR 128
FCT(3,LL)=FCT(3,LLL)	REAR 129
110 IF(I.LT.IOL2)GO TO 101	REAR 130
C COMPUTE POSITION & VELOCITY AT INTERGRATOR TIME	REAR 131
111 CALL INTRP(0.00,H2,IORDER,NEON,Y,FCT,M,SUM)	REAR 132
C REPLACE BACK VALUES	REAR 133
DO 115 L=1,NEON	REAR 134
LM=L*M	REAR 135
DO 115 I=1,IOL2	REAR 136
LL=LM-I	REAR 137
LLL=LL-IOL2	REAR 138
FCT(1,LL)=FCT(1,LLL)	REAR 139
FCT(2,LL)=FCT(2,LLL)	REAR 140
FCT(3,LL)=FCT(3,LLL)	REAR 141
115 CONTINUE	REAR 142
H=DELTAH	REAR 143
H2=DELTAH	REAR 144
IF(.NOT.VAR)H1=DELTAH	REAR 145
J3=2	REAR 146
C RECOMPUTE SUMS	REAR 147
GO TO 1000	REAR 148
120 CONTINUE	REAR 149
RETURN	REAR 150
1000 CONTINUE	REAR 151
HS=H**2	REAR 152
DO 1030 N=1,NEON	REAR 153
NM=N*M	REAR 154
LLL=NM +1	REAR 155
DO 1030 J=1,3	REAR 156
A=0.00	REAR 157
B=0.00	REAR 158
DO 1029 I=1,IOL2	REAR 159
LL=LLL-I	REAR 160
A=A-COEFV(I)*FCT(J,LL)	REAR 161
B=B-COEFV(I)*FCT(J,LL)	REAR 162
1029 CONTINUE	REAR 163
1028 A=A-COEFV(IOL1)*FCT(J,NM-IOL2)+FCT(J,NM)	REAR 164
A=Y(J+3,N)/H+A	REAR 165
SUM(1,J,N)=A	REAR 166
SUM(2,J,N)=Y(J,N)/HS +A+B	REAR 167
1030 CONTINUE	REAR 168
GO TO (2,120),J3	REAR 169
RETURN	REAR 170
END	REAR 171





REFCOR

DESCRIPTION

Subroutine REFCOR rotates a vector between the true equator and equinox of a reference date and the true equator and equinox of date.

This routine invokes subroutines PRECES and NUTATE at the start of each arc to obtain the precession rotation matrix from the reference to 1950.0 and the nutation rotation matrix from mean to true equator and equinox of the reference date. These latter are then inverted and combined to rotate from the mean equator and equinox of 1950.0 to the reference time.

The routine obtains rotation matrices from PRECES and NUTATE for consecutive days, stepping in time as required by the input time. These matrices are combined with the above rotation matrix from mean equator and equinox of 1950.0 to the reference time to produce a rotation matrix for 0 hours on each consecutive day.

REFCOR linearly interpolates between these two consecutive rotation matrices to obtain the appropriate rotation matrix for the time of date. The input vector is then rotated either to or from the reference system.

NAME **REFCOR**

PURPOSE **TO PRECESS A VECTOR BETWEEN THE TRUE EQUATOR AND EQUINOX OF A REFERENCE TIME AND THE TRUE EQUATOR AND EQUINOX OF DATE**

CALLING SEQUENCE **CALL REFCOR(DAY,TO,X)**

SYMBOL	TYPE	DESCRIPTION
DAY	DP	INPUT - NUMBER OF DAYS FROM JAN 0.0 OF REFERENCE YEAR
TO	L	INPUT - .TRUE. - CONVERT FROM TRUE OF DATE TO TRUE OF REFERENCE DATE .FALSE. - CONVERT FROM TRUE OF REFERENCE DATE TO TRUE OF DATE
X (3)	DP	INPUT - VECTOR TO BE CONVERTED OUTPUT - CONVERTED VECTOR

SUBROUTINES USED **PRECES NUTATE MULMAT**

COMMON BLOCKS **INITBK CTIME**



INPUT FILES **NONE**


OUTPUT FILES **NONE**

REFERENCES ***GEO DYN SYSTEMS DESCRIPTION*
 VOLUME 1 - GEO DYN DOCUMENTATION**

— SUBROUTINE REFCOR(DAY,TO,X)	REFC	3
REAL*8 P(3,3),N(3,3),REF(3,3),PRE(3,3,2),X(3),Y(3),DAY,DAYP,	REFC	3
DAYREF,PREINT(3,3)	REFC	3
REAL*8 DAYSV/-1.000/	REFC	3
INTEGER IDAY(2)/2*-1/	REFC	4
COMMON/INITBK/IG2(S4),NOT1ST,IG3(2)	REFC	4
LOGICAL NOT1ST	REFC	4
COMMON/CTIME/DATAEP(2),DAYPEF,G1(19)	REFC	4
LOGICAL TO	REFC	4
IF(.NOT.NOT1ST) DAYSV=-1.000	REFC	4
IF(DAY.EQ.DAYSV) GO TO 72	REFC	4
IF(.NOT1ST) GO TO 20	REFC	4
NOT1ST=.TRUE.	REFC	4
C GET PRECESSION & NUTATION FOR REFERENCE TIME	REFC	4
CALL PRECES(DAYREF,P)	REFC	5
CALL NUTATE(DAYREF,N)	REFC	5
C PREMULTIPLY & TRANSPD	REFC	5
DO 10 I=1,3	REFC	5
DO 10 J=1,3	REFC	5
REF(I,J)=0.00	REFC	5

DO 10 K=1,3	REFC 56
10 REF(I,J)=P(J,K)*N(K,I)+REF(I,J)	REFC 57
C ARE VALUES NEEDED FOR INTERPOLATION ?	REFC 58
20 ID=DAY	REFC 59
IF(ID.EQ.IDAY(1)) GO TO 60	REFC 60
INFED=1	REFC 61
C 2 VALUES OR ONLY THE SECOND ?	REFC 62
IF(ID.NE.IDAY(2)) GO TO 35	REFC 63
INEED=2	REFC 64
C I NEED ONLY THE SECOND...SAVE THE PREVIOUS VALUES	REFC 65
DO 30 I=1,3	REFC 66
DO 30 J=1,3	REFC 67
30 PRE(I,J,1)=PRE(I,J,2)	REFC 68
IDAY(1)=IDAY(2)	REFC 69
C GET PRECESSION & NUTATION VALUES FOR DAY OF INTEREST	REFC 70
35 DO 50 M=INEED,2	REFC 71
IDAY(M)=ID+M-1	REFC 72
DAYP=IDAY(M)	REFC 73
CALL PRECES(DAYP,P)	REFC 74
CALL NUTATE(DAYP,N)	REFC 75
C PREMULTIPLY WITH REFERENCE VALUES	REFC 76
50 CALL MULMAT(PRE(1,1,M),REF,P,N)	REFC 77
C OBTAIN INTERPOLATED ROTATION MATRIX	REFC 78
60 DAYP=DAY-DFLOAT(IDAY(1))	REFC 79
DO 70 I=1,3	REFC 80
DO 70 J=1,3	REFC 81
70 PREINT(I,J)=PRE(I,J,1)+DAYP*(PRE(I,J,2)-PRE(I,J,1))	REFC 82
C ROTATE INPUT VECTOR	REFC 83
72 IF(TO) GO TO 80	REFC 84
C ROTATE FROM REFERENCE SYSTEM	REFC 85
DO 75 I=1,3	REFC 86
Y(I)=0.000	REFC 87
DO 75 J=1,3	REFC 88
75 Y(I)=Y(I)+PREINT(J,I)*X(J)	REFC 89
GO TO 90	REFC 90
C ROTATE TO REFERENCE SYSTEM	REFC 91
80 DO 85 I=1,3	REFC 92
Y(I)=0.000	REFC 93
DO 85 J=1,3	REFC 94
85 Y(I)=Y(I)+PREINT(I,J)*X(J)	REFC 95
C STORE RESULT IN INPUT VECTOR	REFC 96
90 DO 100 I=1,3	REFC 97
100 X(I)=Y(I)	REFC 98
DAYSV=DAY	REFC 99
RETURN	REFC 100
END	REFC 101





NAME REFION
PURPOSE DUMMY IONOSPHERIC REFRACTION SUBROUTINE
CALLING SEQUENCE X=REFION(MTYPE,ISTA,DAY)
SYMBOL TYPE DESCRIPTION
MTYPE I INPUT - MEASUREMENT TYPE
ISTA I INPUT - STATION INDEX
DAY DP INPUT - TIME IN DAYS FROM JAN 0.3 OF THE REFERENCE
YEAR
REFION R OUTPUT - IONOSPHERIC REFRACTION VARIATION
SUBROUTINES USED NONE
COMMON BLOCKS NONE
INPUT FILES NONE
OUTPUT FILES NONE

FUNCTION REFION(MTYPE,ISTA,DAY)
REFION=
RETURN
END

FFF I 29
REF I 30
REF I 31
REF I 32

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

NAME RESPAR
ENTRY POINT PURPOSE
RESPRI INITIALIZATION
RESPAR TO CALCULATE THE PARTIALS OF GEOPOTENTIAL
RESONANCE COEFFICIENTS REQUESTING ADJUSTMENT

CALLING SEQUENCE CALL RESPRI(INDXCS, GPSIG, GPPAR)

SYMBOL	TYPE	DESCRIPTION
INDXCS (3,1)	I*2	INPUT - INDICES OF ADJUSTED GEOPOTENTIAL COEFFICIENTS
GPSIG (1)	DP	INPUT - SIGMAS ON ADJUSTED GEOPOTENTIAL COEFFICIENTS
GRPAR (3,1)	DP	OUTPUT - PARTIALS OF FORCE MODEL PARAMETERS

CALLING SEQUENCE CALL RESPAR

SUBROUTINES USED NONE

COMMON BLOCKS CPARAM INTRK VRBLCK XYZ

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES 'GEO DYN SYSTEMS DESCRIPTION'
VOLUME 1 - GEO DYN DOCUMENTATION

SUBROUTINE RESPRI(INDXCS, GPSIG, GRPAR)	RESP	39
IMPLICIT REAL*8 (A-H, C-Z)	RESP	40
LOGICAL CMPGPR	RESP	41
INTEGER*2 INDXCS	RESP	42
DIMENSION INDXCS(3,1), GPSIG(1), GPPAR(3,1)	RESP	43
COMMON/CPARAM/NSTA, NMAST, NSTEST, NDIR, MBIAS, NGPC1, NGPC2, NGPCOM,	RESP	44
NCSEST, CMPGPR, LIM1, LIM2, NOEN, NDNST, NTIDST, NTIDEN, INNRSW,	RESP	45
NCONST, NDCONS	RESP	46
COMMON/INTRK/THDOT1(53), NEON(2), ADDR(7)	RESP	47
COMMON/VRBLCK/XY50, CSM(31,2), UID(7), P(33,30), AGRN(30),	RESP	48
TPSIM(39)	RESP	49
COMMON/XYZ/X, Y, Z, XDOT, YDOT, ZDOT, R, RSQ, ISAT, IFORCE(2)	RESP	50
EQUIVALENCE (CP, P(2,1))	RESP	51
RETURN	RESP	52
ENTRY RESPAR	RESP	53
C1=TPSIM(2)/RSQ	RESP	54
C2=CP/P	RESP	55

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

II=NEON(1SAT)-NCSEST-7-NTIDEN	RFSP	56
DO 100 I=1,NCSEST	RFSP	57
IC=INDXCS(1,I)	RFSP	58
IF(IC.LT.1) GO TO 100	RFSP	59
II=II+1	RFSP	60
ICC=3-IC	RFSP	61
N=INDXCS(2,I)	RFSP	62
M=INDXCS(3,I)	RFSP	63
M1=M+1	RFSP	64
C3=AORN(N)*GPSIG(I)	RFSP	65
C4=C3*CSML(M1,IC)	RFSP	66
FCP=-C4*(P(M1,N)*DFLOAT(N+1))/R	RFSP	67
FCL=C3*CSML(M1,ICC)*P(M1,N)*DFLOAT(M)	RFSP	68
IF(IC.EQ.1) FCL=-FCL	RFSP	69
FCP=C4*(P(M1+1,N)-TPS14(M1)*P(M1,N))	RFSP	70
C3=FCP/R	RFSP	71
C4=C3-C1*FCP	RFSP	72
C5=FCL/XYSO	RFSP	73
GRPAR(1,II)=X*C4-Y*C5	RFSP	74
GRPAR(2,II)=Y*C4+X*C5	RFSP	75
GRPAR(3,II)=Z*C3+FCP*C2	RFSP	76
100 CONTINUE	RFSP	77
RETURN	RFSP	78
END	RFSP	79

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

NAME RFTMCD

PURPOSE TO CHECK AN 80 CHARACTER ALPHANUMERIC STRING TO
 DETERMINE IF THE FIRST 6 CHARACTERS ARE NUMERALS
 AND THE REMAINING 74 CHARACTERS ARE BLANKS

CALLING SEQUENCE X=RFTMCD(A)

SYMBOL	TYPE	DESCRIPTION
A	I*2	INPUT - CHARACTER STRING
RFTMCD	L	OUTPUT - .TRUE. WHEN FIRST 6 CHARACTERS ARE NUMERALS AND REMAINING 74 ARE BLANKS

SUBROUTINES USED NONE

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE

LOGICAL FUNCTION RFTMCD(A)	RFTM	27
INTEGER*2 A(1),NUMBR5(10),BL	RFTM	28
DATA NUMBR5/1H0, 1H1, 1H2, 1H3, 1H4, 1H5, 1H6, 1H7, 1H8, 1H9/.3L/1H /	RFTM	29
RFTMCD=.FALSE.	RFTM	30
DO 20 I=1,6	RFTM	31
DO 10 J=1,10	RFTM	32
IF(A(I).EQ.NUMBR5(J)) GO TO 20	RFTM	33
10 CONTINUE	RFTM	34
RETURN	RFTM	35
20 CONTINUE	RFTM	36
DO 30 I=7,80	RFTM	37
IF(A(I).NE.BL) RETURN	RFTM	38
30 CONTINUE	RFTM	39
RFTMCD=.TRUE.	RFTM	40
RETURN	RFTM	41
END	RFTM	42

REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR.

NAME RMSCMP

PURPOSE TO COMPUTE RMS, RND, AND MEAN FROM SUMMED INFORMATION

CALLING SEQUENCE CALL RMSCMP(NSUM,ASUM,NTYPE)

SYMBOL	TYPE	DESCRIPTION
NSUM (3,1)	I	INPUT - MEASUREMENT TYPE, NUMBER OF RESIDUALS, AND NUMBER OF WEIGHTED RESIDUAL RATIOS FOR THIS PARTICULAR STATION, SATELLITE, AND MEASUREMENT TYPE
ASUM (8,1)	R	INPUT - SUMMING ARRAYS FOR THIS PARTICULAR STATION SATELLITE & MEASUREMENT TYPE
NTYPE	I	INPUT - MAXIMUM NUMBER OF POSSIBLE TYPES FOR THIS STATION & SATELLITE (MAX. = 4 FOR TYPES 1-14 = 2 FOR TYPES 27-30)

SUBROUTINES USED NONE

COMMON BLOCKS CSTINF

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES *GEODYN SYSTEMS DESCRIPTION*
VOLUME 1 - GEODYN DOCUMENTATION

SUBROUTINE RMSCMP(NSUM,ASUM,NTYPE)	RMSC	37
DIMENSION NSUM(3,1),ASUM(8,1)	RMSC	38
COMMON/CSTINF/MEASNO(4),NORS(4),RDMEAN(4),RMSO(4),RND(4),	RMSC	39
MEASWT(4),WTMEAN(4),RMSWTO(4),WTRND(4),TYPRMS(30),NOTYPE(2,30),	RMSC	40
BSUM(5,12),PHSALL(20),NDALL(30),NDWTOB,LSASE	RMSC	41
DO 300 I=1,NTYPE	RMSC	42
J=NSUM(3,I)	RMSC	43
IF(J.NE.0) GO TO 100	RMSC	44
NORS(I)=0	RMSC	45
RETURN	RMSC	46
100 NORS(I)=NSUM(1,I)	RMSC	47
MEASWT(I)=NSUM(2,I)	RMSC	48
MEASNO(I)=J	RMSC	49
TYPRMS(J)=TYPRMS(J)+ASUM(4,I)	RMSC	50
NOTYPE(2,J)=NOTYPE(2,J)+MEASWT(I)	RMSC	51
XN=NORS(I)	RMSC	52
RDMEAN(I)=ASUM(1,I)/XN	RMSC	53
RMSO(I)=0.	RMSC	54
IF(NORS(I).LT.10) GO TO 200	RMSC	55

RMSO(I)=SQRT(ASUM(3,I)/(XN-1.))	RMSC	56
RND(I)=(ASUM(3,I)-ASUM(1,I)**2/XN)/ASUM(5,I)	RMSC	57
RND(I)=(2.*RND(I)-1.)/SQRT((XN-2.)/(XN**2-1.))	RMSC	58
200 IF(MEASWT(I).EQ.0) GO TO 300	RMSC	59
XN=MEASWT(I)	RMSC	60
WTMEAN(I)=ASUM(2,I)/XN	RMSC	61
IF(MEASWT(I).LT.10) GO TO 300	RMSC	62
RMSWTO(I)=SQRT(ASUM(4,I)/(XN-1.))	RMSC	63
WTRND(I)=(ASUM(4,I)-ASUM(2,I)**2/XN)/ASUM(6,I)	RMSC	64
WTRND(I)=(2.*WTRND(I)-1.)/SQRT((XN-2.)/(XN**2-1.))	RMSC	65
300 CONTINUE	RMSC	66
RETURN	RMSC	67
END	RMSC	68

NAME ROTMAT
PURPOSE TO GENERATE A ROTATION MATRIX FROM AN ANGLE AND
AXIS OF ROTATION

CALLING SEQUENCE CALL ROTMAT(THETA,I1,X)

SYMBOL	TYPE	DESCRIPTION
THETA	DP	INPUT - THE ROTATION ANGLE
I1	I	INPUT - THE ROTATION AXIS
X (3,3)	DP	OUTPUT - THE ROTATION MATRIX

SUBROUTINES USED NONE

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE

SUBROUTINE ROTMAT(THETA,I1,X)

REAL*8 THETA,X(3,3)

I2=MOD(I1,3)+1

I3=MOD(I2,3)+1

X(I1,I1)=1.00

X(I1,I2)=0.00

X(I1,I3)=0.00

X(I2,I1)=0.00

X(I3,I1)=0.00

X(I2,I2)=DCOS(THETA)

X(I3,I3)=X(I2,I2)

X(I2,I3)=DSIN(THETA)

X(I3,I2)=-X(I2,I3)

RETURN

END

POTM 28
ROTM 29
ROTM 30
ROTM 31
ROTM 32
ROTM 33
ROTM 34
ROTM 35
ROTM 36
ROTM 37
ROTM 38
ROTM 39
POTM 40
ROTM 41
ROTM 42

NAME SATCLC

PURPOSE TO APPLY SATELLITE CLOCK CORRECTIONS TO GEOS-I OPTICAL DATA

CALLING SEQUENCE SATCLC(FIRST, DAY)

SYMBOL TYPE DESCRIPTION

FIRST L INPUT & OUTPUT - INITIALIZATION SWITCH

DAY DP INPUT - UNCORRECTED MEASUREMENT TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR FOR THE ARC

SATCLC R OUTPUT - CLOCK CORRECTION IN SECONDS

SUBROUTINES USED YMCAY

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE

FUNCTION	SATCLC(FIRST, DAY)	SATC	28
DIMENSION	DAYNO(126), DAYNO2(37), CLC(126), CLC2(37)	SATC	29
EQUIVALENCE	(DAYNO(127), DAYNO2(1)), (CLC(127), CLC2(1))	SATC	30
REAL*8	LAUNCH, CDAY, ODAY, DAY, YMDAY	SATC	31
LOGICAL	FIRST	SATC	32
DATA	NCL7CK/163/	SATC	33
DATA	DAYNO/	SATC	34
•	9.000, 9.721, 10.641, 13.562, 14.562, 15.304, 15.567	SATC	35
•	16.571, 17.579, 17.579, 19.312, 19.575, 20.579, 20.579	SATC	36
•	21.583, 22.500, 23.587, 24.508, 25.513, 26.513, 27.429	SATC	37
•	29.000, 30.442, 31.442, 32.446, 33.450, 34.367, 47.409	SATC	38
•	48.054, 49.054, 50.125, 51.329, 52.152, 52.241, 54.805	SATC	39
•	55.895, 58.900, 60.095, 60.829, 60.829, 64.835, 65.845	SATC	40
•	66.845, 67.845, 67.845, 68.212, 68.762, 70.771, 70.771	SATC	41
•	71.771, 71.771, 73.779, 73.779, 74.056, 74.696, 74.696	SATC	42
•	78.704, 78.704, 79.712, 79.988, 80.713, 85.642, 85.642	SATC	43
•	85.837, 86.650, 89.033, 89.654, 93.575, 93.575, 94.571	SATC	44
•	94.571, 97.071, 97.591, 100.600, 101.520, 101.520, 102.516	SATC	45
•	102.516, 103.346, 103.521, 105.441, 105.795, 106.446, 106.800	SATC	46
•	108.904, 108.804, 109.371, 109.720, 110.725, 110.725, 111.737	SATC	47
•	111.921, 112.733, 113.733, 113.773, 114.387, 114.746, 114.929	SATC	48
•	115.737, 116.395, 116.745, 117.658, 117.658, 118.291, 118.658	SATC	49
•	119.566, 119.566, 122.075, 122.654, 135.191, 135.538, 136.745	SATC	50
•	136.545, 137.455, 137.455, 138.104, 138.104, 138.545, 139.545	SATC	51
•	154.104, 154.929, 170.937, 170.937, 171.962, 171.962, 177.792	SATC	52
DATA	DAYNO2/	SATC	53
•	177.792, 191.108, 191.562, 261.092, 261.092, 262.092, 268.452	SATC	54
•	269.020, 282.896, 282.896, 283.891, 283.891, 284.900, 284.900	SATC	55

•	289.263,	289.883,	316.916,	317.916,	324.854,	324.854,	326.858,	SATC	56
•	326.858,	358.550,	358.550,	373.396,	373.396,	375.137,	375.229,	SATC	57
•	377.054,	378.050,	379.052,	379.052,	382.983,	382.983,	382.983,	SATC	58
•	383.983,	388.000/						SATC	59

DATA CLC/

•	-7.20,	-8.20,	-0.35,	-0.03,	2.80,	2.70,	1.10,	SATC	61
•	1.10,	1.10,	0.20,	0.20,	1.00,	1.00,	1.30,	SATC	62
•	1.10,	1.10,	1.00,	2.10,	2.20,	2.28,	0.10,	SATC	63
•	-0.10,	-0.90,	-0.90,	0.10,	0.30,	-0.20,	0.20,	SATC	64
•	-3.80,	-0.85,	-0.70,	-0.70,	-0.30,	-0.30,	-1.10,	SATC	65
•	0.10,	0.10,	-0.10,	-0.60,	-0.10,	0.50,	0.80,	SATC	66
•	0.10,	0.40,	0.10,	0.10,	0.10,	0.50,	-0.10,	SATC	67
•	0.30,	-0.30,	-0.30,	-0.10,	-2.10,	-2.00,	0.10,	SATC	68
•	-0.30,	0.25,	0.20,	0.20,	-0.30,	-0.60,	0.20,	SATC	69
•	0.20,	0.20,	0.10,	-0.10,	-0.10,	-2.50,	-2.50,	SATC	70
•	-0.20,	-0.30,	-0.20,	0.30,	-2.60,	-0.50,	3.60,	SATC	71
•	0.20,	0.20,	-0.20,	-2.00,	0.20,	0.20,	-0.50,	SATC	72
•	-0.40,	0.20,	0.20,	0.30,	0.90,	0.40,	0.40,	SATC	73
•	0.10,	-0.60,	0.30,	-0.40,	-0.20,	-0.20,	-0.20,	SATC	74
•	-0.50,	-0.45,	0.10,	0.40,	-0.10,	0.10,	0.20,	SATC	75
•	0.60,	-0.40,	0.10,	-0.20,	-0.10,	-0.30,	-0.30,	SATC	76
•	1.70,	1.70,	-0.10,	0.10,	1.70,	1.70,	0.30,	SATC	77
•	0.10,	0.10,	0.40,	-0.50,	-0.50,	-0.40,	-0.30,	SATC	78

DATA CLC2/

•	0.0,	-0.30,	-0.10,	-0.30,	5.20,	0.20,	0.10,	SATC	79
•	0.10,	-0.20,	-5.00,	-4.90,	0.50,	0.50,	-0.10,	SATC	80
•	-0.10,	-0.10,	-0.40,	-0.20,	-0.40,	-0.20,	-0.40,	SATC	81
•	+0.20,	-0.45,	+0.10,	-0.32,	-0.01,	-0.03,	+0.05,	SATC	82
•	+0.07,	-0.50,	-0.50,	+0.23,	+0.40,	+0.22,	+0.30,	SATC	83
•	+0.24,	+0.50/						SATC	84

C DEFINE REFERENCE TIME

	IF(FIRST) LAUNCH=YMDAY(651108.0,0,0,0)	SATC	85
	FIRST=.FALSE.	SATC	86
	ODAY=DAY+8.6404	SATC	87
	CDAY=IDINT(ODAY+2.00)/4*4	SATC	88
C SET TIME TO MULTIPLE OF 4 SECONDS	SATCLC=CDAY-CDAY+.50-3	SATC	89
	REFTIM=DAY-LAUNCH	SATC	90
	DO 10 I=2,NLOCK	SATC	91
	IF(REFTIM.GT.DAYNO(I)) GO TO 10	SATC	92

C INTERPOLATE

20	RATE=(CLC(I)-CLC(I-1))/(DAYNO(I)-DAYNO(I-1))	SATC	93
	SATCLC=SATCLC+(RATE*(REFTIM-DAYNO(I-1))+CLC(I-1))/1.E3	SATC	94
	RETURN	SATC	95
10	CONTINUE	SATC	96
	RETURN	SATC	97
	END	SATC	98

NAME SATCL2

PURPOSE TO APPLY SATELLITE CLOCK CORRECTIONS TO GEOS-II OPTICAL DATA

CALLING SEQUENCE X=SATCL2(FIRST, DAY)

SYMBOL TYPE DESCRIPTION

FIRST L INPUT & OUTPUT - INITIALIZATION SWITCH

DAY DP INPUT - UNCORRECTED MEASUREMENT TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR FOR THE APC

SATCL2 R OUTPUT - CLOCK CORRECTION IN SECONDS

SUBROUTINES USED SATC21 SATC22 YMDAY

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE

FUNCTION SATCL2(FIRST, DAY)

DIMENSION DAYNO(500), DAYNO1(115), DAYNO2(51), DAYNO3(114),
 DAYNO4(114), DAYNO5(114), CLC(500), CLC1(115), CLC2(51), CLC3(114),
 CLC4(114), CLC5(114), DAYS(3)

EQUIVALENCE (DAYNO1(1), DAYNO(1)), (DAYNO2(1), DAYNO(116)),
 (DAYNO3(1), DAYNO(167)), (DAYNO4(1), DAYNO(268)),
 (DAYNO5(1), DAYNO(369)), (CLC1(1), CLC(1)),
 (CLC2(1), CLC(116)), (CLC3(1), CLC(167)), (CLC4(1), CLC(268)),
 (CLC5(1), CLC(369))

DATA DAYS/232.2979, 457.4813, 789.9319/
 DATA DAYNO1/
 51.0254, 52.0396, 52.7632, 53.2529, 54.0647, 55.0798, SATC 28
 55.5479, 55.6514, 56.1146, 56.6511, 57.0335, 57.6750, SATC 29
 58.0437, 58.6910, 59.0589, 59.0750, 61.0873, 63.7278, SATC 30
 63.6237, 64.6238, 64.6236, 65.0425, 66.6472, 66.7208, SATC 31
 67.8568, 67.6523, 69.1007, 68.5224, 68.5751, 69.0375, SATC 32
 70.0625, 71.1035, 72.0722, 72.0625, 72.6451, 73.0160, SATC 33
 73.0903, 73.6583, 74.0285, 74.1242, 74.6715, 75.0417, SATC 34
 75.1174, 76.6847, 76.0549, 75.1312, 76.6215, 77.0574, SATC 35
 77.6347, 77.7118, 78.0819, 78.4826, 78.7250, 79.0907, SATC 36
 79.6618, 80.1076, 80.5945, 80.6750, 81.6118, 81.6887, SATC 37
 82.643, 82.6251, 82.771, 83.0715, 84.6513, 84.6514, SATC 38
 85.0779, 85.5496, 85.6533, 85.5723, 86.1111, 86.6021, SATC 39
 87.6153, 87.6017, 88.0614, 88.6225, 88.7048, SATC 40
 89.6417, 89.7187, 90.0882, 90.1665, 90.6518, SATC 41
 91.6830, 91.6887, 92.1146, 92.6155, 92.6819, SATC 42
 93.0521, 93.1278, 93.5977, 94.1417, 94.6310, SATC 43
 95.785, 95.6472, 95.6496, 95.7222, 96.0933, SATC 44

•	96.7375.	97.5965.	97.6722.	97.7486.	99.1130.	99.6111.	SATC	56
•	98.5847.	99.0562.	99.1312.	99.6222.	99.6296.	100.0687/	SATC	57
	DATA DAYNO2/						SATC	58
•	100.1451.	100.6354.	101.2917.	101.5486.	101.7253.	102.0951.	SATC	59
•	102.5418.	103.6750.	103.7531.	104.1215.	104.5172.	104.8937.	SATC	60
•	105.1347.	105.6257.	105.7021.	106.1486.	106.6339.	106.7153.	SATC	61
•	107.3354.	107.1619.	107.6576.	108.6694.	108.5697.	109.6940.	SATC	62
•	110.4965.	110.6972.	111.7104.	112.7236.	113.6611.	114.6670.	SATC	63
•	114.6637.	115.6743.	115.6743.	115.7501.	116.1285.	116.6201.	SATC	64
•	117.1417.	117.6333.	117.7090.	118.1548.	118.6465.	119.7222.	SATC	65
•	117.6646.	119.7361.	120.1055.	120.6715.	120.7493.	121.1197.	SATC	66
•	121.5861.	121.7623/					SATC	67
	DATA DAYNO3/						SATC	68
•	122.1319.	122.6993.	122.7764.	123.1451.	123.6353.	123.7125.	SATC	69
•	124.1593.	124.6550.	124.7326.	125.1722.	125.6632.	125.7389.	SATC	70
•	126.1690.	126.6764.	126.7529.	127.1222.	127.6336.	127.7660.	SATC	71
•	128.1354.	128.7028.	129.7111.	129.7792.	129.1449.	129.6403.	SATC	72
•	129.7160.	130.0868.	130.1618.	130.7292.	131.0973.	131.1750.	SATC	73
•	131.5567.	132.6502.	132.6799.	132.7562.	133.1257.	133.6931.	SATC	74
•	133.7594.	134.1289.	134.7002.	134.7426.	135.1521.	135.6437.	SATC	75
•	135.7194.	135.7250.	136.1660.	136.6569.	136.7326.	137.1785.	SATC	76
•	137.5701.	137.7465.	138.1160.	138.1924.	138.6443.	139.1202.	SATC	77
•	139.6265.	139.7729.	140.1423.	140.7097.	140.7869.	141.1555.	SATC	78
•	141.6479.	141.7229.	142.1627.	142.5611.	142.7364.	143.1910.	SATC	79
•	143.6736.	143.7500.	144.1194.	144.6964.	144.7632.	145.1326.	SATC	80
•	145.7000.	146.1451.	146.7129.	146.8310.	147.1533.	147.7271.	SATC	81
•	147.9042.	148.1722.	148.6646.	148.7403.	149.7465.	149.1854.	SATC	82
•	149.6778.	149.7535.	149.7613.	150.1386.	150.6910.	150.7667.	SATC	83
•	151.1361.	151.7042.	151.7915.	152.1436.	152.7183.	152.7937.	SATC	84
•	152.1518.	152.7305.	152.7486.	153.1989.	153.6412.	153.7569.	SATC	85
•	156.2021.	156.6944.	156.7706.	156.7767.	157.1336.	157.7076.	SATC	86
•	157.7840.	158.1521.	158.7218.	158.7972.	159.1553.	159.7340/	SATC	87
	DATA DAYNO4/						SATC	88
•	159.9111.	160.1785.	160.6722.	160.7472.	160.7562.	161.1924.	SATC	89
•	161.6959.	161.7611.	162.2056.	162.6984.	162.7743.	163.1431.	SATC	90
•	163.1137.	163.7118.	164.1555.	164.2326.	164.7250.	165.1687.	SATC	91
•	165.7382.	166.8146.	166.1074.	166.1917.	166.7514.	167.1208.	SATC	92
•	167.1954.	167.7646.	168.2040.	169.7774.	169.2222.	169.7153.	SATC	93
•	169.7917.	170.1597.	170.2361.	170.7256.	170.7352.	171.1722.	SATC	94
•	171.7417.	171.8191.	172.1814.	172.7549.	172.8312.	173.1997.	SATC	95
•	173.6931.	173.7681.	174.2125.	174.7903.	175.1530.	175.2257.	SATC	96
•	175.7187.	176.1632.	176.2396.	176.7312.	177.1764.	177.2528.	SATC	97
•	177.7451.	178.1489.	178.7590.	178.8354.	179.2329.	179.7222.	SATC	98
•	180.1410.	180.2160.	180.7554.	181.2292.	181.7229.	181.7996.	SATC	99
•	182.1467.	182.2430.	182.7361.	182.7472.	183.1700.	183.2562.	SATC	100
•	183.7493.	183.8257.	184.1971.	184.2701.	184.7625.	184.8390.	SATC	101
•	185.1319.	185.2062.	185.7757.	186.7022.	187.2326.	187.8029.	SATC	102
•	188.1701.	188.2465.	188.7306.	188.7437.	189.7543.	190.1955.	SATC	103
•	190.2736.	190.7560.	191.7792.	192.2220.	192.7230.	192.7970.	SATC	104
•	193.1611.	193.2761.	193.7340.	193.8062.	194.1736.	194.2500.	SATC	105
•	194.7437/	195.7569.	195.8326.	196.8100.	196.2771.	196.7701.	SATC	106
•	197.7233.	198.1531.	198.2264.	198.7265.	199.2326.	199.7340/	SATC	107
	DATA DAYNO5/						SATC	108
•	199.3097.	200.1778.	200.7472.	200.8229.	201.1703.	201.7624.	SATC	109
•	202.7730.	202.8520.	203.1471.	203.2167.	203.2914.	203.7858.	SATC	110
•	203.2532.	204.1555.	204.2269.	204.8001.	204.8771.	205.1690.	SATC	111

•	205.2430.	205.7375.	205.8132.	205.1812.	206.2540.	206.8271.	SATC 112	
•	207.1944.	207.2701.	207.7539.	207.8403.	208.1340.	208.2050.	SATC 113	
•	208.2340.	208.7771.	208.8535.	208.2201.	209.2370.	209.7003.	SATC 114	
•	209.4574.	210.1597.	210.2333.	210.3118.	210.7295.	210.8042.	SATC 115	
•	211.1732.	211.2473.	211.7417.	211.8174.	212.1847.	212.2604.	SATC 116	
•	212.7549.	212.8316.	212.1970.	212.2734.	212.7541.	212.8437.	SATC 117	
•	213.4472.	214.1222.	214.1471.	214.2111.	214.2875.	214.7812.	SATC 118	
•	214.4576.	215.1507.	215.2243.	215.2347.	215.3014.	215.7044.	SATC 119	
•	215.7098.	215.8709.	215.1622.	215.2373.	215.3153.	215.8076.	SATC 120	
•	217.1757.	217.2507.	217.7441.	217.8208.	218.1049.	218.2630.	SATC 121	
•	218.7593.	220.7989.	221.2278.	221.3249.	221.7070.	222.2410.	SATC 122	
•	220.8111.	220.8882.	220.1700.	220.2542.	220.8213.	220.1924.	SATC 123	
•	224.2473.	224.8375.	225.2912.	225.7750.	225.4514.	226.2187.	SATC 124	
•	226.7482.	226.8546.	227.2243.	227.3083.	227.8014.	228.2044.	SATC 125	
•	228.3146.	228.8917.	229.1840.	229.3278.	230.1945.	230.2715.	SATC 126	
•	230.4417.	231.2090.	231.2847.	231.8549.	232.2222.	232.2970.	SATC 127	
•	DATA CLC1/						SATC 128	
•	0.390.	0.326.	0.266.	0.200.	0.110.	-0.105.	SATC 129	
•	-0.120.	-0.050.	-0.250.	-0.334.	-0.401.	-0.477.	SATC 130	
•	-0.542.	-0.489.	-0.960.	-0.300.	0.450.	0.450.	SATC 131	
•	0.733.	0.673.	-0.150.	-0.169.	-0.150.	-0.150.	SATC 132	
•	0.406.	0.502.	0.575.	0.643.	0.575.	0.127.	SATC 133	
•	0.190.	0.130.	0.184.	0.188.	0.212.	0.254.	SATC 134	
•	0.277.	0.211.	0.352.	0.237.	-0.255.	-0.762.	SATC 135	
•	-0.770.	-0.197.	-0.038.	-0.060.	-0.060.	0.021.	SATC 136	
•	0.007.	0.031.	0.060.	0.063.	0.108.	0.154.	SATC 137	
•	0.131.	0.164.	0.151.	0.152.	0.141.	0.164.	SATC 138	
•	0.113.	0.155.	0.138.	0.093.	0.093.	7.244.	7.251.	SATC 139
•	7.202.	7.277.	7.275.	0.072.	0.049.	0.107.	SATC 140	
•	0.132.	0.131.	0.156.	0.150.	0.125.	0.131.	SATC 141	
•	0.167.	0.160.	0.184.	0.174.	0.178.	0.175.	SATC 142	
•	0.230.	0.256.	0.291.	0.300.	0.304.	0.336.	SATC 143	
•	0.345.	0.355.	0.437.	0.486.	0.412.	0.430.	SATC 144	
•	0.462.	0.515.	-0.360.	-0.325.	-0.314.	-0.248.	SATC 145	
•	-0.246.	-0.164.	-0.146.	-0.135.	-0.105.	-0.040.	SATC 146	
•	-0.073.	0.017.	0.030.	0.025.	-0.039.	0.154.	SATC 147	
•	DATA CLC2/						SATC 148	
•	0.167.	0.146.	0.220.	0.197.	0.225.	0.250.	SATC 149	
•	0.320.	0.349.	0.360.	0.465.	0.511.	0.195.	SATC 150	
•	0.255.	0.296.	0.331.	0.370.	0.452.	0.450.	SATC 151	
•	0.471.	0.482.	0.020.	0.021.	0.022.	0.245.	SATC 152	
•	-0.085.	-0.066.	0.729.	0.088.	-0.100.	-0.100.	SATC 153	
•	15.253.	15.253.	-0.257.	-0.263.	-0.281.	-0.242.	SATC 154	
•	-0.217.	-0.198.	-0.143.	-0.153.	-0.171.	-0.141.	SATC 155	
•	-0.130.	-0.151.	-0.140.	-0.147.	-0.045.	-0.076.	SATC 156	
•	-0.057.	-0.045.	-0.030.				SATC 157	
•	DATA CLC3/						SATC 158	
•	-0.052.	-0.003.	0.050.	-0.025.	-0.014.	-0.054.	SATC 159	
•	-0.045.	-0.073.	-0.070.	-0.135.	-0.080.	-0.086.	SATC 160	
•	-0.142.	-0.142.	-0.148.	-0.023.	-0.037.	-0.034.	SATC 161	
•	-0.311.	0.088.	-0.077.	-0.040.	-0.019.	-0.011.	SATC 162	
•	0.000.	0.066.	0.078.	0.070.	0.117.	0.127.	SATC 163	
•	0.070.	0.150.	0.175.	0.179.	0.206.	0.272.	SATC 164	
•	0.040.	0.003.	0.004.	0.007.	0.007.	0.005.	SATC 165	
•	0.157.	0.101.	0.004.	0.151.	0.157.	0.134.	SATC 166	
•	0.125.	0.105.	0.057.	0.123.	0.090.	0.103.	SATC 167	

•	0.071.	0.098.	-0.059.	-0.078.	-0.044.	-0.140.	SATC 168
•	-0.188.	-0.177.	-0.140.	-0.164.	-0.158.	-0.151.	SATC 169
•	-0.190.	-0.201.	-0.214.	-0.239.	-0.243.	-0.314.	SATC 170
•	-0.330.	-0.362.	-0.342.	-0.347.	-0.351.	-0.348.	SATC 171
•	-0.354.	-0.418.	-0.372.	-0.361.	0.476.	0.420.	SATC 172
•	-0.432.	0.449.	-0.076.	-0.150.	-0.035.	-0.043.	SATC 173
•	-0.026.	-0.029.	-0.012.	-0.051.	-0.016.	-0.010.	SATC 174
•	0.031.	0.016.	0.134.	-0.137.	-0.219.	-0.260.	SATC 175
•	-0.266.	-0.415.	-0.448.	-0.004.	-0.029.	-0.140.	SATC 176
•	-0.154.	-0.115.	-0.264.	-0.249.	-0.240.	-0.330.	SATC 177
•	DATA CLC47						SATC 178
•	-0.258.	-0.295.	-0.325.	-0.318.	-0.033.	-0.076.	SATC 179
•	-0.059.	-0.050.	-0.082.	-0.103.	-0.100.	-0.164.	SATC 180
•	-0.112.	-0.128.	-0.123.	-0.140.	-0.144.	-0.172.	SATC 181
•	-0.125.	-0.113.	-0.250.	-0.237.	-0.180.	-0.198.	SATC 182
•	-0.212.	-0.175.	-0.207.	-0.013.	-0.195.	-0.208.	SATC 183
•	-0.213.	-0.237.	-0.176.	-0.188.	0.245.	0.220.	SATC 184
•	0.212.	0.192.	0.192.	0.154.	0.213.	0.129.	SATC 185
•	-0.291.	0.191.	0.117.	0.342.	0.374.	0.314.	SATC 186
•	0.337.	0.313.	0.299.	0.304.	0.307.	0.266.	SATC 187
•	0.344.	0.309.	0.264.	0.366.	0.236.	0.053.	SATC 188
•	-0.285.	-0.116.	-0.136.	-0.254.	-0.474.	-0.420.	SATC 189
•	-0.450.	-0.463.	-0.506.	0.232.	0.150.	0.124.	SATC 190
•	0.017.	0.012.	-0.105.	-0.147.	-0.284.	-0.303.	SATC 191
•	-0.355.	-0.363.	-0.384.	0.451.	0.314.	0.026.	SATC 192
•	0.118.	0.113.	0.458.	0.485.	-0.223.	-0.244.	SATC 193
•	-0.244.	-0.186.	0.456.	0.505.	0.312.	-0.434.	SATC 194
•	-0.476.	-0.415.	-0.372.	-0.368.	-0.332.	-0.320.	SATC 195
•	-0.281.	-0.313.	-0.332.	-0.274.	-0.249.	-0.212.	SATC 196
•	-0.108.	-0.130.	-0.100.	-0.408.	-0.473.	-0.447.	SATC 197
•	DATA CLC57						SATC 198
•	-0.445.	-0.421.	-0.402.	-0.405.	-0.419.	-0.327.	SATC 199
•	-0.150.	-0.155.	-0.188.	-0.150.	-0.122.	-0.140.	SATC 200
•	-0.175.	-0.200.	-0.174.	-0.141.	-0.134.	-0.148.	SATC 201
•	-0.183.	-0.151.	-0.158.	-0.175.	-0.159.	-0.109.	SATC 202
•	-0.165.	-0.148.	-0.137.	-0.067.	-0.075.	-0.067.	SATC 203
•	-0.080.	-0.098.	-0.034.	-0.062.	-0.045.	-0.005.	SATC 204
•	-0.001.	0.018.	0.007.	-0.009.	-0.047.	-0.012.	SATC 205
•	-0.030.	0.023.	-0.004.	0.006.	0.052.	0.061.	SATC 206
•	0.065.	0.059.	0.056.	0.099.	0.035.	0.131.	SATC 207
•	0.014.	0.019.	-0.172.	-0.185.	-0.218.	-0.200.	SATC 208
•	-0.146.	-0.140.	-0.177.	-0.476.	-0.472.	-0.491.	SATC 209
•	0.052.	0.078.	0.041.	0.051.	0.057.	0.076.	SATC 210
•	0.131.	0.110.	0.109.	0.158.	0.130.	0.163.	SATC 211
•	0.183.	-0.420.	-0.356.	-0.382.	-0.417.	-0.353.	SATC 212
•	-0.354.	-0.343.	-0.313.	-0.291.	-0.285.	-0.251.	SATC 213
•	-0.251.	-0.237.	-0.126.	-0.187.	-0.180.	-0.115.	SATC 214
•	-0.108.	-0.076.	-0.064.	-0.145.	-0.069.	-0.044.	SATC 215
•	-0.039.	-0.002.	-0.052.	0.021.	0.041.	0.040.	SATC 216
•	0.005.	0.099.	0.056.	0.119.	0.106.	0.125.	SATC 217
•	REAL*P LAUNCH,CDAY,DDAY,DAY,MO DAY						SATC 218
•	LOGICAL FIRST						SATC 219
•	DATA NCLCK/DRY						SATC 220
•	C DEFINE REFERENCE TIME						SATC 221
•	IF (FIRST) LAUNCH*YMDAY(000101.0.0.00)						SATC 222
•	FIRST=.FALSE.						SATC 223

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

```
DDAY=DAY+2.6404
CDAY=(INT(DDAY+2.00)/4+4
C SET TIME TO MULTIPLE OF 4 SECONDS
SATCL2=CDAY-CDAY+.50-2
REFTIM=LAY-LAUNCH
IF(REFTIM.GT.DAYS(3)) RETURN
N=1
IF(REFTIM.GE.DAYS(N).AND.REFTIM.LT.DAYS(N+1)) GO TO 300
N=2
IF(REFTIM.GE.DAYS(N).AND.REFTIM.LT.DAYS(N+1)) GO TO 400
DO 10 I=2,N,CLOCK
IF(REFTIM.GT.DAYNO(I)) GO TO 10
C INTERPOLATE
RATE=(CLC(I)-CLC(I-1))/(DAYNO(I)-DAYNO(I-1))
SATCL2=SATCL2+(RATE*(REFTIM-DAYNO(I-1))+CLC(I-1))*L.F-3
RETURN
10 CONTINUE
RETURN
300 SATCL2=SATCL2+SATC21(REFTIM)
RETURN
400 SATCL2=SATCL2+SATC22(REFTIM)
RETURN
END
```

SATC 224
SATC 225
SATC 226
SATC 227
SATC 228
SATC 229
SATC 230
SATC 231
SATC 232
SATC 233
SATC 234
SATC 235
SATC 236
SATC 237
SATC 238
SATC 239
SATC 240
SATC 241
SATC 242
SATC 243
SATC 244
SATC 245
SATC 246

NAME SATC21
PURPOSE PART 2 OF SATELLITE CLOCK CORRECTION FOR GEOS 2 (SATCL2)
CALLING SEQUENCE X=SATC21(RFFTIM)
SYMBOL TYPE DESCRIPTION
RFFTIM R INPUT - REFERENCE TIME
SATC21 P OUTPUT - CLOCK CORRECTION IN SECONDS
SUBROUTINES USED NONE
COMMON BLOCKS NONE
INPUT FILES NONE
OUTPUT FILES NONE

```

FUNCTION SATC21(RFFTIM)
  DIMENSION DAYNO(456), CLC(456), DAYNO6(114), DAYNO7(111),
  * DAYNO9(114), DAYNO8(114), CLC6(114), CLC7(111), CLC8(114),
  * CLC9(114)
  EQUIVALENCE (DAYNO(1), DAYNO(1)), (DAYNO(11), DAYNO(2)),
  * (DAYNO7(1), DAYNO7(11)), (DAYNO8(1), DAYNO(22)),
  * (DAYNO9(1), DAYNO(34)), (CLS1, CLC(1)), (CLC6(1), CLC(2)),
  * (CLC7(1), CLC(11)), (CLC8(1), CLC(22)), (CLC9(1), CLC(34))
  DATA DAYNO, CLS1/232.2979, 0.125/
  DATA DAYNO6/
  * 232.7917, 233.2554, 237.0049, 237.0912, 234.2185, 234.9191,
  * 234.9059, 235.1931, 235.2614, 235.4313, 235.8354, 235.9093,
  * 236.2750, 236.8451, 237.2112, 237.2492, 237.8543, 237.8629,
  * 238.2257, 234.3021, 238.7951, 239.2380, 239.7153, 240.8215,
  * 241.1917, 241.2553, 241.8354, 242.2742, 242.2735, 242.9486,
  * 243.2165, 243.2167, 243.7917, 247.9617, 243.9614, 244.2709, 244.3055,
  * 244.8750, 245.3187, 245.3119, 245.8987, 246.2582, 246.3326, 246.8257,
  * 246.4292, 247.2587, 247.7465, 247.9793, 248.2087, 248.2626,
  * 248.4521, 248.8555, 249.2218, 249.2958, 249.8553, 250.2333,
  * 250.3090, 250.8785, 251.2465, 251.7220, 251.8160, 252.2007,
  * 252.3731, 252.8292, 253.2729, 253.3500, 253.8433, 254.2118,
  * 254.2561, 254.7798, 254.8555, 254.8547, 255.2247, 255.2003,
  * 255.7230, 255.8587, 255.8729, 256.2377, 256.8812, 257.2505,
  * 257.4193, 257.8250, 258.2612, 258.4325, 258.9030, 259.2764,
  * 259.2158, 260.2896, 260.8547, 261.2243, 261.2027,
  * 261.5721, 262.2410, 262.2167, 262.8453, 263.7512, 263.3284,
  * 263.8213, 264.8262, 264.2573, 264.2475, 264.4702, 265.2605,
  * 265.7574, 265.8691, 266.2192, 266.5737, 266.9673, 267.2324,
  * 267.2069, 267.8755, 268.3201, 268.8130, 269.2575, 269.3733/

```

DATA DAYNO7/							SATC 701
269.5319.	270.2708.	271.7472.	272.8389.	273.2839.	274.3611.	275.3103.	SATC 702
271.9521.	272.2971.	273.3735.	274.3662.	275.2361.	276.3103.	277.3374.	SATC 703
273.4792.	274.2485.	275.7235.	276.3227.	277.2617.	278.3374.	279.3645.	SATC 704
275.3355.	276.2749.	277.3576.	278.4423.	279.2474.	280.3645.	281.3916.	SATC 705
277.3555.	278.3203.	279.3784.	280.4637.	281.2402.	282.3916.	283.4187.	SATC 706
279.3826.	280.2867.	281.2527.	282.3247.	283.2958.	284.3916.	285.4187.	SATC 707
281.3438.	282.4325.	283.3739.	284.3124.	285.3542.	286.4457.	287.4721.	SATC 708
283.3221.	284.2915.	285.3547.	286.2757.	287.3047.	288.4721.	289.4990.	SATC 709
284.3492.	285.3171.	286.3853.	287.3624.	288.2561.	289.3311.	290.4091.	SATC 710
286.3792.	287.3442.	288.4359.	289.3124.	290.3574.	291.4491.	292.4755.	SATC 711
288.3255.	289.3713.	290.4623.	291.3387.	292.3081.	293.4755.	294.4919.	SATC 712
290.3526.	291.3813.	292.4697.	293.3447.	294.3209.	295.4919.	296.5057.	SATC 713
293.3479.	294.3394.	295.4158.	296.2859.	297.3611.	298.4200.	299.4350.	SATC 714
295.3750.	296.3663.	297.3113.	298.2792.	299.3553.	300.4350.	301.4500.	SATC 715
297.3923.	298.3992.	299.3381.	300.3103.	301.3903.	302.4500.	303.4650.	SATC 716
299.3430.	300.4187.	301.3645.	302.3326.	303.3020.	304.4650.	305.4800.	SATC 717
301.3784.	302.4594.	303.3152.	304.3423.	305.4804.	306.5004.	307.5154.	SATC 718
303.3958.	304.4906.	305.3573.	306.3416.	307.4949.	308.5205.	309.5355.	SATC 719
305.3547.	306.4221.	307.3997.					SATC 720
							SATC 721
							SATC 722
							SATC 723
							SATC 724
							SATC 725
							SATC 726
							SATC 727
							SATC 728
							SATC 729
							SATC 730
							SATC 731
							SATC 732
							SATC 733
							SATC 734
							SATC 735
							SATC 736
							SATC 737
							SATC 738
							SATC 739
							SATC 740
							SATC 741
							SATC 742
							SATC 743
							SATC 744
							SATC 745
							SATC 746
							SATC 747
							SATC 748
							SATC 749
							SATC 750
							SATC 751
							SATC 752
							SATC 753
							SATC 754
							SATC 755
							SATC 756
							SATC 757
							SATC 758
							SATC 759
							SATC 760
							SATC 761
							SATC 762
							SATC 763
							SATC 764
							SATC 765
							SATC 766
							SATC 767
							SATC 768
							SATC 769
							SATC 770
							SATC 771
							SATC 772
							SATC 773
							SATC 774
							SATC 775
							SATC 776
							SATC 777
							SATC 778
							SATC 779
							SATC 780
							SATC 781
							SATC 782
							SATC 783
							SATC 784
							SATC 785
							SATC 786
							SATC 787
							SATC 788
							SATC 789
							SATC 790
							SATC 791
							SATC 792
							SATC 793
							SATC 794
							SATC 795
							SATC 796
							SATC 797
							SATC 798
							SATC 799
							SATC 800

2-2

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

•	433.5444,	434.0374,	434.4805,	434.5575,	435.0505,	435.1277,	SATC 757
•	435.4937,	436.0627,	436.4326,	436.5858,	437.0776,	437.4451,	SATC 758
•	437.5208,	438.0908,	438.4567,	438.5739,	439.1074,	439.4714,	SATC 759
•	439.5479,	455.1013,	456.4692,	456.5440,	457.0388,	457.4810/	SATC 760
	DATA CL05/						SATC 761
•	0.132,	0.152,	0.246,	0.261,	0.255,	0.276,	SATC 762
•	0.293,	-7.546,	-20.321,	-20.727,	0.118,	0.157,	SATC 763
•	0.144,	0.209,	-40.382,	-40.430,	-56.356,	0.327,	SATC 764
•	-37.320,	-40.744,	0.314,	-2.875,	-7.750,	0.012,	SATC 765
•	-0.016,	-0.010,	-0.005,	0.0	-0.030,	-0.028,	SATC 766
•	-0.024,	5.226,	6.286,	6.286,	0.324,	0.337,	SATC 767
•	0.387,	0.361,	-31.096,	0.380,	0.407,	0.358,	SATC 768
•	0.090,	0.129,	0.148,	0.120,	-1.374,	-1.389,	SATC 769
•	-1.442,	0.159,	0.026,	0.192,	0.137,	0.221,	SATC 770
•	0.239,	0.196,	0.271,	0.270,	0.275,	0.215,	SATC 771
•	0.315,	0.332,	0.352,	0.347,	0.394,	0.393,	SATC 772
•	0.409,	0.418,	0.438,	0.269,	0.253,	0.257,	SATC 773
•	0.229,	0.218,	-0.119,	-0.100,	-0.170,	-0.190,	SATC 774
•	-7.163,	-0.200,	-0.153,	-0.126,	-0.142,	-0.192,	SATC 775
•	-0.197,	-0.252,	-0.209,	-0.211,	-0.251,	-0.234,	SATC 776
•	-0.251,	-0.257,	-0.267,	-0.264,	-0.390,	-0.370,	SATC 777
•	-1.409,	0.384,	0.359,	0.362,	0.291,	0.258,	SATC 778
•	0.279,	0.212,	0.115,	0.106,	0.058,	0.083,	SATC 779
•	0.119,	0.069,	0.035,	0.025,	0.023,	-0.011/	SATC 780
	DATA CL07/						SATC 781
•	-1.049,	-0.072,	-0.090,	-0.105,	-0.125,	-0.128,	SATC 782
•	-0.128,	-0.149,	-0.137,	-0.157,	-0.161,	-0.157,	SATC 783
•	-0.143,	-0.147,	-0.148,	-0.178,	-0.152,	-0.163,	SATC 784
•	-0.205,	-0.184,	-0.188,	-0.224,	-0.237,	-0.257,	SATC 785
•	-0.276,	-0.302,	-0.276,	-0.351,	-0.327,	-0.310,	SATC 786
•	-0.382,	0.229,	0.260,	0.285,	0.224,	-1.237,	SATC 787
•	-1.115,	-1.104,	-1.122,	0.259,	0.230,	0.215,	SATC 788
•	0.317,	0.272,	0.293,	0.279,	0.259,	0.209,	SATC 789
•	0.218,	0.165,	0.168,	0.212,	0.137,	0.135,	SATC 790
•	0.150,	0.094,	0.140,	0.145,	0.111,	0.138,	SATC 791
•	0.119,	0.088,	0.152,	0.137,	0.089,	0.105,	SATC 792
•	0.117,	0.046,	0.065,	0.054,	0.017,	0.010,	SATC 793
•	-0.010,	-0.054,	-0.054,	-0.007,	0.025,	-0.012,	SATC 794
•	-0.005,	-0.032,	-0.011,	-0.035,	-0.015,	-0.023,	SATC 795
•	-0.015,	-0.064,	-0.062,	-0.125,	-0.074,	-0.073,	SATC 796
•	-0.123,	-0.111,	-0.163,	-0.160,	-0.157,	-0.125,	SATC 797
•	-0.122,	-0.176,	-0.168,	-0.147,	-0.154,	-0.190,	SATC 798
•	-0.175,	-1.041,	-1.066,	-1.040,	-1.170,	-1.105,	SATC 799
•	-1.132,	-1.168,	0.335/				SATC 800
	DATA CL04/						SATC 801
•	0.162,	0.168,	0.027,	0.150,	0.137,	0.147,	SATC 802
•	0.123,	0.175,	0.193,	0.169,	0.202,	0.185,	SATC 803
•	0.205,	0.175,	0.232,	0.188,	0.195,	0.213,	SATC 804
•	0.252,	0.193,	0.238,	0.227,	0.231,	0.277,	SATC 805
•	0.225,	0.209,	0.256,	0.303,	0.305,	0.288,	SATC 806
•	0.355,	0.174,	0.190,	0.143,	0.179,	0.090,	SATC 807
•	0.098,	0.119,	0.150,	0.206,	0.238,	0.270,	SATC 808
•	0.211,	0.147,	0.154,	0.112,	0.125,	0.021,	SATC 809
•	0.211,	0.052,	0.010,	0.006,	0.016,	0.042,	SATC 810
•	0.051,	0.058,	0.070,	0.090,	0.102,	0.112,	SATC 811
•	0.126,	0.132,	0.150,	0.172,	0.125,	0.205,	SATC 812

•	0.195.	0.192.	0.200.	0.199.	0.215.	0.212.	SATC 413
•	0.228.	0.233.	0.249.	0.240.	0.261.	0.258.	SATC 414
•	0.085.	0.091.	0.106.	0.108.	0.127.	0.152.	SATC 415
•	0.140.	0.170.	0.181.	0.205.	0.221.	0.226.	SATC 416
•	0.256.	0.264.	0.235.	0.246.	0.309.	0.253.	SATC 417
•	0.260.	0.310.	0.112.	0.044.	0.139.	0.158.	SATC 418
•	0.164.	0.098.	0.190.	0.204.	0.223.	0.241.	SATC 419
•	0.257.	0.277.	0.285.	0.273.	0.297.	0.291/	SATC 420
DATA CLC7/							SATC 421
•	0.297.	0.006.	0.019.	0.027.	0.023.	0.024.	SATC 422
•	0.043.	0.043.	0.040.	0.040.	0.040.	0.054.	SATC 423
•	0.078.	0.092.	0.102.	0.091.	0.117.	0.114.	SATC 424
•	0.119.	0.104.	0.110.	0.108.	0.132.	0.095.	SATC 425
•	0.089.	0.090.	0.090.	0.075.	0.074.	0.075.	SATC 426
•	0.059.	0.069.	0.055.	0.052.	0.058.	0.053.	SATC 427
•	0.060.	0.076.	0.078.	0.071.	0.091.	0.091.	SATC 428
•	0.090.	0.084.	0.090.	0.081.	0.092.	0.093.	SATC 429
•	0.088.	0.100.	0.087.	0.101.	0.103.	0.054.	SATC 430
•	0.076.	0.063.	0.055.	0.087.	0.112.	0.120.	SATC 431
•	0.165.	0.161.	0.175.	0.173.	0.274.	0.223.	SATC 432
•	0.222.	0.238.	0.254.	0.252.	0.275.	0.292.	SATC 433
•	0.237.	0.305.	0.309.	0.232.	0.344.	0.240.	SATC 434
•	0.148.	0.113.	0.110.	0.105.	0.090.	0.090.	SATC 435
•	0.077.	0.062.	0.047.	0.045.	0.099.	0.023.	SATC 436
•	0.017.	0.003.	0.005.	0.0	0.012.	0.005.	SATC 437
•	0.018.	0.049.	0.081.	0.077.	0.102.	0.141.	SATC 438
•	0.135.	0.170.	0.200.	0.202.	0.228.	0.250.	SATC 439
•	0.243.	-0.358.	-0.389.	-0.392.	-0.444.	-0.481/	SATC 440
DATA NCLOCK/4567							SATC 441
DO 10 I=2,NCLOCK							SATC 442
IF (REFTIM.GT.DAYNO(I)) GT TO 10							SATC 443
C INTERPLATE							SATC 444
RATE=(CLC(I)-CLC(I-1))/(DAYNO(I)-DAYNO(I-1))							SATC 445
SATC21=(RATE*(REFTIM-DAYNO(I-1))+CLC(I-1))*1.E-3							SATC 446
RETURN							SATC 447
1: CONTINUE							SATC 448
FFURN							SATC 449
END							SATC 450

NAME SATC22

PURPOSE PART 3 OF SATELLITE CLOCK CORRECTION FOR GEOS 2 (SATCL3)

CALLING SEQUENCE X=SATC22(PEFTIM)

SYMBOL TYPE DESCRIPTION

PEFTIM R INPUT - REFERENCE TIME

SATC22 F OUTPUT - CLOCK CORRECTION IN SECONDS

SUBROUTINES USED NONE

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE

FUNCTION SATC22(PEFTIM)	SATC 475
DIMENSION DAYNO(489), CLC(489), DAYN10(114), DAYN11(114),	SATC 476
DAYN12(114), DAYN13(114), DAYN14(32), CL10(114), CL11(114),	SATC 477
CL12(114), CL13(114), CL14(32)	SATC 478
EQUIVALENCE (DAYS2, DAYNO(1)), (DAYN10(1), DAYNO(2)),	SATC 479
(DAYN11(1), DAYNO(116)), (DAYN12(1), DAYNO(232)),	SATC 480
(DAYN13(1), DAYNO(348)), (DAYN14(1), DAYNO(464)),	SATC 481
(CL10, CLC(1)), (CL12(1), CLC(2)), (CL11(1), CLC(116)),	SATC 482
(CL12(1), CLC(232)), (CL13(1), CLC(348)), (CL14(1), CLC(464))	SATC 483
DATA DAYS2, CLS2/457.4819, -0.4817	SATC 484
DATA DAYN10/	SATC 485
457.4373, 457.5581, 458.0520, 458.4956, 458.5720, 459.1416,	SATC 486
459.5083, 459.5859, 460.0794, 460.4478, 460.5215, 460.5269,	SATC 487
461.0216, 461.4602, 461.5354, 462.0291, 462.4734, 462.5486,	SATC 488
463.1179, 463.4861, 463.5518, 464.1318, 464.4993, 464.5757,	SATC 489
465.0686, 465.5125, 465.5901, 466.0818, 466.1532, 466.5256,	SATC 490
467.1721, 467.4644, 467.5389, 468.1032, 468.4771, 468.5520,	SATC 491
469.0457, 469.0437, 469.0559, 470.1353, 470.5077, 470.5701,	SATC 492
471.0720, 471.1484, 471.5159, 472.0452, 472.5291, 472.6069,	SATC 493
473.0784, 473.4709, 473.5422, 474.1116, 474.4912, 474.5554,	SATC 494
475.1250, 475.4937, 475.5503, 476.1287, 476.5068, 476.5825,	SATC 495
477.0757, 477.1519, 477.5210, 477.5942, 478.0849, 478.5325,	SATC 496
479.4104, 479.1021, 479.4732, 479.5657, 480.1152, 480.4546,	SATC 497
481.5496, 481.1254, 481.5776, 482.1424, 482.5123, 482.5151,	SATC 498
483.5859, 483.1791, 483.1575, 483.5734, 484.0735, 484.1687,	SATC 499
485.3366, 484.6139, 485.1055, 485.4756, 485.5408, 485.6211,	SATC 500
486.1137, 486.6988, 486.5530, 487.1319, 487.5012, 487.5762,	SATC 501
488.1050, 488.5144, 488.6701, 489.1599, 489.5249, 489.6073,	SATC 502
490.0957, 490.1721, 490.5470, 491.1392, 491.1453, 491.5532,	SATC 503

•	492.1221.	492.1992.	492.5667.	493.1353.	493.5046.	493.5798/	SATC 504
	DATA DAYN11/						SATC 505
•	494.0728.	494.1514.	494.5935.	495.0959.	495.1538.	495.5313.	SATC 506
•	495.1753.	495.5137.	496.6206.	497.1123.	497.1927.	497.5560.	SATC 507
•	497.5095.	498.1255.	498.2019.	498.4963.	498.5701.	499.1387.	SATC 508
•	499.5098.	499.5940.	500.1519.	500.5215.	500.5972.	501.0994.	SATC 509
•	501.1650.	501.5347.	502.1025.	502.5479.	502.6243.	503.1160.	SATC 510
•	503.1943.	503.5510.	504.1222.	504.2075.	504.5742.	505.1423.	SATC 511
•	505.5125.	505.5874.	506.0798.	506.1555.	506.6006.	507.0930.	SATC 512
•	507.1687.	507.5145.	508.1810.	508.5513.	508.6277.	509.1187.	SATC 513
•	509.1958.	509.5652.	510.1318.	510.2090.	510.5734.	511.1459.	SATC 514
•	511.5166.	511.5908.	512.1532.	512.1589.	512.5213.	512.5040.	SATC 515
•	512.5006.	513.1721.	513.5422.	513.6179.	513.6145.	514.1553.	SATC 516
•	514.5547.	514.6311.	515.5679.	516.1353.	516.2124.	516.5796.	SATC 517
•	516.5762.	517.3574.	517.7214.	517.7969.	517.3699.	517.7346.	SATC 518
•	517.8101.	517.3933.	517.7471.	517.9240.	517.3965.	517.7603.	SATC 519
•	517.8379.	517.3302.	517.6997.	517.7742.	517.3435.	517.7124.	SATC 520
•	517.7974.	517.2917.	517.7249.	517.8018.	517.2942.	517.7400.	SATC 521
•	517.7074.	517.7512.	517.8274.	517.3977.	517.7455.	517.8442.	SATC 522
•	517.3337.	517.7061.	517.7709.	517.3450.	517.7185.	517.7009.	SATC 523
•	517.2854.	517.3506.	517.8040.	517.3777.	517.7449.	517.8179/	SATC 524
	DATA DAYN12/						SATC 525
•	518.3108.	518.7546.	518.8311.	518.3240.	518.4014.	518.7676.	SATC 526
•	518.3374.	518.7273.	518.7813.	518.3506.	518.7209.	518.7042.	SATC 527
•	518.3645.	518.7324.	518.8081.	518.7456.	518.8213.	518.3909.	SATC 528
•	518.7533.	518.7715.	518.9484.	518.7408.	518.7107.	518.7847.	SATC 529
•	518.3540.	518.7261.	518.7383.	518.4679.	518.7305.	518.8115.	SATC 530
•	518.3911.	518.7400.	518.8247.	518.7179.	518.7425.	518.9395.	SATC 531
•	518.3311.	518.7256.	518.3442.	518.7383.	518.8657.	518.3574.	SATC 532
•	518.3019.	518.3713.	518.7400.	518.3149.	518.3142.	518.7534.	SATC 533
•	518.3281.	518.4026.	518.7559.	518.4420.	518.3345.	518.4109.	SATC 534
•	518.7791.	518.3477.	518.7929.	518.4591.	518.3508.	518.4380.	SATC 535
•	518.3066.	518.3740.	518.7471.	518.4134.	518.3123.	518.7588.	SATC 536
•	518.3503.	518.3247.	518.4014.	518.7749.	518.3379.	518.4146.	SATC 537
•	518.7454.	518.4275.	518.7293.	518.3545.	518.4414.	518.8101.	SATC 538
•	518.7777.	518.7476.	518.4228.	518.3904.	518.7632.	518.9350.	SATC 539
•	518.3241.	518.4048.	518.8491.	518.3416.	518.4177.	518.7866.	SATC 540
•	518.3547.	518.4309.	518.7998.	518.3673.	518.4456.	518.4130.	SATC 541
•	518.3811.	518.4583.	518.3262.	518.4947.	518.7655.	518.4394.	SATC 542
•	518.3811.	518.4375.	518.6525.	518.3450.	518.4211.	518.7900.	SATC 543
•	518.3582.	518.8025.	518.3713.	518.4478.	518.8179.	518.3845/	SATC 544
	DATA DAYN13/						SATC 545
•	518.4517.	518.8296.	518.3225.	518.7708.	518.8428.	518.4109.	SATC 546
•	518.7513.	518.4560.	518.3494.	518.4253.	518.7935.	518.4414.	SATC 547
•	518.3065.	518.3338.	518.3198.	518.4382.	518.4092.	518.8333.	SATC 548
•	518.3014.	518.7722.	518.3664.	518.3790.	518.4146.	518.3521.	SATC 549
•	518.4277.	518.5005.	518.3652.	518.3123.	518.3734.	518.4548.	SATC 550
•	518.4250.	518.3916.	518.4640.	518.3531.	518.4045.	518.4812.	SATC 551
•	518.3409.	518.3423.	518.4140.	518.4519.	518.4312.	518.6027.	SATC 552
•	518.3779.	518.2579.	518.4343.	518.4145.	518.3811.	518.4575.	SATC 553
•	518.3943.	518.4714.	518.9401.	518.4067.	518.4075.	518.4533.	SATC 554
•	518.3506.	518.4214.	518.3672.	518.4745.	518.8311.	518.4476.	SATC 555
•	518.4174.	518.3345.	518.3373.	518.3777.	518.4475.	518.4100.	SATC 556
•	518.3567.	518.4341.	518.6716.	518.4616.	518.3373.	518.4512.	SATC 557
•	518.3213.	518.4680.	518.8747.	518.4014.	518.8484.	518.4146.	SATC 558
•	518.3504.	518.4277.	518.4747.	518.4434.	518.9832.	518.4575.	SATC 559

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

•	752.9319.	759.4707.	759.8331.	761.4044.	761.9506.	761.4180.	SATC 560
•	761.9638.	762.4312.	762.8770.	763.4443.	763.8909.	764.4575.	SATC 561
•	764.9048.	765.4714.	765.8430.	766.3543.	767.8672.	767.9470.	SATC 562
•	768.4346.	768.8804.	769.4473.	769.3943.	770.3253.	770.9075.	SATC 563
•	771.3984.	771.8464.	772.4116.	772.4592.	773.4248.	773.8706/	SATC 564
	DATA DAYN14/						SATC 565
•	774.4409.	774.8838.	775.4512.	775.4084.	776.4644.	776.9116.	SATC 566
•	777.3805.	777.9241.	778.4915.	778.5611.	779.4222.	779.8755.	SATC 567
•	780.4443.	780.8875.	781.4542.	781.3038.	782.4717.	782.9150.	SATC 568
•	787.4039.	783.9277.	784.4972.	784.3560.	785.4458.	785.8792.	SATC 569
•	786.4478.	786.8909.	787.4617.	787.3041.	788.4714.	788.9172.	SATC 570
•	789.4495.	789.8919/					SATC 571
	DATA CL10/						SATC 572
•	-0.132.	-0.124.	-0.178.	-0.218.	-0.215.	-0.257.	SATC 577
•	-0.305.	-0.300.	0.349.	-0.473.	-0.375.	0.050.	SATC 574
•	0.015.	-0.024.	-0.027.	-0.074.	-0.104.	-0.100.	SATC 575
•	-0.139.	-0.190.	-0.184.	-0.322.	-0.264.	-0.258.	SATC 576
•	-0.316.	-0.352.	-0.346.	-0.371.	-0.385.	-0.420.	SATC 577
•	-0.459.	-0.485.	-0.490.	-0.494.	-0.557.	-0.555.	SATC 578
•	0.165.	0.122.	0.099.	0.060.	0.035.	0.031.	SATC 579
•	-0.012.	-0.015.	-0.048.	-0.096.	-0.134.	-0.138.	SATC 580
•	-0.176.	-0.210.	-0.211.	-0.245.	-0.277.	-0.277.	SATC 581
•	-0.313.	-0.343.	-0.346.	-0.377.	-0.431.	-0.415.	SATC 582
•	-0.459.	-0.456.	-0.473.	0.306.	0.291.	0.276.	SATC 583
•	0.271.	0.254.	0.236.	0.233.	0.211.	0.193.	SATC 584
•	0.196.	0.182.	-0.270.	-0.203.	-0.209.	0.473.	SATC 585
•	0.477.	0.468.	0.466.	0.464.	0.468.	0.460.	SATC 586
•	0.454.	0.461.	0.477.	0.422.	0.423.	0.434.	SATC 587
•	0.422.	0.411.	0.360.	0.221.	0.210.	0.211.	SATC 588
•	0.190.	0.175.	0.179.	0.178.	0.163.	0.176.	SATC 589
•	0.167.	0.168.	0.164.	0.169.	0.172.	0.172.	SATC 590
•	0.178.	0.180.	0.184.	0.183.	0.183.	0.197/	SATC 591
	DATA CL11/						SATC 592
•	0.194.	0.199.	0.203.	0.204.	0.215.	0.216.	SATC 593
•	0.232.	0.205.	0.212.	0.220.	0.227.	0.230.	SATC 594
•	0.237.	0.245.	0.240.	0.247.	0.256.	0.252.	SATC 595
•	0.244.	0.250.	0.226.	0.218.	0.224.	0.220.	SATC 596
•	0.228.	0.221.	0.229.	0.232.	0.242.	0.240.	SATC 597
•	0.240.	0.249.	0.260.	0.257.	0.254.	0.280.	SATC 598
•	0.279.	0.280.	0.323.	0.314.	0.324.	0.251.	SATC 599
•	0.261.	0.254.	0.249.	0.246.	0.249.	0.238.	SATC 600
•	0.237.	0.234.	0.233.	0.234.	0.275.	0.230.	SATC 601
•	0.272.	0.234.	0.234.	-8.900.	-8.910.	-8.920.	SATC 602
•	0.352.	0.334.	0.319.	0.322.	-0.040.	-0.040.	SATC 603
•	-0.043.	0.0	-0.324.	-0.322.	-0.322.	-0.324.	SATC 604
•	-0.021.	-0.139.	-0.150.	-0.141.	-0.135.	-0.146.	SATC 605
•	-0.139.	-0.152.	-0.144.	-0.138.	-0.131.	-0.127.	SATC 606
•	-0.135.	-0.151.	-0.150.	-0.147.	-0.142.	0.0	SATC 607
•	-0.155.	-0.163.	-0.172.	-0.204.	-0.204.	-0.203.	SATC 608
•	-0.197.	-0.203.	-0.189.	-0.184.	-0.177.	-0.185.	SATC 609
•	-0.182.	-0.187.	-0.184.	-0.181.	-0.177.	-0.185.	SATC 610
•	-0.174.	-0.187.	-0.203.	-0.215.	-0.211.	-0.215/	SATC 611
	DATA CL12/						SATC 612
•	-0.224.	0.0	-0.231.	-0.230.	-0.237.	-0.240.	SATC 613
•	-0.229.	-0.227.	-0.246.	-0.241.	0.122.	0.103.	SATC 614
•	0.138.	0.160.	0.176.	0.164.	0.140.	0.166.	SATC 615

•	0.162.	0.152.	0.142.	0.145.	0.154.	0.157.	SATC 616
•	0.163.	0.135.	0.145.	0.184.	0.141.	0.140.	SATC 617
•	0.145.	0.124.	0.126.	0.103.	0.077.	0.079.	SATC 618
•	0.075.	0.083.	0.095.	0.095.	0.079.	0.107.	SATC 619
•	0.196.	0.114.	0.113.	0.112.	0.0	0.0	SATC 620
•	-0.145.	-0.050.	-0.055.	-0.047.	-0.055.	-0.070.	SATC 621
•	-0.070.	-0.075.	-0.083.	-0.085.	-0.083.	-0.090.	SATC 622
•	-0.095.	-0.100.	-0.110.	-0.110.	-0.115.	-0.116.	SATC 623
•	-0.107.	-0.107.	-0.118.	-0.105.	-0.125.	-0.132.	SATC 624
•	-0.147.	-0.149.	-0.164.	-0.175.	-0.174.	-0.184.	SATC 625
•	-0.196.	-0.194.	-0.211.	-0.210.	-0.224.	-0.230.	SATC 626
•	-0.232.	-0.225.	-0.040.	-0.025.	-0.035.	-0.036.	SATC 627
•	-0.064.	-0.081.	-0.097.	-0.097.	-0.110.	-0.111.	SATC 628
•	-0.121.	-0.132.	-0.133.	-0.144.	-0.168.	-0.173.	SATC 629
•	-0.186.	-0.176.	0.240.	0.229.	0.236.	0.226.	SATC 630
•	0.210.	0.194.	0.192.	0.196.	0.175.	0.179/	SATC 631
•	DATA CL13/						SATC 632
•	0.171.	0.165.	0.154.	0.144.	0.149.	0.144.	SATC 633
•	0.133.	0.134.	0.132.	0.135.	0.125.	0.116.	SATC 634
•	0.117.	0.110.	0.092.	0.099.	0.094.	0.090.	SATC 635
•	0.081.	0.066.	0.065.	0.070.	0.083.	0.053.	SATC 636
•	0.057.	0.034.	0.045.	0.024.	0.031.	0.079.	SATC 637
•	0.031.	0.050.	0.032.	0.032.	0.015.	0.002.	SATC 638
•	0.027.	0.010.	0.017.	0.017.	0.009.	0.010.	SATC 639
•	0.028.	0.018.	0.013.	0.002.	0.015.	0.015.	SATC 640
•	0.023.	0.015.	0.009.	0.020.	4.730.	4.740.	SATC 641
•	0.120.	0.124.	0.115.	0.116.	0.124.	0.127.	SATC 642
•	0.116.	0.125.	0.115.	0.119.	0.109.	0.122.	SATC 643
•	0.122.	0.124.	0.118.	0.124.	0.112.	0.109.	SATC 644
•	0.106.	0.097.	0.095.	0.103.	0.095.	0.090.	SATC 645
•	0.0	0.095.	0.089.	0.097.	0.081.	0.066.	SATC 646
•	0.055.	0.056.	0.043.	0.071.	0.037.	0.037.	SATC 647
•	0.029.	0.037.	0.028.	0.030.	0.029.	0.028.	SATC 648
•	0.024.	0.023.	0.029.	0.033.	0.035.	0.024.	SATC 649
•	0.350.	0.350.	0.305.	0.309.	0.304.	0.325.	SATC 650
•	0.324.	0.307.	0.316.	-0.093.	-0.092.	-0.094/	SATC 651
•	DATA CL13/						SATC 652
•	-0.092.	-0.100.	-0.091.	-0.090.	-0.094.	-0.092.	SATC 653
•	-0.089.	-0.090.	-0.079.	-0.091.	-0.083.	-0.091.	SATC 654
•	-0.091.	-0.053.	-0.101.	-0.104.	-0.105.	-0.115.	SATC 655
•	-0.120.	-0.120.	-0.096.	-0.130.	-0.130.	-0.143.	SATC 656
•	-0.132.	-0.125.	-0.117.	-0.123.	-0.118.	-0.118.	SATC 657
•	-0.126.	-0.113/					SATC 658
•	DATA NCLOCK/489/						SATC 659
•	DJ 13 I=3,NCLOCK						SATC 660
•	IF(PRETIM.GT.DAYNO(I)) GO TO 10						SATC 661
•	C INTERPOLATE						SATC 662
•	RATE=(CLC(I)-CLC(I-1))/(DAYNO(I)-DAYNO(I-1))						SATC 663
•	SATC22=(RATE*(PRETIM-DAYNO(I-1))+CLC(I-1))*1.E-3						SATC 664
•	RETURN						SATC 665
•	10 CONTINUE						SATC 666
•	RETURN						SATC 667
•	END						SATC 668

NAME SIMRD

PURPOSE TO READ SIMULATED DATA TAPES

CALLING SEQUENCE CALL SIMRD(NSTARD)

SYMBOL TYPE DESCRIPTION

NSTARD I INPUT-- NUMBER OF STATIONS THAT WERE READ

SUBROUTINES USED RANDWR TDIF YMDAY NUMBR2 BIAS

COMMON BLOCKS APARAM CEPHEM CGEOS CONSTS CPARAM
CSTINF CTIME INTBLK PREBLK SIGBLK
STANUM

INPUT FILES IOBS - DATA TAPE

OUTPUT FILES PRINTER

REFERENCE 'GEODYN PROGRAM OPERATIONS DESCRIPTION' - APPENDIX C
VOLUME 3 - GEODYN DOCUMENTATION

```

SUBROUTINE SIMRD(NSTARD)
  IMPLICIT REAL*8 (A-H,O-Z)
  LOGICAL*4 OKSAT,VHFCHN,PREPRC
  LOGICAL NOFATE,SATSAT
  INTEGER*2 CULL,CHANL,NVEAS,MTYPE,PRETYR,INTYPL,ISTAC,ISATNO,
  ITYPE,ESTANO,ISTARD,ISTANO,STANCS,JBASE,KBASE
  INTEGER RECON
  REAL TCIF,SIGSTO,DAYINT,SIGCHG,SGPRNT
  DOUBLE PRECISION JNAME,NAM
  DIMENSION SIG(2)
  COMMON/APARAM/INPAR,INPAR1,NSIAS,NSTSTA,NSAT,NGPARC(5)
  COMMON/CEPHEM/JNAME(361),ISTARC(361),ESTANO(361),ISTANO(366)
  COMMON/CGEOS/ISAT2(2),IPRPR(250),NPRE,NSIG,NCULL,SIGCHG(50),
  INTYPE(50),ISTNO(50),CULL(2,100)
  COMMON/CONSTS/OPI,BTWCP1,D2F,S2R
  COMMON/CPARAM/NSTA,NMAST(15)
  COMMON/CSTINF/JBASE(283),KBASE(283),LBASE
  COMMON/CTIME/LATAEP,DAYREF,CAYE,CAYSTR,CAYINT(15)
  COMMON/INTBLK/INTBK1(53),NOFATE,INTBK2(78)
  COMMON/PREBLK/DAY,CES1,CES2,SIG1,SIG2,SFFNDX,ISTA,MTYPE,NVEAS,
  ISATNO,PRETYR,CHANL,VHFCHN,PREPRC,RECON
  COMMON/SIGBLK/SIGSTO(30),SGPRNT(30),IOBS,IOAPE(3)
  COMMON/STANUM/NAM(280),STANCS(220),NCSTOF
  EQUIVALENCE (SIG(1),SIG1),(KKSAT,SIG2)
  AITIME(DAY)=TCIF(4,3,2AY)/8.64E4
C INITIALIZE
  IF(IOBS.EQ.0) IOBS=21

```

SIMR 27
SIMR 28
SIMR 29
SIMR 30
SIMR 31
SIMR 32
SIMR 33
SIMR 34
SIMR 35
SIMR 36
SIMR 37
SIMR 38
SIMR 39
SIMR 40
SIMR 41
SIMR 42
SIMR 43
SIMR 44
SIMR 45
SIMR 46
SIMR 47
SIMR 48
SIMR 49
SIMR 50
SIMR 51
SIMR 52
SIMR 53

```

NN1=0
NN2=0
NSTAT=FALSE
P=1
VHFCH=0
VFCH=0
C 30 31-31 1000
10 30 31 (IF (C.DES.1=SEC.1) IYMD,IMH,SEC,C.DES.1,C.DES.2,SIG1,SIG2,ISN,
  MTYPE,NO.2,NO.3,NTYPE,NO.1,NO.2,NO.3) GO TO 30
  IF (ISN.EQ.0) GO TO 30
  GO TO 30
C 32 32 OPERATIONS
20 32 31 (IF (C.DES.1=SEC.1) IYMD,IMH,SEC,C.DES.1,C.DES.2,SIG1,SIG2,ISN,
  MTYPE,NO.2,NO.3,NTYPE,NO.1,NO.2,NO.3) GO TO 32
  IF (ISN.EQ.0) GO TO 32
  GO TO 32
C 33 33 CHECK SATELLITE
30 33 31=FALSE
  DO 40 I=1,NSAT
    IF (LKSATI) GO TO 40
    ISATNO=I
    OKSAT=ISAT1.EQ.1.CATE2(I)
  40 CONTINUE
    IF (.NOT.OKSAT) GO TO 20
C 34 34 CHECK SECOND SATELLITE IF SAT-SAT TRACKING
  IF (NTYPE.NO.2.N.MTYPE.NO.3) GO TO 48
  SATSAT=KKSAT.GT.0
  IF (.NOT.SATSAT) GO TO 48
  IF (NSAT.LT.2) GO TO 20
  OKSAT=FALSE
  DO 45 I=1,NSAT
    OKSAT=KKSAT.EQ.ISAT2(I)
    IF (.NOT.OKSAT) GO TO 45
    KKSAT=I
  45 CONTINUE
  GO TO 48
  48 NN1=0
  NN2=0
  IF (NSIG.LE.0) GO TO 60
C 35 35 CHECK FOR SIG44 CHANGE
  DO 50 I=1,NSIG
    IF (ISN.NO.1,ISTNO(I).AND.1STNO(I).NO.0) GO TO 50
    IF (NTYPE.EQ.1,NTYPE(I).OR.1NTYPE(I).EQ.0) NN1=I
    IF (NTYPE.GT.7) GO TO 50
    IF (NTYPE+7.EQ.1,NTYPE(I).OR.1NTYPE(I).EQ.0) NN2=I
  50 CONTINUE
  60 ISTA=ISN
C 36 36 COMPUTE TIME IN A.1 DAYS FROM JAN 0.0
  DAY=YMDAY(IYMD,IMH,SEC)
  DAY=DAY+ATIME(DAY)
  IF (DAY.LT.0) GO TO 20
  IF (DAY.GT.0) GO TO 200
C 37 37 CHECK FOR STATION PRESENT
  IF (ISN.EQ.0) GO TO 100
  ISN=NUMB2(ISTK,ISTANG,NSTA)
  IF (ISN.GT.0) GO TO 90
  ISN=NUMB2(ISTA,ISTARD,NSTARD)
  IF (ISN.GT.0) GO TO 80

```



SIMR 41
SIMR 42
SIMR 43
SIMR 44
SIMR 45
SIMR 46
SIMR 47
SIMR 48
SIMR 49
SIMR 50
SIMR 51
SIMR 52
SIMR 53
SIMR 54
SIMR 55
SIMR 56
SIMR 57
SIMR 58
SIMR 59
SIMR 60
SIMR 61
SIMR 62
SIMR 63
SIMR 64
SIMR 65
SIMR 66
SIMR 67
SIMR 68
SIMR 69
SIMR 70
SIMR 71
SIMR 72
SIMR 73
SIMR 74
SIMR 75
SIMR 76
SIMR 77
SIMR 78
SIMR 79
SIMR 80
SIMR 81
SIMR 82
SIMR 83
SIMR 84
SIMR 85
SIMR 86
SIMR 87
SIMR 88
SIMR 89
SIMR 90
SIMR 91
SIMR 92
SIMR 93
SIMR 94
SIMR 95
SIMR 96
SIMR 97
SIMR 98
SIMR 99
SIMR 100
SIMR 101
SIMR 102
SIMR 103
SIMR 104
SIMR 105
SIMR 106
SIMR 107
SIMR 108
SIMR 109



REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR


```
ISN2=NY (ISTAR,ISTARS,NOSTAR) SIMR 110
IF (ISN2.GT.0) GO TO 111 SIMR 111
IF (ISTAR.GT.0) GO TO 112 SIMR 112
PRINT 10,ISTAR SIMR 113
GO TO 114 SIMR 114
6. NSTAR=ISTAR SIMR 115
ISTAR=ISTAR SIMR 116
ISN2=0 SIMR 117
90 ISTAR=0 SIMR 118
IF (NTYP.LT.27) GO TO 100 SIMR 119
C CHECK FOR TRANSMITTING STATION IF VALUE OF THE DATA SIMR 120
ISTAR=0 SIMR 121
ISN2=CUMUL(ISTAR,ISTANC,NSTA) SIMR 122
IF (ISN2.GT.0) GO TO 94 SIMR 123
IGI2=NUM(0,ISTAR,ISTAR-NSTAR) SIMR 124
IF (ISN2.GT.0) GO TO 92 SIMR 125
ISN2=NUM(0,ISTAR,ISTARS,NOSTAR) SIMR 126
IF (ISN2.GT.0) GO TO 92 SIMR 127
PRINT 10,ISTAR SIMR 128
GO TO 20 SIMR 129
92 NSTAR=NSTAR+1 SIMR 130
ISN2=NSTAR SIMR 131
ISTANC(ISTAR)=ISTAR SIMR 132
94 CHANNEL=ISTAR SIMR 133
IF (LEAS.LT.0) GO TO 95 SIMR 134
GO TO LEAS SIMR 135
IF (ISN2.NE.UBAS(1)) GO TO 96 SIMR 136
IF (ISN2.NE.KRAS(1)) GO TO 96 SIMR 137
GO TO 100 SIMR 138
96 CONTINUE SIMR 139
98 LEAS=LEAS+1 SIMR 140
UBAS(LEAS)=ISN SIMR 141
KRAS(LEAS)=ISN2 SIMR 142
100 IF (NN1.LT.0) GO TO 150 SIMR 143
C CONVERT MEASUREMENT SIGMAS TO PROPER UNITS SIMR 144
IF (NTYP.LT.14) GO TO 300 SIMR 145
GO TO (110,115,120,115,130,135,110),NTYPE SIMR 146
110 SIG1=SIGCHG(NN1)*S2R/DCCS(OBS2) SIMR 147
GO TO 150 SIMR 148
115 SIG1=SIGCHG(NN1) SIMR 149
GO TO 150 SIMR 150
120 SIG1=0.0100*SIGCHG(NN1) SIMR 151
GO TO 150 SIMR 152
130 SIG1=OBS1**2 SIMR 153
IF (SIG1.LT.1.000) SIG1=DSQRT(1.000-SIG1) SIMR 154
SIG1=SIGCHG(NN1)*1.00-3/SIG1 SIMR 155
GO TO 150 SIMR 156
135 SIG1=SIGCHG(NN1)*S2R SIMR 157
150 IF (NN2.LT.0) GO TO 170 SIMR 158
GO TO (155,170,170,150,165,155,155),NTYPE SIMR 159
155 SIG2=SIGCHG(NN2)*S2R SIMR 160
GO TO 170 SIMR 161
160 SIG2=SIGCHG(NN2)*0.0100 SIMR 162
GO TO 170 SIMR 163
165 SIG2=OBS2**2 SIMR 164
IF (SIG2.LT.1.000) SIG2=DSQRT(1.000-SIG2) SIMR 165
```

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

SIG=SIGCH4(NM)PI.0-178132	SIMR 184
GO TO 175	SIMR 187
180 MTYPE=C	SIMR 188
IF(NM=0)GO TO 27	SIMR 189
GO TO (170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200,201,202,203,204,205)	SIMR 190
192	SIMR 191
315 SIG=SIGCH4(NM)PI.0-178132	SIMR 192
GO TO 175	SIMR 193
320 SIG=SIGCH4(NM)PI.0-178132	SIMR 194
GO TO 175	SIMR 195
325 SIG=SIGCH4(NM)PI.0-178132	SIMR 196
GO TO 175	SIMR 197
327 SIG=SIGCH4(NM)PI.0-178132	SIMR 198
175 IF(NCULL=0) GO TO 198	SIMR 199
C CHECK FOR OBSERVATIONS COLLECTED	SIMR 200
GO 185 J=1,NM	SIMR 201
GO 185 J=1,NM	SIMR 202
IF(NUMBER+J-CULL(1,1)) 185,186,175	SIMR 203
175 IF(NUMBER+J-CULL(2,1)) GO TO 198	SIMR 204
180 SIG(J)=0.0	SIMR 205
181 CONTINUE	SIMR 206
185 NUMP=NUMP+NM	SIMR 207
RECNO=RECNO+1	SIMR 208
C CHECK FOR STAGES	SIMR 209
IF(NM=0) CALL DIAS	SIMR 210
C STORE OBSERVATIONS	SIMR 211
CALL PANWR	SIMR 212
GO TO 20	SIMR 213
200 MTYPE=C	SIMR 214
187	SIMR 215
C LAST RECORD STORE TYPE AND STATION ZERO	SIMR 216
RECNO=RECNO+1	SIMR 217
CALL PANWR	SIMR 218
PRINT 200,NUMBER,IGBS	SIMR 219
DAYSTRDAY	SIMR 220
RETURN	SIMR 221
1000 FORMAT(1X,'STATION ',I4,' NOT FOUND IN FILE')	SIMR 222
2000 FORMAT(1H//25X,I6,' OBSERVATIONS SELECTED FROM MASTER SIMULATED',	SIMR 223
DATA TAPE NUMBER',I3)	SIMR 224
END	SIMR 225







SQUANT

DESCRIPTION

SQUANT is a subroutine which computes quantities related to the Earth-fixed station positions. The first call has a different processing than the subsequent or normal processing.

The first call processing for each station consists of

- convert ϕ , the geodetic latitude, and λ , the east longitude, to radians.
- compute the Cartesian coordinates.
- If the station is to be adjusted, compute the matrix of partial derivatives of the rectangular coordinates with respect to the geodetic spherical coordinates.
- compute the \hat{N} , \hat{Z} , and \hat{E} with vectors describing the horizontal topocentric coordinate system.

Subsequent processing consists

- converting the adjusted rectangular station positions to spherical geodetic coordinates, and their covariances as well (using subroutine PLHOUT), and
- recompute the \hat{N} , \hat{Z} , and \hat{E} unit vectors on the basis of the new (adjusted) positions.

NAME SQUANT
 ENTRY POINT PURPOSE
 SQUANT: INITIAL IZATION
 SQUANT TO CONVERT STATION POSITIONS TO X,Y,Z COORDINATE SYSTEM ON FIRST CALL AND TO PHI,LAMBDA,H COORDINATE SYSTEM ON SUBSEQUENT CALLS AND TO COMPUTE PARTIALS OF X,Y,Z WITH RESPECT TO LAMBDA, PHI,H
 CALLING SEQUENCE CALL SQUANT(H,PARTL,PLHSIG,STAXYZ,ENAT,NHAT,ZHAT,THPRIM,FLAT,FLON)

SYMBOL	TYPE	DESCRIPTION
H (1)	DP	INPUT & OUTPUT - STATION HEIGHT
PARTL (3.3.1)	H	OUTPUT - PARTIALS OF X,Y,Z WITH RESPECT TO PHI, LAMBDA, AND H
PLHSIG (3.3.1)	P	INPUT - STATION SPHERICAL COORDINATE SIGMAS
STAXYZ (3.1)	DP	INPUT - TRACKING STATION EARTH FIXED CARTESIAN COORDINATES
ENAT (3.1)	DP	OUTPUT - STATION EAST UNIT VECTOR
NHAT (3.1)	DP	OUTPUT - STATION NORTH UNIT VECTOR
ZHAT (3.1)	DP	OUTPUT - STATION LOCAL VERTICAL UNIT VECTOR
THPRIM (2.1)	DP	OUTPUT - SINE AND COSINE OF THE DIFFERENCE BETWEEN GEODETIC AND GEOCENTRIC LATITUDES
FLAT (1)	DP	INPUT & OUTPUT - TRACKING STATION LATITUDE
FLON (1)	DP	INPUT & OUTPUT - TRACKING STATION LONGITUDE

CALLING SEQUENCE CALL SQUANT(NSTA,NSTEST,FRSTIM)

SYMBOL	TYPE	DESCRIPTION
NSTA	I	INPUT - INTERNAL NUMBER OF STATIONS
NSTEST	I	INPUT - NUMBER OF ADJUSTED STATIONS

FRSTIM INPUT - SWITCH IS TRUE ON FIRST CALL TO SQUANT FOR A CASE



SUBROUTINES USED CLEAR PLMOUT
 COMMON BLOCKS INTRLK
 INPUT FILES NONE
 OUTPUT FILES NONE

SUBROUTINE SQUANT(H, PARTL, PLHSIG, STAXYZ, EPHAT, NHAT, ZHAT, THPRIM,	SQUA	70
PLAT, PLON, ISTDND)	SQUA	71
IMPLICIT REAL*8 (A-H, C-7)	SQUA	72
DIMENSION H(1), PARTL(3,3,1), PLHSIG(3,3,1), STAXYZ(3,1,1), EPHAT(3,1),	SQUA	73
NHAT(3,1), ZHAT(3,1), THPRIM(3,1), PLAT(1), PLON(1)	SQUA	74
INTEGER*2 ISTDND	SQUA	75
DIMENSION ISTDND(1)	SQUA	76
LOGICAL FRSTIM	SQUA	77
REAL PARTL, PLHSIG	SQUA	78
DOUBLE PRECISION NHAT	SQUA	79
COMMON/INTRLK/IGS(8), AH, AESQ, FLAT, FFSQ(5)	SQUA	80
RETURN	SQUA	81
ENTRY SQUANT(NSTA, NSTEST, FRSTIM)	SQUA	82
C CALCULATE CONSTANTS ON FIRST CALL	SQUA	83
IF(.NOT. FRSTIM) GO TO 10	SQUA	84
ESSQ=FLAT	SQUA	85
ESQ1=(1.00-ESSQ)**2	SQUA	86
ESQ=1.00-ESQ1	SQUA	87
CALL CLEAR(STAXYZ, 6, NSTA)	SQUA	88
CALL CLEAR(EPHAT, 6, NSTA)	SQUA	89
CALL CLEAR(NHAT, 6, NSTA)	SQUA	90
CALL CLEAR(ZHAT, 6, NSTA)	SQUA	91
CALL CLEAR(THPRIM, 4, NSTA)	SQUA	92
10 DO 100 I=1, NSTA	SQUA	93
IF(ISTDND(I).EQ.-4) GO TO 100	SQUA	94
IF(FRSTIM) GO TO 20	SQUA	95
C ON ALL CALLS AFTER FIRST, CONVERT ADJUSTED X,Y,Z TO PHI, LAMBDA, H	SQUA	96
CALL PLMOUT(STAXYZ(1, I), PARTL(1, I), PLHSIG(1, I), PLAT(1),	SQUA	97
PLON(1), H(1))	SQUA	98
C CALCULATE SINE AND COSINE OF GEODETIC LATITUDE AND EAST LONGITUDE	SQUA	99
20 SLATG=DSIN(PLAT(1))	SQUA	100
SLTG=SLATG**2	SQUA	101
CLATG=DSQRT(1.000-SLTG)	SQUA	102
SINLON=DSIN(PLON(1))	SQUA	103
COSLON=DCOS(PLON(1))	SQUA	104
H=AH/DT(1)*GDD-ESQ*SLTG	SQUA	105
PCL=CLATG*(H(1)+H)	SQUA	106
C ON FIRST CALL, CALCULATE STATION X,Y,Z	SQUA	107
IF(.NOT. FRSTIM) GO TO 10	SQUA	108
STAXYZ(1, 1)=PCL*COSLON	SQUA	109
STAXYZ(2, 1)=PCL*SINLON	SQUA	110
STAXYZ(3, 1)=CLATG*(H(1)+ESQ*H)	SQUA	111



```

C IF FIRST CALL AND STATION TO BE ADJUSTED, CALCULATE PARTIALS
  IF(I.GT.NSTEST) GO TO 4C
C INTERMEDIATE CONSTANTS
  ESOPHI=ESQ*SLATG**2
  E1PHI=1.00-ESOPHI
  CPCL=CLATG*CSLON
  CPSL=CLATG*SINLON
  ESC1P=ESQ*ESQ1
  HR=H(I)+P
C PARTIALS OF X,Y,Z WITH RESPECT TO PHI
  PARTL(1,1,1)=-SLATG*CSLON*(HR-(ESQ**2*CLATG**2)/E1PHI)
  PARTL(1,2,1)=SINLON*PARTL(1,1,1)/CSLON
  PARTL(1,3,1)=CLATG*(H(I)+ESQ**2*(1.00+ESOPHI/E1PHI))
C PARTIALS OF X,Y,Z WITH RESPECT TO LAMBDA
  PARTL(2,1,1)=-STAXYZ(2,1)
  PARTL(2,2,1)=STAXYZ(1,1)
  PARTL(2,3,1)=0.
C PARTIALS OF X,Y,Z WITH RESPECT TO HEIGHT
  PARTL(3,1,1)=CPCL
  PARTL(3,2,1)=CPSL
  PARTL(3,3,1)=SLATG
C SINE AND COSINE OF DIFFERENCE OF GEOCENTRIC & GEODETIC LATITUDES
  4C THETPR=BLAT(1)-DATAN(STAXYZ(3,1)/PCL)
  THPRIM(1,1)=COS(THETPR)
  THPRIM(2,1)=CCOS(THETPR)
C FORM UNIT VECTORS AT STATION
C ...NORTH VECTOR
  NHAT(1,1)=-SLATG*CSLON
  NHAT(2,1)=-SLATG*SINLON
  NHAT(3,1)=CLATG
C ...LOCAL VERTICAL
  ZHAT(1,1)=CLATG*CSLON
  ZHAT(2,1)=CLATG*SINLON
  ZHAT(3,1)=SLATG
C ...EAST VECTOR
  EHAT(1,1)=-SINLON
  EHAT(2,1)=CSLON
  EHAT(3,1)=0.000
100 CONTINUE
  RETURN
  END
SQUA 112
SQUA 113
SQUA 114
SQUA 115
SQUA 116
SQUA 117
SQUA 118
SQUA 119
SQUA 120
SQUA 121
SQUA 122
SQUA 123
SQUA 124
SQUA 125
SQUA 126
SQUA 127
SQUA 128
SQUA 129
SQUA 130
SQUA 131
SQUA 132
SQUA 133
SQUA 134
SQUA 135
SQUA 136
SQUA 137
SQUA 138
SQUA 139
SQUA 140
SQUA 141
SQUA 142
SQUA 143
SQUA 144
SQUA 145
SQUA 146
SQUA 147
SQUA 148
SQUA 149
SQUA 150
SQUA 151
SQUA 152

```



REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR



STAINF

DESCRIPTION

Subroutine STAINF computes the statistical information derived from the measurement residuals. These statistics are primarily for printout purposes, being composed of those statistics in the residual summary printout. In addition, STAINF computes the weighted RMS considering the degrees of freedom removed due to the regression.

There are four entries; their function in the order of computation is to

- 1) Initialize (zero all storage areas),
- 2) Sum weighted and unweighted measurements for each type of statistic,
- 3) Compute statistics for each station by measurement type, and
- 4) Compute statistics for all weighted measurements either in the arc or in the entire run.

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

NAME STAINF
 ENTRY POINT PURPOSE
 STAIF1 INITIALIZATION
 STAIF2 TO CORRECT STATISTICAL INFORMATION FOR ELECTRONIC
 BIAS COMPENSATION
 STAINF TO COMPUTE STATISTICAL INFORMATION
 CALLING SEQUENCE CALL STAIF1(ASUM, NSUM, NSTA, NSAT, CSUM, MSUM, NBASE,
 LSUM)

SYMBOL	TYPE	DESCRIPTION
ASUM	I	INPUT - SUMMING ARRAYS FOR PARTICULAR STATIONS, (8.4, NSTA, NSAT) SATELLITES, AND MEASUREMENT TYPES
NSUM	I	INPUT - NUMBER OF RESIDUALS AND WEIGHTED RESIDUAL (3.4, NSTA, NSAT) RATIOS FOR PARTICULAR STATIONS, - SATELLITES, AND MEASUREMENT TYPES
NSTA	I	INPUT - NUMBER OF TRACKING STATIONS
NSAT	I	INPUT - NUMBER OF SATELLITES
CSUM	I	INPUT - SCRATCH SPACE FOR SUMMING STATISTICAL (8.2, NBASE, 1) INFORMATION
MSUM	I	INPUT - SCRATCH SPACE FOR SUMMING STATISTICAL (3.2, NBASE, 1) INFORMATION
NBASE	I	INPUT - NUMBER OF BASE LINES ASSOCIATED WITH MEASUREMENTS INVOLVING TWO STATIONS
LSUM	I	INPUT - SCRATCH SPACE FOR SUMMING STATISTICAL (NBASE, 1) INFORMATION

CALLING SEQUENCE CALL STAIF2(ISTAE, MTYPE, SIGE, EBIAS, NEPASS, NOSAT)

SYMBOL	TYPE	DESCRIPTION
ISTAE	I	INPUT - STATION NUMBER USED IN COMPENSATING FOR ELECTRONIC BIAS
MTPE	I	INPUT - MEASUREMENT TYPE FOR ELECTRONIC BIAS COMPENSATION
SIGE	I	INPUT - SIGMA FOR ELECTRONIC BIAS COMPENSATION
EBIAS	I	INPUT - BIAS WHICH IS COMPENSATED FOR
NEPASS	I	INPUT - NUMBER OF MEASUREMENTS IN THE PASS FOR ELECTRONIC BIAS COMPENSATION

NOSAT I INPUT - SATELLITE NUMBER FOR ELECTRONIC BIAS
 COMPENSATION

CALLING SEQUENCE CALL STAINF(I,TYPE,ISTA,M,TYPE,SIG,RESID,LINNER,
 ISAT,ISTA2)

SYMBOL	TYPE	DESCRIPTION
I,TYPE	I	INPUT - DETERMINES IF INITIALIZATION, ADDITION, STATION SUMMARY, OR TOTAL SUMMARY, IS REQUESTED
ISTA	I	INPUT - INTERNAL STATION NUMBER
M,TYPE	J	INPUT - MEASUREMENT TYPE
SIG	R	INPUT - SIGMA
RESID	R	INPUT - RESIDUAL
LINNER	L	INPUT - .TRUE. - LAST INNER ITERATION
ISAT	I	INPUT - SATELLITE NUMBER
ISTA2	I	INPUT - SECOND STATION WHEN MEASUREMENT INVOLVES TWO STATIONS

SUBROUTINES USED SMSTAT CLEAR RMSAMP

COMMON BLOCKS CSTINF CSTAT

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES 'GEO-DYN SYSTEMS DESCRIPTION'
 VOLUME 1 - GEO-DYN DOCUMENTATION

SUBROUTINE STAI1(ASUM,NSUM,NSTA,NSAT,CRUM,MSUM,NBASE,LSUM)	STAI 08
INTEGER ISAT	STAI 09
DIMENSION ASUM(8,4),NSTA,NSAT),NSUM(8,4),NSTA,NSAT),	STAI 100
CSUM(8,2,NBASE,1),MSUM(8,2,NBASE,1),SUM(NBASE,1)	STAI 101
COMMON/CSTINF/VEASND(4),NOIS(4),DDMEAN(-),RMSD(4),RND(4),	STAI 102
MEASRT(4),WTRMEAN(4),RMSHTC(4),WTRNO(4),TYPRMS(2),NOTYPE(2,30),	STAI 103
NSUM(8,13),RMSALL(3),NOALL(3),NOWTR,NBASE	STAI 104
COMMON/CSTAT/RESID1,SIG1,NMTOT,ITSUMT	STAI 105
NONSTA=NSAT	STAI 106
LAST=NBASE	STAI 107
IST=ISTA	STAI 108
RETURN	STAI 109
COMPENSATE FOR ELECTRONIC BIAS REMOVAL	STAI 110
ENTRY STAI2(ISTA2,M,TYPE,SIG,RESID,NSAT)	STAI 111

ISAT=0	STAI 112
500 ISAT=ISAT+1	STAI 113
DO 600 I=1,4	STAI 114
MNO=NSUM(1,1,ISTA,ISAT)	STAI 115
IF(MNO-MTYPE) 600,625,600	STAI 116
525 ASUM(1,1,ISTA,ISAT)=ASUM(1,1,ISTA,ISAT)	STAI 117
-ERIAS*FLOAT(NEPASS)	STAI 118
ASUM(3,1,ISTA,ISAT)=ASUM(3,1,ISTA,ISAT)	STAI 119
-FLOAT(NEPASS)*ERIAS*2	STAI 120
ASUM(7,1,ISTA,ISAT)=0.	STAI 121
RATIO=ERIAS/SIG	STAI 122
R2A=FLOAT(NEPASS)*RATIO	STAI 123
ASUM(2,1,ISTA,ISAT)=ASUM(2,1,ISTA,ISAT)-R2N	STAI 124
R2N=R2N+RATIO	STAI 125
ASUM(4,1,ISTA,ISAT)=ASUM(4,1,ISTA,ISAT)-R2N	STAI 126
WTSUMT=WTSUMT-R2N	STAI 127
ASUM(3,1,ISTA,ISAT)=0.	STAI 128
600 CONTINUE	STAI 129
IF(ISAT.LT.NCSAT) GO TO 500	STAI 130
RETURN	STAI 131
ENTRY STAINF(ITYPE,ISTA,MTYPE,SIG,RESID,LINNER,ISAT,ISTA)	STAI 132
LOGICAL LINNER	STAI 133
SIG=SIG	STAI 134
RESID=RESID	STAI 135
C DETERMINE IF INITIALIZATION, ADDITION, STATION SUMMARY, OR TOTAL	STAI 136
C SUMMARY	STAI 137
GO TO (1,101,301,401),ITYPE	STAI 138
C ZERO STORAGE AREAS	STAI 139
1 CALL CLEAR(ASUM,32,N)	STAI 140
CALL CLEAR(NSUM,12,N)	STAI 141
CALL CLEAR(TYPRMS,146,1)	STAI 142
IF(LBASE.LE.0) GO TO 10	STAI 143
CALL CLEAR(CSUM,15,LBASE*N)	STAI 144
CALL CLEAR(MSUM,6,LBASE*N)	STAI 145
CALL CLEAR(LSUM,LBASE,N)	STAI 146
10 WTSUMT=0	STAI 147
WTSUMT=0.	STAI 148
IF(ISTA.NE.0) RETURN	STAI 149
NMALL=0	STAI 150
WTSUMA=0.	STAI 151
NODEGF=1	STAI 152
CALL CLEAR(RMSALL,30,2)	STAI 153
RETURN	STAI 154
C SUM STATISTICS	STAI 155
101 IF(MTYPE.GT.14) GO TO 201	STAI 156
C TYPES 1 - 14	STAI 157
DO 200 I=1,4	STAI 158
MNO=NSUM(1,1,ISTA,ISAT)	STAI 159
IF(MNO.GT.0) IF(MNO-MTYPE) 200,125,200	STAI 160
NSUM(3,1,ISTA,ISAT)=MTYPE	STAI 161
125 CALL SRTSTAT(ASUM(1,1,ISTA,ISAT),NSUM(1,1,ISTA,ISAT))	STAI 162
CONTINUE	STAI 163
200 CONTINUE	STAI 164
RETURN	STAI 165
201 IF(MTYPE.GT.26) GO TO 210	STAI 166
C TYPES 15 - 26	STAI 167

```

NTYPE=NTYPE-14
CALL SMSTAT(RSUM(1,NTYPE),NOTYPE(1,NTYPE))
RETURN
C TYPES 27 - 30
210 IND=(ISAT-1)*LSTA+ISTA
DO 275 J=1,LPASE
KSTA=LSUM(J,IND)
IF(KSTA.GT.0) IF(KSTA-ISTA?) 275,225,275
LSUM(J,IND)=ISTA2
225 DO 275 I=1,2
MNI=MSUM(3,I,J,IND)
IF(MNI.GT.0) IF(MNI-MTYPE) 270,250,270
MSUM(3,I,J,IND)=NTYPE
250 CALL SVSTAT(CSUM(1,I,J,IND),MSUM(1,I,J,IND))
RETURN
270 CONTINUE
275 CONTINUE
RETURN
C STATION SUMMARY
301 IF(ISTA.EQ.0) GO TO 351
IF(ISTA2.GT.0) GO TO 325
C SINGLE STATION MEASUREMENTS
CALL PRSUMP(NSUM(1,1,ISTA,ISAT),ASUM(1,1,ISTA,ISAT),4)
RETURN
C TWO STATION MEASUREMENTS
325 IND=(ISAT-1)*LSTA+ISTA
DO 350 J=1,LPASE
KSTA=LSUM(J,IND)
IF(KSTA.NE.ISTA2) GO TO 350
CALL PRSUMP(NSUM(1,1,J,IND),CSUM(1,1,J,IND),2)
NORS(3)=0
RETURN
350 CONTINUE
NORS(1)=0
RETURN
C MEASUREMENTS NOT INVOLVING STATIONS
351 DO 400 I=1,12
J=I+74
K=NOTYPE(1,J)
IF(K.EQ.0) GO TO 400
XN=K
IF(K.LT.10) GO TO 395
RSUM(5,1)=(RSUM(3,1)-RSUM(1,1)**2/XN)/RSUM(5,1)
RSUM(5,1)=(2.*RSUM(5,1)-1.)/SQRT((XN-2.)/(XN**2-1.))
RSUM(3,1)=SQRT(RSUM(3,1)/(XN-1.))
395 RSUM(2,1)=RSUM(1,1)/XN
K=NOTYPE(2,J)
IF(K.EQ.0) GO TO 400
XN=K
TYCDMS(1)=RSUM(4,1)
IF(K.LT.10) GO TO 395
RSUM(7,1)=(RSUM(5,1)-RSUM(3,1)**2/XN)/RSUM(7,1)
RSUM(7,1)=(2.*RSUM(7,1)-1.)/SQRT((XN-2.)/(XN**2-1.))
RSUM(5,1)=SQRT(RSUM(5,1)/(XN-1.))
395 RSUM(2,1)=RSUM(2,1)/XN
400 CONTINUE

```

STAI 169
STAI 160
STAI 170
STAI 171
STAI 172
STAI 173
STAI 174
STAI 175
STAI 176
STAI 177
STAI 178
STAI 179
STAI 180
STAI 181
STAI 182
STAI 183
STAI 184
STAI 185
STAI 186
STAI 187
STAI 188
STAI 189
STAI 190
STAI 191
STAI 192
STAI 193
STAI 194
STAI 195
STAI 196
STAI 197
STAI 198
STAI 199
STAI 200
STAI 201
STAI 202
STAI 203
STAI 204
STAI 205
STAI 206
STAI 207
STAI 208
STAI 209
STAI 210
STAI 211
STAI 212
STAI 213
STAI 214
STAI 215
STAI 216
STAI 217
STAI 218
STAI 219
STAI 220
STAI 221
STAI 222
STAI 223

RETURN	STAI 224
C TOTAL SUMMARY	STAI 225
401 IF (ISTA.EQ.0) GO TO 467	STAI 226
NTYPE=NMTOT	STAI 227
IF (.NOT.LINNER) GO TO 410	STAI 228
NMALL=NMALL+NMTOT	STAI 229
NODEGF=NODEGF+ISTA	STAI 230
WTSUMA=WTSUMA+WTSUMT	STAI 231
410 SIG=SQRT (WTSUMT/FLDAT (C.MTOT-ISTA-1))	STAI 232
DO 450 I=1,30	STAI 233
IF (LINNER) NCALL (I)=NCALL (I)+NCTYPE (2,I)	STAI 234
IF (LINNER) RMSALL (I)=RMSALL (I)+TYRMS (I)	STAI 235
450 IF (NCTYPE (2,I).NE.0) TYRMS (I)=SQRT (TYRMS (I)/FLOAT (NCTYPE (2,I)))	STAI 236
RETURN	STAI 237
460 NODEGF=NODEGF+NTYPE	STAI 238
NTYPE=NMALL	STAI 239
SIG=SQRT (WTSUMA/FLOAT (NMALL-NODEGF))	STAI 240
DO 470 I=1,30	STAI 241
470 IF (NCALL (I).NE.0) TYRMS (I)=SQRT (RMSALL (I)/FLOAT (NCALL (I)))	STAI 242
RETURN	STAI 243
END	STAI 244
DIMENSION NCTYPE (2), SUM (8)	STAI 245
COMMON/STAT /RESID, SIG, NMTOT, WTSUMT	STAI 246
IF (NCTYPE (1).EQ.0) SUM (7)=RESID	STAI 247
NCTYPE (1)=NCTYPE (1)+1	STAI 248
SUM (1)=SUM (1)+RESID	STAI 249
SUM (3)=SUM (3)+RESID**2	STAI 250
SUM (5)=SUM (5)+(RESID-SUM (7))**2	STAI 251
SUM (7)=RESID	STAI 252
IF (SIG.EQ.0.) RETURN	STAI 253
R=RESID/SIG	STAI 254
R2=R**2	STAI 255
NMTOT=NMTOT+1	STAI 256
WTSUMT=WTSUMT+R2	STAI 257
IF (NCTYPE (2).EQ.0) SUM (8)=R	STAI 258
NCTYPE (2)=NCTYPE (2)+1	STAI 259
SUM (2)=SUM (2)+R	STAI 260
SUM (4)=SUM (4)+R2	STAI 261
SUM (6)=SUM (6)+(R-SUM (8))**2	STAI 262
SUM (8)=R	STAI 263
RETURN	STAI 264
END	STAI 265

STAINP

DESCRIPTION

STAINP reads tracking station positions from the GEODYN Input Cards. STAINP can read tracking stations in geodetic (latitude, longitude, height) or Cartesian coordinates.

STAINP assumes the station position to be in Cartesian coordinates. If, however, the station vector is considerably smaller in magnitude than one Earth radius STAINP considers the station position to be given in geodetic coordinates.

STAINP will not accept duplicate stations and prints an error message upon any such encounters.

This subroutine also loads the variable storage station arrays selecting station coordinates from the input values supplemented by station coordinate values stored in the block data subroutine STAPOS.

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

NAME STAINP
 PURPOSE TO READ GEDDYN TRACKING STATIONS AND CONVERT FROM
 CARTESIAN COORDINATES WHENEVER NECESSARY
 CALLING SEQUENCE CALL STAINP(NSTA,KSTAND,STANAM,STALAT,STALON,
 HEIGHT,NSTARD,NSTEST,ESTAND)

SYMBOL	TYPE	DESCRIPTION
NSTA	I	INPUT - NUMBER OF STATIONS USED
KSTAND (1)	I*2	OUTPUT - STATION NUMBERS
STANAM (1)	DP	OUTPUT - STATION NAMES
STALAT (1)	DP	OUTPUT - STATION LATITUDES
STALON (1)	DP	OUTPUT - STATION LONGITUDES
HEIGHT (1)	DP	OUTPUT - STATION HEIGHTS
NSTARD	I	INPUT - NUMBER OF STATIONS THAT WERE READ FROM CARDS
NSTEST	I	INPUT - NUMBER OF STATIONS REQUESTING ADJUSTMENT
ESTAND	I*2	OUTPUT - MASTER STATION NUMBERS FOR ESTIMATED STATIONS

SUBROUTINES USED	NUMBR2	ERROR	DARCTN		
COMMON BLOCKS	CEPHEM	STANUM	STAPOS	CONSTS	INTRLK
	TPEBLK				

INPUT FILES INTP - GEDDYN INPUT CARDS

OUTPUT FILES NONE



REFERENCES 'GEDDYN SYSTEMS DESCRIPTION'
 VOLUME 1 - GEDDYN DOCUMENTATION



SUBROUTINE STAINP(NSTA,KSTAND,STANAM,STALAT,STALON,HEIGHT,
 NSTARD,NSTEST,ESTAND)
 IMPLICIT REAL*8 (A-H,C-Z)
 DIMENSION KSTAND(1),STANAM(1),STALAT(1),STALON(1),HEIGHT(1),
 ESTAND(1),NAMEB(9)

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

LOGICAL*1 NAME8,NAME7	STAI 56
INTEGER*2 KSTAND,ISTAND,STANDS,ESTAND,ISTARD,ESTAND	STAI 57
INTEGER OUTP	STAI 58
DOUBLE PRECISION NAME,L	STAI 59
COMMON/CEPHEM/DNAME(3P1),ISTARD(381),ESTAND(381),ISTAND(356)	STAI 60
COMMON/STANUM/SNAME(280),STANDS(280),NOSTOR	STAI 61
COMMON/STAPDS/RLAT(280),RLON(280),H(280)	STAI 62
COMMON/CONSTS/PI,TWOP1,DRAD,RSFC	STAI 63
COMMON/INTBLK/THCOT(4),AE,AE50,FLAT,FS032(59)	STAI 64
COMMON/TPEBLK/INTP,OUTP,IDUM(10)	STAI 65
DATA DATA/6HDATA /,STAPDS/6HSTAPDS/	STAI 66
DATA ALTMTR/6HALTMTR/	STAI 67
DATA DELTA/1.00-3/	STAI 68
EQUIVALENCE (JSTAND,DSTAND),(NAME,NAME8(1)),(NAME7,NAME8(7))	STAI 69
IF(NSTA.LE.0) RETURN	STAI 70
DO 10 I=1,NSTA	STAI 71
10 KSTAND(I)=0	STAI 72
IF(NSTARD.LE.0) GO TO 1000	STAI 73
REWIND INTP	STAI 74
READ(INTP,5000)	STAI 75
ESQ=(1.000-FLAT)**2	STAI 76
ESQ=1.000-ESQ	STAI 77
20 READ(INTP,10005) NAME	STAI 78
IF(NAME.NE.STAPDS) GO TO 20	STAI 79
C READ STATION POSITION CARDS.	STAI 80
100 READ(INTP,10005,FND=900) NAME,JSTAND,P,L,HT,NAME7	STAI 81
IF(NAME.FQ.DATA) GO TO 1000	STAI 82
ISN=NUMB02(JSTAND,ISTAND,NSTA)	STAI 83
C SKIP CARD IF STATION NOT USED	STAI 84
IF(ISN.LE.0) GO TO 100	STAI 85
IF(KSTAND(ISN).LE.0) GO TO 150	STAI 86
CALL ERROR(3,DSTAND)	STAI 87
GO TO 100	STAI 88
C CHECK COORDINATE SYSTEMS	STAI 89
150 XYSQ=P*P+L*L	STAI 90
RSQ=XYSQ+HT*HT	STAI 91
IF(RSQ.LT.3.5D13) GO TO 400	STAI 92
C CONVERT CARTESIAN TO SPHERICAL COORDINATES	STAI 93
T=ESQ*HT	STAI 94
C ...HEIGHT	STAI 95
DO 200 J=1,25	STAI 96
ZT=HT+T	STAI 97
H1=DSQRT(XYSQ+ZT**2)	STAI 98
SINPHI=ZT/H1	STAI 99
ESQSP=ESQ*SINPHI	STAI 100
H2=AE/DSQRT(1.000-ESQSP*SINPHI)	STAI 101
T1=H2*ESQSP	STAI 102
IF(DABS(T1-T).LT.DELTA) GO TO 300	STAI 103
200 T=T1	STAI 104
300 HT=H1-H2	STAI 105
RTXYSQ=DSQRT(XYSQ)	STAI 106
RSQ=DATAN2(ZT,RTXYSQ)	STAI 107
C ...LONGITUDE	STAI 108
L=CARCTN(L,P)	STAI 109
C ...LATITUDE	STAI 110
P=RSQ	STAI 111

GO TO 500	STAI 112
400 JLATD=P*1.0D-4	STAI 113
P=P-JLATD*1.0D+4	STAI 114
JLATM=P*1.0D-2	STAI 115
TSLAT=P-JLATM*1.0D+2	STAI 116
JLCND=L*1.0D-4	STAI 117
L=L-JLOND*1.0D+4	STAI 118
JLONM=L*1.0D-2	STAI 119
TSLCN=L-JLCNM*1.0D+2	STAI 120
P=(DFLOAT(JLATD)+DFLOAT(JLATM)/6.0D1+TSLAT/3.6D3)*ORAD	STAI 121
L=(DFLOAT(JLCND)+DFLOAT(JLONM)/6.0D1+TSLCN/3.6D3)*ORAD	STAI 122
C STORE STATION POSITION IN VARIABLE STORAGE ARRAYS	STAI 123
500 KSTANO(ISN)=JSTANO	STAI 124
STANAM(ISN)=NAME	STAI 125
STALAT(ISN)=P	STAI 126
STALON(ISN)=L	STAI 127
HEIGHT(ISN)=HT	STAI 128
IF(ISN.LE.NSTEST) ESTANO(ISN)=ESTANO(ISN)	STAI 129
GO TO 100	STAI 130
900 CALL ERROR(6,DSTANO)	STAI 131
RETURN	STAI 132
1000 DO 2000 I=1,NSTA	STAI 133
IF(KSTANO(I).NE.0) GO TO 2000	STAI 134
JSTANO=ISTANC(I)	STAI 135
ISN=NUMB2(JSTANO,STANOS,NOSTOR)	STAI 136
IF(ISN.GT.0) GO TO 1500	STAI 137
IF(JSTANO.NE.-4) GO TO 1200	STAI 138
KSTANO(I)=-4	STAI 139
STANAM(I)=ALTMTR	STAI 140
STALAT(I)=0.0D0	STAI 141
STALON(I)=0.0D0	STAI 142
HEIGHT(I)=0.0D0	STAI 143
GO TO 2000	STAI 144
1200 CALL ERROR(9,DSTANO)	STAI 145
RETURN	STAI 146
C IF STATION FROM ELOCK DATA THEN TRANSFER TO VARIABLE STORAGE ARRAYS	STAI 147
1500 KSTANO(I)=JSTANO	STAI 148
STANAM(I)=SNAME(ISN)	STAI 149
STALAT(I)=RLAT(ISN)	STAI 150
STALON(I)=RLCN(ISN)	STAI 151
HEIGHT(I)=H(ISN)	STAI 152
IF(I.LE.NSTEST) ESTANO(I)=ESTANO(I)	STAI 153
2000 CONTINUE	STAI 154
C ZERO COORDINATES FOR ALTIMETER	STAI 155
RETURN	STAI 156
5000 FORMAT(1X/1X/1X)	STAI 157
10005 FORMAT(A5,I4,3D15.8,A1)	STAI 158
END	STAI 159





REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

NAME SMSTAT
 PURPOSE TO SUM STATISTICS
 CALLING SEQUENCE CALL SMSTAT(SUM,NOTYPE)

SYMBOL	TYPE	DESCRIPTION
SUM (8)	R	INPUT & OUTPUT - ARRAY THAT INFORMATION GETS SUMMED INTO
NOTYPE (2)	I	INPUT & OUTPUT - ARRAY CONTAINING NUMBER OF MEASUREMENTS

SUBROUTINES USED NONE

COMMON BLOCKS CSTAT

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES 'GEODYN SYSTEMS DESCRIPTION'
 VOLUME 1 - GEODYN DOCUMENTATION

```

SUBROUTINE SMSTAT(SUM,NOTYPE)
DIMENSION NOTYPE(2),SUM(8)
COMMON/CSTAT /RESID,SIG,NMTOT,WTSUMT
IF(NOTYPE(1).EQ.0) SUM(7)=RESID
NOTYPE(1)=NOTYPE(1)+1
SUM(1)=SUM(1)+RESID
SUM(3)=SUM(3)+RESID**2
SUM(5)=SUM(5)+(RESID-SUM(7))**2
SUM(7)=RESID
IF(SIG.LE.0.) RETURN
R=RESID/SIG
R2=R**2
NMTOT=NMTOT+1
WTSUMT=WTSUMT+R2
IF(NOTYPE(2).EQ.0) SUM(8)=R
NOTYPE(2)=NOTYPE(2)+1
SUM(2)=SUM(2)+R
SUM(4)=SUM(4)+R2
SUM(6)=SUM(6)+(R-SUM(8))**2
SUM(8)=R
RETURN
END
  
```

STAI 273
 STAI 274
 STAI 275
 STAI 276
 STAI 277
 STAI 278
 STAI 279
 STAI 280
 STAI 281
 STAI 282
 STAI 283
 STAI 284
 STAI 285
 STAI 286
 STAI 288
 STAI 289
 STAI 290
 STAI 291
 STAI 292
 STAI 293
 STAI 294

395	B SUM(2,I)=B SUM(2,I)/XN	STAI	232
400	CONTINUE	STAI	233
	RETURN	STAI	234
C	TOTAL SUMMARY	STAI	235
401	IF(ISTA.EQ.0) GO TO 460	STAI	236
	MTYPE=NMTOT	STAI	237
	IF(.NOT.LINNER) GO TO 410	STAI	238
	NMALL=NMALL+NMTOT	STAI	239
	NODEGF=NODEGF+I STA	STAI	240
	WTSUMA=WTSUMA+WTSUMT	STAI	241
410	SIG=SQRT(WTSUMT/FLOAT(NMTOT-ISTA-1))	STAI	242
	DO 450 I=1,30	STAI	243
	IF(LINNER) NOALL(I)=NCALL(I)+NOTYPE(2,I)	STAI	244
	IF(LINNER) RMSALL(I)=RMSALL(I)+TYPRMS(I)	STAI	245
450	IF(NOTYPE(2,I).NE.0) TYPRMS(I)=SQRT(TYPRMS(I)/FLOAT(NOTYPE(2,I)))	STAI	246
	RETURN	STAI	247
460	NODEGF=NODEGF+MTYPE	STAI	248
	MTYPE=NMALL	STAI	249
	SIG=SQRT(WTSUMA/FLOAT(NMALL-NODEGF))	STAI	250
	DO 470 I=1,30	STAI	251
470	IF(NOALL(I).NE.0) TYPRMS(I)=SQRT(RMSALL(I)/FLOAT(NOALL(I)))	STAI	252
	RETURN	STAI	253
	END	STAI	254

NAME STAPOS
PURPOSE BLOCK DATA STORAGE OF STATION POSITIONS
UNITS FOR COORDINATES
LAT - RADIANS
LON - RADIANS
H - METERS

COMMON BLOCKS STAPOS STANUM

BLOCK DATA										STAP	14
DOUBLE PRECISION LAT, LON, HT, NAME										STAP	15
INTEGER*2 NUMBER										STAP	16
COMMON/ST APOS/LAT(280), LON(280), HT(280)										STAP	17
COMMON/ST ANUM/NAME(280), NUMBER(280), NSTOR										STAP	18
DATA NSTOR/254/										STAP	19
DOUBLE PRECISION NAME 1										STAP	20
DIMENSION NAME 1(114)										STAP	21
EQUIVALENCE (NAME (1), NAME 1(1))										STAP	22
DATA NAME 1/										STAP	23
• 7H4IL3	• 7HGBM3	• 7HADA3	• 7HANG3	• 7HCY13	• 7HACN3					STAP	24
• 7H4AD3	• 7HCRD3	• 7HGWM3	• 7HHSK3	• 7HHAW3	• 7HGRS3					STAP	25
• 7HGYM3	• 7HTEX3	• 7HGTG3	• 7HPIOD	• 7HECHO	• 7HVENO					STAP	26
• 7H4AP3	• 7H4QMD	• 7HTID3	• 7HJOB3	• 7HR3BD	• 7HIC3D					STAP	27
• 7H4RTD	• 7HKOURDU	• 7HRPOINT	• 7HFTMYR6	• 7HQUIT06	• 7HLIMAP6					STAP	28
• 7HSNTAG6	• 7HNEWFL6	• 7HCOLLEGE	• 7HGFRPK5	• 7HWNKFL6	• 7HJORUR6					STAP	29
• 7HMOJAV6	• 7HDOOMERA	• 7HULASK5	• 7HHPIN	• 7HFTMYR	• 7HMAOGA6					STAP	30
• 7H1OOMER	• 7H1OUI10	• 7H1LIMAP	• 7H1SATAG	• 7H1MOJAV	• 7H1JOBUP					STAP	31
• 7H1NEWFL	• 7H1COLLEG	• 7H1GFORK	• 7H1WNKFL	• 7H1ULASK	• 7H1ROSMN					STAP	32
• 7H1OPORL	• 7H1ROSMA	• 7H1TANAV	• 7HORORA6	• 7HMADGARS	• 7HRDSPANS					STAP	33
• 7HSNTAGR5	• 7HALASKR5	• 7HCARVONS	• 7HMADGARV	• 7HR0SRANV	• 7HSNTAGRV					STAP	34
• 7HALASKRV	• 7HCARVONV	• 7HANTIGA	• 7HGRNVLE	• 7HGRVILL	• 7HUSAFAC					STAP	35
• 7HEDFRD	• 7HSEMME5	• 7H5WANIS	• 7HGRDTRK	• 7HURACO	• 7HTRNOAD					STAP	36
• 7HGRANFK	• 7HTWINDK	• 7HPOTHGR	• 7HATHNGR	• 7HTDRRSP	• 7HCHOFUJ					STAP	37
• 7H1INDLY	• 7HEDWADS	• 7HHUNTER	• 7HJUPRAF	• 7HABERDN	• 7HHOME5T					STAP	38
• 7HCHYWY4	• 7HETRGR6	• 7HETPCAK	• 7HETRPRE	• 7HETRPAT	• 7HETRANT					STAP	39
• 7HETRGR7	• 7HETFMRT	• 7HETRGR8	• 7HWSH122	• 7HWSC113	• 7HWSH123					STAP	40
• 7H4SS127	• 7HWSP124	• 7HWST125	• 7HWTRPP5	• 7HWTRTR2	• 7HWTRTR1					STAP	41
• 7H4TRPP3	• 7HWTRVAN	• 7HEGLINE	• 7HEGLINE	• 7H4MRSN5	• 7H4MRSN6					STAP	42
DOUBLE PRECISION NAME 2										STAP	43
DIMENSION NAME 2(114)										STAP	44
EQUIVALENCE (NAME (115), NAME 2(1))										STAP	45
DATA NAME 2/										STAP	46
• 7HPMRV1	• 7HPMRV2	• 7HPMRV41	• 7HPMRV42	• 7HPMRSN2	• 7HPMRSN3					STAP	47
• 7HPMRSN4	• 7HPMRSN3	• 7HPMRSN4	• 7HPMRK1	• 7HFDAPB3	• 7HNELHAR					STAP	48
• 7HNELYV	• 7HNWI#2A	• 7HNWI#2B	• 7HNWI#3A	• 7HNWI#3B	• 7HNRER34					STAP	49
• 7HNTANAN	• 7HWTEKAU	• 7HNRBEROS	• 7HNCARNV	• 7HNWALI5	• 7HNWALI3					STAP	50
• 7HNDJ3R	• 7HCAFAB1	• 7HCAFAB5	• 7HNDURTN	• 7HSANSAL	• 7HPTARGO					STAP	51
• 7HHERNDN	• 7HCUEICAL	• 7HLAFSON	• 7HWRTON	• 7HGRENV	• 7HVAUIHA					STAP	52
• 7H4ALSEC	• 7HETKART	• 7HPINTAFB	• 7HPMREFL	• 7HRELTVL	• 7HVOLGGD					STAP	53
• 7HEREVAN	• 7HKIEVAA	• 7HKRASND	• 7HNQVQSI	• 7HRIAZAN	• 7HTARTU2					STAP	54
• 7HTASHK1	• 7HASTRMD	• 7HLYNNLK	• 7HTIMENS	• 7H1UNDAK	• 7H1EDINA					STAP	55

• 7HICOLRA	• 7HIBERMD	• 7HIPURID	• 7HIGSFCO	• 7HIGSFCP	• 7HICKVLE	• STAP	56							
• 7HIDENVR	• 7HIJUM24	• 7HIJUM40	• 7HIJUPC1	• 7HIJURC4	• 7HISDRR	• STAP	57							
• 7HIJAMAC	• 7HIGSFCN	• 7HWALMOT	• 7HICARVN	• 7HGDOLAS	• 7HRNSLAS	• STAP	58							
• 7HWALLAS	• 7HYOILAS	• 7HCRMLAS	• 7HHOVLAS	• 7HHOMLA2	• 7HMOBLA2	• STAP	59							
• 7HSENLAS	• 7HSFLLAS	• 7HCRGLAS	• 7HOLILAS	• 7HARELAS	• 7HHOPLAS	• STAP	60							
• 7HNATLAS	• 7HGRELAS	• 7HHRNSCH	• 7HDELFTH	• 7HZIMWLD	• 7HMALVRN	• STAP	61							
• 7HROYORS	• 7HATHENS	• 7HHAUTEP	• 7HNICEFR	• 7HVICLAS	• 7HSALLAS	• STAP	62							
• 7HMIDONI	• 7HEDINBH	• 7HMUNICH	• 7HFRANKF	• 7HSANFLR	• 7HHAUTLR	• STAP	63							
• 7HIORGAN	• 7HIOLFAN	• 7HWODMER	• 7HISPAIN	• 7HITOKYO	• 7HINATAL	• STAP	64							
• 7HIQUIPA	• 7HISRAZ	• 7HICURAC	• 7HIJUPTR	• 7HIVILDO	• 7HMAUID	• STAP	65							
DOUBLE PRECISION NAME 3						• STAP	66							
DIMENSION NAME 3(26)						• STAP	67							
EQUIVALENCE (NAME (229),NAME 3(1))						• STAP	68							
DATA NAME 3/						• STAP	69							
• 7HDAKARO	• 7HHOPKIN	• 7HAUSBAK	• 7HODDAIR	• 7HOFZEIT	• 7HNATALR	• STAP	70							
• 7HCOMRIV	• 7HBRAZIL	• 7HJUPGED	• 7HAGASSI	• 7HZIMMER	• 7HRIGALA	• STAP	71							
• 7HUZHGR	• 7HGRFECE	• 7HCALBAK	• 7HCOLAKE	• 7HOSLONR	• 7HJOHNST	• STAP	72							
• 7HMTJOHN	• 7HSANVTD	• 7HEDWAFB	• 7HPOTDAM	• 7HZVENIG	• 7HHELSIK	• STAP	73							
• 7HDAKAR	• 7HSANVIT					• STAP	74							
INTEGER*2 NUMB 1						• STAP	75							
DIMENSION NUMB 1(247)						• STAP	76							
EQUIVALENCE (NUMBER(1),NUMB 1(1))						• STAP	77							
DATA NUMB 1/						• STAP	78							
• 1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	12.	13.	STAP	79
• 14.	15.	511.	512.	513.	514.	541.	542.	551.	561.	562.	571.	1000.	STAP	80
• 1001.	1003.	1005.	1006.	1008.	1012.	1013.	1014.	1015.	1016.	1017.	1018.	1019.	STAP	81
• 1021.	1022.	1023.	1024.	1025.	1026.	1028.	1029.	1031.	1032.	1033.	1034.	1035.	STAP	82
• 1036.	1037.	1039.	1040.	1043.	1121.	1123.	1125.	1127.	1128.	1132.	1022.	1026.	STAP	83
• 1627.	1629.	1652.	3106.	3333.	3334.	3400.	3401.	3402.	3404.	3405.	3406.	3407.	STAP	84
• 3451.	3452.	3453.	3463.	3464.	3465.	3471.	3472.	3448.	3449.	3457.	3461.	3402.	STAP	85
• 4040.	4041.	4050.	4060.	4061.	4081.	4082.	4083.	4142.	4143.	4144.	4145.	4146.	STAP	86
• 4151.	4240.	4241.	4242.	4260.	4280.	4340.	4341.	4400.	4401.	4402.	4403.	4440.	STAP	87
• 4441.	4442.	4443.	4444.	4445.	4446.	4450.	4540.	4610.	4690.	4732.	4733.	4734.	STAP	88
• 4735.	4740.	4741.	4742.	4760.	4761.	4840.	4860.	4946.	4948.	4949.	4954.	4006.	STAP	89
• 4003.	5001.	5200.	5201.	5202.	5333.	5411.	5508.	5648.	5649.	5861.	6002.	6014.	STAP	90
• 6018.	6023.	6027.	6035.	6042.	6051.	6052.	6100.	6107.	6113.	7034.	7036.	7037.	STAP	91
• 7039.	7040.	7042.	7043.	7044.	7045.	7071.	7072.	7073.	7074.	7075.	7076.	7077.	STAP	92
• 7078.	7079.	7050.	7051.	7052.	7053.	7054.	7055.	7056.	7057.	7058.	7804.	7901.	STAP	93
• 7902.	7907.	7921.	7929.	7930.	8004.	8009.	8010.	8011.	8013.	8014.	8015.	8019.	STAP	94
• 8021.	8022.	8030.	8031.	8032.	8033.	7804.	7815.	9001.	9002.	9003.	9004.	9005.	STAP	95
• 9006.	9007.	9008.	9009.	9010.	9011.	9012.	9020.	9021.	9023.	9025.	9028.	9029.	STAP	96
• 9031.	9039.	9049.	9050.	9056.	9074.	9077.	9091.	9113.	9114.	9115.	9117.	9119/	STAP	97
INTEGER*2 NUMB 2													• STAP	98
DIMENSION NUMB 2(7)													• STAP	99
EQUIVALENCE (NUMBER(248),NUMB 2(1))													• STAP	100
DATA NUMB 2/													• STAP	101
• 9120.	9423.	9429.	9430.	9435.	9020.	9120/							• STAP	102
DOUBLE PRECISION LAT 1													• STAP	103
DIMENSION LAT 1(76)													• STAP	104
EQUIVALENCE (LAT (1),LAT 1(1))													• STAP	105
DATA LAT 1/													• STAP	106
• 0.4775001100.	0.4648276700.	0.5646344700.	0.2959950200.										• STAP	107
• 0.4465773600.	-0.1334472900.	0.7160312400.	-0.4347009500.										• STAP	108
• 0.2123114700.	-0.6210474400.	0.3261777700.	0.6168287300.										• STAP	109
• 0.4980150300.	0.4826454000.	0.6806514000.	0.6176670500.										• STAP	110
• 0.6160726300.	0.6151878900.	0.6182781100.	-0.5477174200.										• STAP	111

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

•-0.6178622100, -0.4518644800, 0.7056151400, 0.7060395900,	STAP 112
• 0.4970598800, 0.0916487200, 0.6707397000, 0.4633516600,	STAP 113
•-0.0108712000, -0.2055467400, -0.5785708000, 0.8332457200,	STAP 114
• 1.1322234700, 0.3281533100, 0.8979059500, -0.4517563700,	STAP 115
• 0.6166243100, -0.5478679600, 1.1340604100, 0.6707397000,	STAP 116
• 0.4633516600, -0.3317705300, -0.5478679600, -0.0108712000,	STAP 117
•-0.2055467400, -0.5785708000, 0.6166243100, -0.4517563700,	STAP 118
• 0.8332457200, 1.1322234700, 0.3281533100, 0.8979059500,	STAP 119
• 1.1340604100, 0.6143911500, -0.6217865500, 0.6143912600,	STAP 120
•-0.3317705300, -0.6217865500, -0.3219736000, 0.6142862800,	STAP 121
•-0.5785708000, 1.1339744700, -0.4346424900, -0.3319372200,	STAP 122
• 0.6142862800, -0.5786094200, 1.1339663000, -0.4346616900,	STAP 123
• 0.2993004800, 0.5843433000, 0.5833881400, 0.6907874800,	STAP 124
• 0.7409771300, 0.5372224600, 0.3037798800, 0.3740273200/	STAP 125
DOUBLE PRECISION LAT 2	STAP 126
DIMENSION LAT 2(76)	STAP 127
EQUIVALENCE (LAT (77),LAT 2(1))	STAP 128
DATA LAT 2/	STAP 129
• 0.2110192200, 0.1875053000, 0.8367821000, 0.6304874800,	STAP 130
• 0.8973779700, 0.6613157600, 0.7066373700, 0.6225417700,	STAP 131
• 0.5651906200, 0.6102373900, 0.5585368700, 0.4716763400,	STAP 132
• 0.6899158000, 0.4451848900, 0.7179078300, 0.4645295500,	STAP 133
• 0.4970375000, -0.4528068400, 0.4926431000, 0.2992148800,	STAP 134
• 0.3745930300, 0.4961043100, 0.4649886400, 0.5742537300,	STAP 135
• 0.5547535900, 0.5742393000, 0.5901640300, 0.5837297300,	STAP 136
• 0.5776768400, 0.6644563900, 0.6035870600, 0.5035820000,	STAP 137
• 0.6544568600, 0.6050024300, 0.5509574600, 0.5309574600,	STAP 138
• 0.5502530900, 0.5602502200, 0.3863106000, 0.3863176400,	STAP 139
• 0.5955307800, 0.5955426600, 0.5802685800, 0.5902909300,	STAP 140
• 0.5802932900, 0.5955493700, 0.5955560800, 0.3843570100,	STAP 141
• 0.6103339100, 0.6860674500, 0.6860646500, 0.6609076500,	STAP 142
• 0.6609076800, 0.5518311900, 0.6518311900, 0.5645785000,	STAP 143
•-0.3316439300, 0.3861247800, 0.5645760600, -0.4345251300,	STAP 144
• 0.6504548000, 0.6607850300, -0.5378853000, 0.9098049200,	STAP 145
• 0.9093035000, 0.7877440400, 0.4209402400, 0.6035820000,	STAP 146
• 0.6305709500, 0.5728229200, 0.8235294400, 0.7618259900,	STAP 147
• 0.5333900500, 0.3634413500, 0.6607702600, 0.5571432000/	STAP 148
DOUBLE PRECISION LAT 3	STAP 149
DIMENSION LAT 3(76)	STAP 150
EQUIVALENCE (LAT (153),LAT 3(1))	STAP 151
DATA LAT 3/	STAP 152
• 0.5595279600, 0.4448769700, 0.6811590400, 1.0336237700,	STAP 153
• 0.7022041400, 0.8905740200, 0.7858926700, 0.9596402000,	STAP 154
• 0.9535557900, 1.0181087300, 0.7216936500, 0.6811627700,	STAP 155
• 0.9224093400, 0.8476259200, 0.8381533100, 0.4604106900,	STAP 156
• 0.6798179000, 0.5648560900, 0.3186611400, 0.6810291800,	STAP 157
• 0.6310427400, 0.5696861900, 0.6919655700, 0.4715965000,	STAP 158
• 0.4715964300, 0.4715921300, 0.4715992300, 0.6108078400,	STAP 159
• 0.3158378700, 0.6806632100, 0.6509348200, -0.4347004300,	STAP 160
• 0.6810375600, 0.6142927000, 0.6607818000, 0.6810442400,	STAP 161
•-0.4346664600, 0.5530146400, 0.5530141900, 0.6810442400,	STAP 162
• 0.7452701400, 0.6363926200, 0.5659993900, -0.4530906700,	STAP 163
•-0.237384100, 0.5520937800, -0.1034601100, 0.6645937700,	STAP 164
• 0.9177295700, 0.9076039800, 0.8131578900, 0.9100748600,	STAP 165
• 0.9727393400, 0.5630191500, 0.7667694600, 0.7631600000,	STAP 166
• 0.7667700300, -0.5061475400, 0.8518304400, 0.9727393400,	STAP 167

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

• 0.8342934600, 0.9765023500, 0.6363942200, 0.7667606100,	STAP 168
• 0.5558982400, -0.4530906700, -0.5428122900, 0.6364003300,	STAP 169
• 0.6225104300, 0.5124156300, -0.2873844100, 0.5172661300,	STAP 170
• 0.2110155900, 0.4715972200, -0.5575122000, 0.3614076200/	STAP 171
DOUBLE PRECISION LAT 4	STAP 172
DIMENSION LAT 4(26)	STAP 173
EQUIVALENCE (LAT (229),LAT 4(1))	STAP 174
DATA LAT 4/	STAP 175
• 0.2573255900, 0.5529928100, -0.5478715800, 0.6284151100,	STAP 176
• 0.1526712600, -0.1034691100, -0.8008958600, -0.1034610300,	STAP 177
• 0.4715962900, 0.7418664900, 0.8181579000, 0.9939423200,	STAP 178
• 0.8488183800, 0.6646039000, 0.6102377300, 0.9554430200,	STAP 179
• 1.0508735900, 0.2922282700, -0.7677514300, 0.7092800400,	STAP 180
• 0.6102376900, 0.9142272100, 0.9720339200, 1.0500206700,	STAP 181
• 0.2573023300, 0.7092800200/	STAP 182
DOUBLE PRECISION LON 1	STAP 183
DIMENSION LON 1(76)	STAP 184
EQUIVALENCE (LON (1),LON 1(1))	STAP 185
DATA LON 1/	STAP 186
• 4.8748156400, 4.9176780500, 5.1546866900, 5.2053985900,	STAP 187
• 6.0102959000, 6.0331221900, 6.2104237400, 1.9846313600,	STAP 188
• 2.5261452500, 2.6001518600, 3.4965034700, 4.2433522500,	STAP 189
• 4.3507345100, 4.5836045800, 4.9420226300, 4.2437751500,	STAP 190
• 4.2445419400, 4.2447269400, 4.2430749900, 2.3891359000,	STAP 191
• 2.6002125100, 0.4931583300, 6.2090229700, 6.2069493300,	STAP 192
• 4.8770303600, 5.3615589600, 4.9377721400, 4.8543625900,	STAP 193
• 4.9117216900, 4.9366590700, 5.0497895700, 5.3630387100,	STAP 194
• 3.7028942500, 4.5900196200, 5.2710064700, 0.4835832600,	STAP 195
• 4.2428901600, 2.3888473700, 3.7083908500, 4.9377721400,	STAP 196
• 4.8543625900, 0.8255372900, 2.3888473700, 4.9117216900,	STAP 197
• 4.9366590700, 5.0497895700, 4.2428901600, 0.4835832600,	STAP 198
• 5.3630387100, 3.7028942500, 4.5900196200, 6.2710064700,	STAP 199
• 3.7033908500, 4.8367977900, 2.5997399200, 4.8367966300,	STAP 200
• 0.8255372900, 2.5997399200, 0.8255958600, 4.8367233300,	STAP 201
• 5.0499218000, 3.7085923200, 1.9847250900, 0.8255959600,	STAP 202
• 4.9367246800, 5.0498217700, 3.7086198300, 1.9847250600,	STAP 203
• 5.2047634800, 4.6948783600, 4.6964403800, 4.4526176100,	STAP 204
• 5.0392327500, 4.7428920500, 4.8181256700, 5.0414559700/	STAP 205
DOUBLE PRECISION LON 2	STAP 206
DIMENSION LON 2(76)	STAP 207
EQUIVALENCE (LON (77),LON 2(1))	STAP 208
DATA LON 2/	STAP 209
• 5.0917424500, 5.2078834200, 4.5935696800, 4.5864477200,	STAP 210
• 0.1558104500, 0.4143549600, 6.2234366000, 2.4753138500,	STAP 211
• 5.1342554700, 4.2251878600, 4.8667850200, 4.8849533700,	STAP 212
• 4.9555007800, 4.8801485800, 4.4529161300, 4.9157552300,	STAP 213
• 4.8763559600, 0.4949384700, 4.8764594200, 5.2046959500,	STAP 214
• 5.0415948200, 4.3753229900, 4.9171537300, 4.4314026300,	STAP 215
• 4.4266789100, 4.4214025500, 4.4216314100, 4.4309277700,	STAP 216
• 4.4303558100, 4.1451770200, 4.1789950900, 4.1789971000,	STAP 217
• 4.1451601500, 4.1786402400, 4.7682714600, 4.7632683300,	STAP 218
• 4.1971776900, 4.1971752500, 3.4954071100, 3.4954077600,	STAP 219
• 4.2035037000, 4.2035738900, 4.1971630300, 4.1071526100,	STAP 220
• 4.1971412300, 4.2035559800, 4.2035400700, 3.4945102300,	STAP 221
• 4.2249006400, 4.2745365700, 4.2745407300, 4.9563231600,	STAP 222
• 4.9663230200, 4.9584228800, 4.9584228800, 5.1547650300,	STAP 223

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

• 0.8257977900.	3.4964966000.	5.1547640600.	1.9847397900.	STAP 224	
• 4.9557190000.	4.9652965100.	2.3882757200.	6.2034523500.	STAP 225	
• 6.2734428000.	6.2628070200.	4.9828412300.	4.1789870900.	STAP 226	
• 4.9335471100.	4.2388596400.	4.2003567600.	4.6150211600.	STAP 227	
• 4.6954376400.	3.5523405400.	4.9652904000.	4.8595799500.	STAP 228	
DOUBLE PRECISION LON 3				STAP 229	
DIMENSION LON 3(76)				STAP 230	
EQUIVALENCE (LON (153),LON 3(1))				STAP 231	
DATA LON 3/				STAP 232	
• 4.8657695200.	4.9804233100.	4.9423236900.	0.6962166900.	STAP 233	
• 0.7765713200.	0.5323596000.	0.6803002500.	1.4460352900.	STAP 234	
• 0.6933153600.	0.4654211300.	1.2075351300.	4.9423290100.	STAP 235	
• 4.5192343500.	4.8629945900.	4.5900196200.	4.5669820300.	STAP 236	
• 4.6738989900.	5.1547107300.	5.1312820500.	4.9422898000.	STAP 237	
• 4.9422977100.	4.7883901600.	4.4574665200.	4.8949465400.	STAP 238	
• 4.8849473800.	4.8849485300.	4.8849487200.	4.8703917600.	STAP 239	
• 4.9426130800.	4.9420830500.	4.9652047700.	1.98482058200.	STAP 240	
• 4.9422800900.	4.8367247000.	4.9652869500.	4.9422845600.	STAP 241	
• 1.9847200000.	4.3479951300.	4.3479932000.	4.9422822700.	STAP 242	
• 4.9422750100.	6.1748530500.	4.4234815200.	0.4930144300.	STAP 243	
• 5.0353039500.	4.3479950700.	5.6694469900.	0.4176999700.	STAP 244	
• 0.1933445000.	0.0762868400.	0.1302956600.	6.2488280500.	STAP 245	
• 6.2253212000.	0.4142000400.	0.0997006400.	0.1274026500.	STAP 246	
• 0.0935995300.	2.4198500400.	0.0389108600.	6.2268212000.	STAP 247	
• 0.1923814400.	0.1523618300.	6.1748560600.	0.0996931200.	STAP 248	
• 4.4234821100.	0.4930184300.	2.3873331800.	6.1748500300.	STAP 249	
• 2.4335945400.	1.3667563000.	5.0353439500.	0.9166432400.	STAP 250	
• 5.0817419000.	4.8649509200.	5.1468630500.	3.5559829000.	STAP 251	
DOUBLE PRECISION LON 4				STAP 252	
DIMENSION LON 4(26)				STAP 253	
EQUIVALENCE (LON (229),LON 4(1))				STAP 254	
DATA LON 4/				STAP 255	
• 5.9778732100.	4.3479950700.	2.3889930700.	2.4293536200.	STAP 256	
• 0.6799556500.	5.6694469900.	5.1031259000.	5.6694500700.	STAP 257	
• 4.8949481200.	5.0342568300.	0.1302856800.	0.4199076400.	STAP 258	
• 0.3991835900.	0.4177078500.	4.2251877800.	4.3625577100.	STAP 259	
• 0.1876359500.	3.3246037600.	2.9752487200.	0.3115046200.	STAP 260	
• 4.2251898000.	0.2280415800.	0.6417029200.	0.4354858800.	STAP 261	
• 5.9778732100.	0.3115046100.			STAP 262	
DOUBLE PRECISION HT 1				STAP 263	
DIMENSION HT 1(95)				STAP 264	
EQUIVALENCE (HT (1),HT 1(1))				STAP 265	
DATA HT 1/				STAP 266	
• -45.000000.	-46.000000.	-66.000000.	-27.000000.	208.000000.	STAP 267
• 555.000000.	821.000000.	8.000000.	76.000000.	1149.000000.	STAP 268
• 1123.000000.	920.000000.	-28.000000.	-41.000000.	-6.000000.	STAP 269
• 983.000000.	936.000000.	1040.000000.	979.000000.	148.000000.	STAP 270
• 673.000000.	1410.000000.	825.000000.	775.000000.	-41.000000.	STAP 271
• -19.000000.	-53.693000.	-42.000000.	3556.913000.	50.703000.	STAP 272
• 713.892000.	48.000000.	156.367000.	203.162000.	90.410000.	STAP 273
• 1540.977000.	876.254000.	130.403000.	283.125000.	-53.693000.	STAP 274
• -42.000000.	1359.777000.	130.403000.	3556.913000.	50.703000.	STAP 275
• 713.492000.	876.254000.	1540.977000.	48.000000.	156.367000.	STAP 276
• 203.162000.	90.410000.	283.125000.	849.933000.	749.578000.	STAP 277
• 850.063000.	1359.777000.	949.578000.	1382.000000.	918.180000.	STAP 278
• 727.193000.	340.399000.	1.153000.	1381.000000.	314.000000.	STAP 279

• 727.00000.	340.00000.	1.00000.	-59.37800.	-15.90900.	STAP 280
• -16.66400.	2141.20100.	30.06400.	17.43000.	7.44900.	STAP 281
• -61.08200.	-40.73100.	245.51500.	243.17100.	255.67800.	STAP 282
• 368.55500.	59.74100.	635.09400.	69.14900.	-13.33200.	STAP 283
• 752.74400.	-45.05800.	-36.71700.	-54.78500.	-49.89500.	STAP 284
• 1840.19900.	-41.90700.	-40.37800.	1609.20600.	-38.82200/	STAP 285
DOUBLE PRECISION HT 2					STAP 286
DIMENSION HT 2(95)					STAP 287
EQUIVALENCE (HT (96),HT 2(1))					STAP 288
DATA HT 2/					STAP 289
• -3.66000.	-27.03600.	-42.79200.	-45.56900.	1217.35100.	STAP 290
• 1185.61400.	1217.55100.	1484.31300.	1577.09300.	1225.94100.	STAP 291
• 3.27600.	599.25700.	599.23700.	-8.03200.	60.80600.	STAP 292
• -25.14700.	-26.76700.	220.97000.	221.26000.	447.41900.	STAP 293
• 442.98700.	-47.35000.	-47.55800.	218.97100.	220.01200.	STAP 294
• 219.43300.	-47.30700.	-47.30600.	-15.30900.	765.45500.	STAP 295
• 2787.74500.	2776.33600.	-54.06600.	-54.06600.	-60.47400.	STAP 296
• -60.47400.	-36.79500.	1320.10700.	1127.48300.	-35.55200.	STAP 297
• 12.30400.	-50.28500.	-47.72400.	124.34800.	182.55700.	STAP 298
• 182.55500.	41.87300.	-49.62500.	665.88700.	68.33500.	STAP 299
• 76.65300.	322.16000.	426.20100.	-17.16400.	28.27400.	STAP 300
• -49.29300.	-29.38300.	-43.25900.	-43.71600.	-15.25400.	STAP 301
• 150.00000.	960.00000.	184.00000.	40.00000.	150.00000.	STAP 302
• 114.00000.	75.00000.	440.80000.	-14.65400.	293.50800.	STAP 303
• 231.43100.	203.15200.	7.78200.	212.51800.	-27.00000.	STAP 304
• -18.36000.	-6.19300.	-6.09300.	124.72000.	1745.43100.	STAP 305
• -37.67500.	-37.42500.	-38.16600.	-37.51200.	221.00000.	STAP 306
• 404.75600.	-8.80000.	-55.11900.	-12.94100.	3.00000.	STAP 307
• 819.68900.	-60.00000.	-5.20700.	-5.36100.	2317.83300/	STAP 308
DOUBLE PRECISION HT 3					STAP 309
DIMENSION HT 3(64)					STAP 310
EQUIVALENCE (HT (191),HT 3(1))					STAP 311
DATA HT 3/					STAP 312
• 2317.71100.	-4.85600.	177.60600.	38.30000.	1000.80000.	STAP 313
• 1570.20000.	2490.00000.	2339.14100.	25.39000.	490.32800.	STAP 314
• 91.55300.	45.54000.	933.22000.	137.02000.	309.62700.	STAP 315
• 133.59200.	694.32000.	405.22000.	674.87100.	6.29500.	STAP 316
• 190.01000.	309.62700.	961.70800.	193.81900.	55.87500.	STAP 317
• 686.09200.	1615.00000.	1570.00000.	158.11500.	55.44000.	STAP 318
• 80.00000.	1856.00000.	2490.00000.	1563.80000.	-24.00000.	STAP 319
• -23.00000.	636.54000.	3031.81600.	171.00000.	2339.05000.	STAP 320
• 138.40600.	879.00000.	1901.30000.	25.39000.	234.06400.	STAP 321
• 44.00000.	-39.78600.	131.45400.	933.00000.	-14.56000.	STAP 322
• 204.69000.	490.32000.	743.44400.	654.00000.	595.04000.	STAP 323
• -7.00000.	1011.00000.	144.00000.	729.17400.	122.06400.	STAP 324
• 133.92800.	40.00000.	170.67200.	144.01000/		STAP 325
END					STAP 326

START

DESCRIPTION

START returns the array of back values of accelerations plus the values of the first and second sums needed for the integration routine. It iterates using interpolation until the sums converge. The arguments used for iteration are the epoch position and velocity arrays and initial values assigned to the variational partials. Initial predictions are made with a Taylor series approximation.

NAME START
PURPOSE TO START INTEGRATION PROCESS USING INTERPOLATOR
 FORMULAS AND ITERATING UNTIL DESIRED ACCURACY IS
 ACHIEVED

CALLING SEQUENCE CALL START(IORDER,H,FCT,SUM,Y,NN,M1,M2,TIM)

SYMBOL	TYPE	DESCRIPTION
IORDER	I	INPUT - ORDER
H	DP	INPUT - STEPSIZE
FCT (3,1)	DP	OUTPUT - ARRAY OF BACK VALUES OF ACCELERATION
SUM (2,3,1)	DP	OUTPUT - SUM ARRAY USED BY INTEGRATOR AND INTERPOLATOR.
Y (6,1)	DP	INPUT - ARRAY OF STATE AND PARTIALS
NN	I	INPUT - NUMBER OF EQUATIONS
M1	I	INPUT - DISPLACEMENT USED BY COWELL
M2	I	INPUT - DISPLACEMENT USED BY COWELL
TIM	DP	INPUT - EPOCH TIME OUTPUT - INTEGRATOR TIME

SUBROUTINES USED CLEAR F VEVAL COEF DOTPRD
 ERROR

COMMON BLOCKS INTBLK

INPUT FILES NONE

OUTPUT FILES PRINTER

REFERENCES *GEODYN SYSTEMS DESCRIPTION*
 VOLUME 1 - GEODYN DOCUMENTATION

SUBROUTINE START(IORDER,H,FCT,SUM,Y,NN,M1,M2,TIM)	STAR	48
IMPLICIT REAL*8(A-H,O-Z)	STAR	40
DIMENSION Y(6,1),AUX(6)	STAR	50
DIMENSION FCT(3,1),SUM(2,3,1),C(15,15,2)	STAR	51
COMMON/INTBLK/THROT(3),GY,AE(62)	STAR	53
DIMENSION A1(15,15),AS1(15,15)	STAR	54
EQUIVALENCE(C(1,1,1),AS1(1,1)),(C(1,1,2),A1(1,1))	STAR	54
DATA EPS/1.00-13/,MAXK/20/	STAR	55

LOGICAL SWITCH,NEOSW	STAR	56
IKNT=0	STAR	57
C SAVE EPOCH TIME	STAR	58
T=TIM	STAR	59
NEOSW=NN.GT.1	STAR	60
IOL2=IORDER-2	STAR	61
IOL1=IORDER-1	STAR	62
IEVEN=((IORDER+2)/2)*2-IORDER-1	STAR	63
MI=IOL2/2-IEVEN	STAR	64
MID=M1+1+IEVEN	STAR	65
MIDP2=MID+2	STAR	66
IST=M1-IOL2	STAR	67
ISTV=M2-IOL2	STAR	68
CALL CLEAR(SUM,12,NN)	STAR	69
M=IST+MID	STAR	70
MV=ISTV+MID	STAR	71
DO 5 J=1,6	STAR	72
5 AUX(J)=Y(J,1)	STAR	73
IF(.NOT.NEOSW)GOTO 7	STAR	74
DO 6 J=1,6	STAR	75
JPI=J+1	STAR	76
DO 4 N=2,NN	STAR	77
4 Y(J,N)=0.00	STAR	78
Y(J,JPI)=1.00	STAR	79
6 CONTINUE	STAR	80
C ONE TIME CALL AT EPOCH	STAR	81
7 CALL F(T,Y,FCT(1,4)..FALSE.)	STAR	82
1 FORMAT(1H0,6G20,10)	STAR	83
IF(NEOSW 1CALL VEVAL(Y,FCT(1,M1+MV),6..TRUE..M2)	STAR	84
C COMPUTE INTERPOLATOR COEFFICIENTS FOR EACH OF THE BACK VALUE POINTS	STAR	85
C IN ARRAY FCT	STAR	86
DO 10 I=1,IOL2	STAR	87
K=I+2	STAR	88
S=DFLOAT(I-IOL2)	STAR	89
CALL COEF(S,IORDER,A1(1,K),AS1(1,K))	STAR	90
10 CONTINUE	STAR	91
S=-DFLOAT(IOL2)	STAR	92
CALL COEF(S,IORDER,A1(1,2),AS1(1,2))	STAR	93
TIM=T	STAR	94
C PREDICT FORWARD USING TAYLOR SERIES	STAR	95
DO 200 I=1,MI	STAR	96
J=M+I	STAR	97
JV=MV+I	STAR	98
R0=DSQRT(DOTPRD(Y,Y))	STAR	99
R1=DOTPRD(Y,Y(4,1))/R0	STAR	100
DO 19 K=1,3	STAR	101
KP3=K+3	STAR	102
GERK=-GM*(Y(KP3,1)-3.C3*R1*Y(K,1)/R0)/R0**3	STAR	103
Y(K,1)=Y(K,1)+H*(Y(KP3,1)+H*(FCT(K, J-1)*.5D0+H*GERK/6.0D1))	STAR	104
19 Y(KP3,1)=Y(KP3,1)+H*(FCT(K, J-1)+H*.5D0*GERK)	STAR	105
TIM=TIM+H	STAR	106
CALL F(TIM,Y,FCT(1,J)..FALSE.)	STAR	107
IF(.NOT.NEOSW)GO TO 200	STAR	108
K0=M1+JV-1	STAR	109
DO 799 N=2,NN	STAR	110
KK=K0+(N-2)*M2	STAR	111

```

DO 799 K=1,3
  KP3=K+3
  Y(K,N)=Y(K,N)+H*(Y(KP3,N)+H*FCT(K,KK))*500
799 Y(KP3,N)=Y(KP3,N)+H*FCT(K,KK)
  CALL VEVAL(Y,FCT(1,K0+1),6,.TRUE.,M2)
200 CONTINUE
C PREDICT BACKWARD USING TAYLOR SERIES
DO 205 K=1,6
  Y(K,1)=AUX(K)
205 CONTINUE
  IF(.NOT.NEQSW)GO TO 207
  DO 209 J=1,6
  DO 208 N=2,NN
208 Y(J,N)=0.00
209 Y(J,J+1)=1.00
207 CONTINUE
  TIM=T
  DO 21 I=1,MID
  J=M-I
  JV=MV-I
  R0=DSORT(DOTPRD(Y,Y))
  R1=DOTPRD(Y,Y(4,1))/R0
  DO 20 K=1,3
  KP3=K+3
  GERK=-GM*(Y(KP3,1)-3.0)*R1*Y(K,1)/R0/P0**3
  Y(K,1)=Y(K,1)-H*(Y(KP3,1)-H*(FCT(K,J+1)*500-H*GERK/6.00))
20 Y(KP3,1)=Y(KP3,1)-H*(FCT(K,J+1)-H*500*GERK)
  TIM=TIM-H
  CALL F(TIM,Y,FCT(1,J),.FALSE.)
  IF(.NOT.NEQSW)GO TO 21
  K0=M1+JV+1
  DO 800 N=2,NN
  KK=K+(N-2)*M2
  DO 800 K=1,3
  KP3=K+3
  Y(K,N)=Y(K,N)-H*(Y(KP3,N)-H*FCT(K,KK))*500
800 Y(KP3,N)=Y(KP3,N)-H*FCT(K,KK)
  CALL VEVAL(Y,FCT(1,K0-1),6,.TRUE.,M2)
21 CONTINUE
C RESET EPDCH VALUES
DO 206 K=1,6
  Y(K,1)=AUX(K)
206 CONTINUE
  IF(.NOT.NEQSW)GO TO 22
  DO 2206 J=1,6
  DO 2207 N=2,NN
2207 Y(J,N)=0.00
2206 Y(J,J+1)=1.00
22 CONTINUE
  SWITCH=.FALSE.
  KOUNT=0
C COMPUTE SUMS
23 DO 30 N=1,NN
  K3=0
  IF(N.GT.1)K3=1
  K4=K3+1

```

STAR 112
STAR 113
STAR 114
STAR 115
STAR 116
STAR 117
STAR 118
STAR 119
STAR 120
STAR 121
STAR 122
STAR 123
STAR 124
STAR 125
STAR 126
STAR 127
STAR 128
STAR 129
STAR 130
STAR 131
STAR 132
STAR 133
STAR 134
STAR 135
STAR 136
STAR 137
STAR 138
STAR 139
STAR 140
STAR 141
STAR 142
STAR 143
STAR 144
STAR 145
STAR 146
STAR 147
STAR 148
STAR 149
STAR 150
STAR 151
STAR 152
STAR 153
STAR 154
STAR 155
STAR 156
STAR 157
STAR 158
STAR 159
STAR 160
STAR 161
STAR 162
STAR 163
STAR 164
STAR 165
STAR 166
STAR 167



```

      ISTD=K3*(ISTV-IST)+IST
      K0=K3*M1+(N-K4)*M2+ISTD+IOL1
      DO 30 J=1,3
      A=C.D0
      B=C.D0
      DO 29 I=1,IOL2
      KK=K0-I
      A=A-C(I,MID+2,1)*FCT(J,KK)
      B=B-C(I,MID+2,2)*FCT(J,KK)
29  CONTINUE
      A=A-C(IOL1,MID+2,1)*FCT(J,K0-IOL1)
      A=Y(J+3,N)/H+A
      B=Y(J,N)/H**2+DFLOAT(I+M1)*A+B
C CHECK IF SUMS HAVE REACHED DESIRED ACCURACY
      DIFF1=DABS(A-SUM(1,J,N))
      SUM(1,J,N)=A
      DIFF2=DABS(B-SUM(2,J,N))
      SUM(2,J,N)=B
      IF(DABS(A).GT.1.E-50)DIFF1=DIFF1/DABS(A)
      IF(DABS(B).GT.1.E-50)DIFF2=DIFF2/DABS(B)
      IF(DIFF1.GT.EPS)SWITCH=.TRUE.
      IF(DIFF2.GT.EPS)SWITCH=.TRUE.
30  CONTINUE
C USING ABOVE SUMS, RECOMPUTE POSITION & VELOCITY & ACCELERATION
      KOUNT=KOUNT+1
      II=IORDER+1-KOUNT
      IF(II.EQ.MIDP2)GOTO 50
      I=IST+IOL1-KOUNT
      I2=-KOUNT
      DO 45 N=1,NN
      K3=0
      IF(N.GT.1)K3=1
      K4=K3+1
      ISTD=K3*(ISTV-IST)+IST
      K0=K3*M1+(N-K4)*M2+ISTD+IOL1
      DO 45 J=1,3
      A=C.D0
      B=C.D0
      DO 44 K=1,IOL2
      KK=K0-K
      A=A+C(K,II,1)*FCT(J,KK)
      B=B+C(K,II,2)*FCT(J,KK)
44  CONTINUE
      A=A+C(IOL1,II,1)*FCT(J,K0-IOL1)
      A=A+SUM(1,J,N)
      B=B+SUM(1,J,N)*DFLOAT(I2)+SUM(2,J,N)
      Y(J,N) = B*H**2
      Y(J+3,N)= A*H
45  CONTINUE
      TIM=T+DFLOAT(I-M)*H
      CALL F(TIM,Y,FCT(1,I),.FALSE.)
      IF(NEOSW)CALL VEVAL(Y,FCT(1,N1+ISTV+IOL1-KOUNT),5,.TRUE.,M2)
      DO 52 J=1,6
52  Y(J,1)=AUX(J)
      IF(.NOT.NEOSW)GOTO 50
      DO 53 J=1,6

```

STAP 169
 STAR 169
 STAR 170
 STAR 171
 STAR 172
 STAR 173
 STAR 174
 STAR 175
 STAR 176
 STAR 177
 STAR 178
 STAR 179
 STAR 180
 STAR 181
 STAR 182
 STAR 183
 STAR 184
 STAR 185
 STAR 186
 STAR 187
 STAR 188
 STAR 189
 STAR 190
 STAR 191
 STAR 192
 STAR 193
 STAR 194
 STAR 195
 STAR 196
 STAR 197
 STAR 198
 STAR 199
 STAR 200
 STAR 201
 STAR 202
 STAR 203
 STAR 204
 STAR 205
 STAR 206
 STAR 207
 STAR 208
 STAR 209
 STAR 210
 STAR 211
 STAR 212
 STAR 213
 STAR 214
 STAR 215
 STAR 216
 STAR 217
 STAR 218
 STAR 219
 STAR 220
 STAR 221
 STAR 222
 STAR 223

DO 55 K=2,NN	STAR 224
55 Y(J,K)=0.00	STAR 225
53 Y(J,J+1)=1.00	STAR 226
59 IF(KOUNT.LT.10L1)GO TO 23	STAR 227
IKNT=IKNT+1	STAR 228
C EXIT LOOP IF ACCURACY HAS NOT BEEN REACHED AFTER MAXK TRIES	STAR 229
IF(IKNT.GT.MAXK) GO TO 59	STAR 230
C RECOMPUTE SUMS	STAR 231
IF(SWITCH) GO TO 22	STAR 232
GO TO 6)	STAR 233
59 PRINT 61,IKNT,EPS,DIFF1,DIFF2	STAR 234
CALL ERROR(10,DIFF1)	STAR 235
61 FORMAT(1H1,2CX,'INTEGRATION STARTING SUMS NOT CONVERGED AFTER',	STAR 236
• 13,' ITERATIONS'/1H0,15X,'EPS =',D22.16/10X,'DIFF1 =',	STAR 237
• D22.16/16X,'DIFF2 =',D22.16/1H0,2CX,'EXECUTION CONTINUING'///)	STAR 238
60 TIM=T+OFLOAT(MI)*H	STAR 239
RETURN	STAR 240
END	STAR 241



59

PRINT NOT CON-
VERGED MESSAGE;
EXECUTION CON-
TINUING,

ERROR
WRITE ERROR
MESSAGE

NAME STORE
 PURPOSE TO STORE COMMON AND ARC INFORMATION ON DISK
 CALLING SEQUENCE CALL STORE(RECALL,COMPAR)

SYMBOL	TYPE	DESCRIPTION
RECALL	L	INPUT - .FALSE. - STORE INFORMATION .TRUE. - RETRIEVE INFORMATION
COMPAR	L	INPUT - .FALSE. = APC PARAMETER INFORMATION .TRUE. = COMMON PARAMETER INFORMATION

SUBROUTINES USED - NONE

COMMON BLOCKS	APARAM	CELEM	CTIME	FLXBLK	FMODEL
	INITBK	INTBLK	PREBLK	PRIOR!	TPEBLK
	VRBLK				

INPUT FILES SCRC - SCRATCH

OUTPUT FILES SCRC - SCRATCH

SUBROUTINE STORE(RECALL,COMPAR)	STOP	28
IMPLICIT REAL*8 (A-H,C-Z)	STOP	29
LOGICAL RECALL,COMPAR,MISLOG	STOP	30
INTEGER XYZTF,FLCTP,SCRC,FLTP,THETGO,ELEMST	STOP	31
REAL DAYREF	STOP	32
DOUBLE PRECISION MODEL	STOP	33
COMMON/APARAM/INPAR(10)	STOP	34
COMMON/CELEM/ELEMST(53)	STOP	35
COMMON/CTIME/DAYREF(23)	STOP	36
COMMON/FLXBLK/ISTORE(450,9)	STOP	37
COMMON/FMODEL/INDEX(4),CS(30,33),MODEL(8)	STOP	38
COMMON/INITEK/IEPYMD(43),MISLOG(9)	STOP	39
COMMON/INTBLK/THDOT1(27),THETGO(78)	STOP	40
COMMON/PREBLK/DAYSTA,NOPS(15)	STOP	41
COMMON/PRIORI/ELEMIN(56)	STOP	42
COMMON/TPEBLK/INTP(3),XYZTP(3),PLOTP(3),SCRC,FLTP(2)	STOP	43
COMMON/VRBLK/JSTORE(450,5)	STOP	44
IF(RECALL) GO TO 500	STOP	45
DO 100 I=1,9	STOP	46
100 WRITE(SCRC) (ISTORE(J,I),J=1,450)	STOP	47
DO 200 I=1,5	STOP	48
200 WRITE(SCRC), (JSTORE(J,I),J=1,450)	STOP	49
IF(COMPAR) RETURN	STOP	50
WRITE(SCRC) DAYREF,INDEX,DAYSTA,XYZTP,THETGO,ELEMIN,INPAR,IEPYMD,	STOP	51
ELEMST	STOP	52
RETURN	STOP	53
500 DO 700 I=1,9	STOP	54
700 READ(SCRC) (ISTORE(J,I),J=1,450)	STOP	55

DO 800 I=1.5	STOP	56
800 READ(SCRC) (JSTORE(J, I), J=1, 450)	STOP	57
IF(CMPAR) RETURN	STOP	58
READ(SCRC) DAYREF, INDEX, DAYSTA, XYZTP, THE TGO, FLEM(N), INPAR, IEPYMD,	STOP	59
• ELEMST	STOP	60
RETURN	STOP	61
END	STOP	62

NAME SUMMARY
PURPOSE TO PRINT ARC STATISTICAL SUMMARY
CALLING SEQUENCE CALL SUMMARY(ARCNO, INNER, OUTER, LINNER, EDIT, NAME)

SYMBOL	TYPE	DESCRIPTION
ARCNO	I	INPUT - ARC NUMBER
INNER	I	INPUT - INNER ITERATION NUMBER
OUTER	I	INPUT - OUTER ITERATION NUMBER
LINNER	L	INPUT - .TRUE. - LAST INNER ITERATION
EDIT	R	INPUT - EDITING LEVEL
NAME	I	INPUT - STATION NAMES

(1)

SUBROUTINES USED STAINF EPROR
COMMON BLOCKS ALPMRC APARAM CELEM CONSTS CPARAM
 CSTINF INITBK TPERLK
INPUT FILES NONE
OUTPUT FILES OUTP - PRINTER

SUBROUTINE SUMMARY(ARCNO, INNER, OUTER, LINNER, EDIT, NAME)	SUMM	34
LOGICAL*1 BYTE, SLASH	SUMM	35
LOGICAL CMPGPR, SWITCH, LINNER, HYPER	SUMM	36
INTEGER*2 ISAT	SUMM	37
INTEGER RECNO1, OUTP, DATP, ARCNO, OUTER	SUMM	38
DOUBLE PRECISION ITNMS, TIMING, BLANK, ATYPE, UNITS, ELEMST, ORBELA,	SUMM	39
• EPSEC, ORBEL, CONFIG, NAME, ELCUT	SUMM	40
DIMENSION BYTE(8), NAME(1)	SUMM	41
COMMON/ALPMRC/ITNMS(5), TIMING, BLANK, ATYPE(31), UNITS(15), ELCUT,	SUMM	42
• HYPER	SUMM	43
COMMON/APARAM/INPAR, INPARI, NBIAS, ESTSTA, NSAT, NGPARC, RECNO1, NPARAM,	SUMM	44
• NBIAS, MAXPAR	SUMM	45
COMMON/CELEM/ELEMST(6,2), ORFLA(6,2), IG15(4), RMSTOT	SUMM	46
COMMON/CONSTS/PI, P11, T40P1, T40P11, DRAD, DRAD1, RSEC, RSEC1	SUMM	47
COMMON/CPARAM/NSTA, NMAST, NSTEST, NDIM, MBIAS, NGPC1, NGPC2, NGPCOM,	SUMM	48
• ACSEST, CMPGPR, LIM1, LIM2, NOEN, NDNST, NTIDST, NTIDEN, INNRSW,	SUMM	49
• NCONST, NDCONS	SUMM	50
COMMON/CSTINF/MEASND(4), NOTS(4), FDMEAN(4), RMSD(4), PND(4),	SUMM	51
• MEASNT(4), WTRMEAN(4), RMSWTO(4), WTRND(4), TYPRMS(30), NUTYPE(2,30),	SUMM	52
• BSUM(4,12), RMSNJC(30), NOALL(30), NOWTOU, LBASE	SUMM	53
COMMON/INITBK/IEPYMD, IEPHM, CPSEC, IYREF, INNMAX, INNMIN, CONVRG,	SUMM	54
• ORBEL(6,2), EDITN, INSUPR, ICSAT(2), SWITCH(21)	SUMM	55

COMMON/TPEBLK/INTP,OUTP,DATP(10)	SUMM 56
EQUIVALENCE (CONFIG,BYTE(1))	SUMM 57
DATA ASTRSK/1H*//,SLASH/1H//	SUMM 58
L1=LBASF+1	SUMM 59
ISAT=1	SUMM 60
C CALCULATE STATISTICS	SUMM 61
CALL STAINF(3,C,P,P,P,P,ISAT,P)	SUMM 62
C LOOP THROUGH 1500 FOR EACH SATELLITE	SUMM 63
DO 1500 ISAT=1,NSAT	SUMM 64
WRITE(OUTP,5000) ISAT,ARCNO,INNER,OUTER	SUMM 65
LINE=1	SUMM 66
NOBS(1)=0	SUMM 67
IF(NSTA.LE.0) GO TO 1600	SUMM 68
C LOOP THROUGH 1500 FOR EACH STATION (OR BASELINE)	SUMM 69
DO 1500 J=1,NSTA	SUMM 70
DO 1500 LP1=1,L1	SUMM 71
L=LP1-1	SUMM 72
IF(NORS(1).EQ.0) GO TO 100	SUMM 73
WRITE(OUTP,5100)	SUMM 74
LINE=LINE+1	SUMM 75
C OBTAIN STATISTICS FOR EACH STATION (OR BASE LINE FOR TWO STATION	SUMM 76
C DATA) AND EACH SATELLITE	SUMM 77
100 CALL STAINF(3,J,P,P,P,P,ISAT,L)	SUMM 78
C LOOP THROUGH 1400 FOR EACH MEASUREMENT TYPE	SUMM 79
DO 1400 K=1,4	SUMM 80
IF(NOBS(K).EQ.0) GO TO 1500	SUMM 81
IF(LINE.LT.40) GO TO 200	SUMM 82
WRITE(OUTP,5200)	SUMM 83
WRITE(OUTP,5000) ISAT,ARCNO,INNER,OUTER	SUMM 84
LINE=1	SUMM 85
200 LINE=LINE+1	SUMM 86
MTYPE=MEASNO(K)	SUMM 87
C CONVERT UNITS FOR OUTPUT	SUMM 88
IF(MTYPE.LT.14) GO TO 300	SUMM 89
M=MTYPE-26	SUMM 90
GO TO (400,450,700,700),M	SUMM 91
300 M=MTYPE-(MTYPE/8)*7	SUMM 92
GO TO (500,800,700,700,600,500,500),M	SUMM 93
400 RDMEAN(K)=RDMEAN(K)*1.0E+09	SUMM 94
RMSO(K)=RMSO(K)*1.0E+09	SUMM 95
GO TO 800	SUMM 96
450 RDMEAN(K)=RDMEAN(K)*1.0E+6	SUMM 97
RMSO(K)=RMSO(K)*1.0E+6	SUMM 98
GO TO 800	SUMM 99
500 RDMEAN(K)=RDMEAN(K)/RSEC	SUMM 100
RMSO(K)=RMSO(K)/RSEC	SUMM 101
GO TO 800	SUMM 102
600 RDMEAN(K)=RDMEAN(K)*1.0E+03	SUMM 103
RMSO(K)=RMSO(K)*1.0E+03	SUMM 104
GO TO 800	SUMM 105
700 IF(MTYPE.EQ.4) GO TO 800	SUMM 106
RDMEAN(K)=RDMEAN(K)*1.0E+02	SUMM 107
RMSO(K)=RMSO(K)*1.0E+02	SUMM 108
800 CONFIG=PLANK	SUMM 109
IF(MTYPE.LT.27) GO TO 900	SUMM 110
CONFIG=NAME(L)	SUMM 111

```



        BYTE(8)=SLASH
C PRINT STATISTICS
900 WRITE(OUTP,5300) CONFIG,NAME(J),ATYPE(MTYPE),NOBS(K),RDMEAN(K)
    IF(NOBS(K).LT.10) GO TO 1000
    A1=BLANK
    A2=BLANK
    IF(RND(K).GT.1.98) A1=ASTRSK
    IF(RND(K).GT.2.58) A2=1STRSK
    WRITE(OUTP,5400) RND(K),A1,A2,RMSO(K)
1000 IF(MEASWT(K).GT.0) WRITE(OUTP,5500) MEASWT(K),WTMEAN(K)
    IF(MEASWT(K).LT.10) GO TO 1100
    A1=BLANK
    A2=BLANK
    IF(WTRND(K).GT.1.98) A1=ASTRSK
    IF(WTRND(K).GT.2.58) A2=ASTRSK
    WRITE(OUTP,5600) WTRND(K),A1,A2,RMSWT0(K)
1100 NOWTDB=NOWTDB+MEASWT(K)
1400 CONTINUE
1500 CONTINUE
1600 DO 1700 I=15,26
1700 NOWTCB=NOWTCB+NOTYPE(2,I)
    ISAT=1
    IF(NOWTDB.LT.8) CALL EPROR(5,BLANK)
C COMPUTE STATISTICS BY MEASUREMENT TYPE ONLY
    CALL STAINF(4,NPARAM,NM,RMSTOT,P,LINNER,P,P)
    IF(NSTA.LE.0) GO TO 1800
C WRITE TOTAL RMS
    WRITE(OUTP,5700) NM,RMSTOT
    WRITE(OUTP,5200)
C PRINT EDIT LEVEL
1800 IF(EDITN.LT.100.) WRITE(OUTP,5800) EDIT
C PRINT SUMMARY BY MEASUREMENT TYPE
    WRITE(OUTP,5900) ARCNC,INNER,OUTFR
    DO 1900 I=1,7
    IF(NOTYPE(2,I).GT.0) WRITE(OUTP,6000) ATYPE(I),NOTYPE(2,I),
        TYPRMS(I)
        J=I+7
    IF(NOTYPE(2,J).GT.0) WRITE(OUTP,6000) ATYPE(J),NOTYPE(2,J),
        TYPRMS(J)
1900 IF(NOTYPE(2,I)+NOTYPE(2,J).GT.0) WRITE(OUTP,5100)
    DO 2000 I=15,30
    IF(NOTYPE(2,I).LE.0) GO TO 2000
    WRITE(OUTP,6000) ATYPE(I),NOTYPE(2,I),TYPRMS(I)
    WRITE(OUTP,5100)
2000 CONTINUE
    WRITE(OUTP,6100) NM,RMSTOT
    RETURN
5000 FORMAT(1H1,22X,9HSATELLITE,12,4H ARC,13,
    . 32H RESIDUAL SUMMARY BY STATION FOR,
    . 16H INNER ITERATION,13,19H OF OUTER ITERATION,12//
    1 14X,13HSTATION MEAS,13X,31HRESIDUALS FROM ALL OBSERVATIONS,
    2 8X,9HNO OF WTD,3X,10HMEAN RATIO,2X,9HWEIGHTED,3X,9HWEIGHTED/
    3 15X,4HNAME,4X,4HTYPE,9X,4HNUMBER,5X,4HMEAN,7X,7HRND,9X,3HMS,
    4 7X,7HRESIDUALS,4X,3HTO SIGMA,5X,3HRND,9X,5HRMS/)
5100 FORMAT(1X)
5200 FORMAT(1H0,12X,71H* - INDICATES RESIDUALS ARE SIGNIFICANTLY NON-RASUMM


```

SUMM 112
SUMM 113
SUMM 114
SUMM 115
SUMM 116
SUMM 117
SUMM 118
SUMM 119
SUMM 120
SUMM 121
SUMM 122
SUMM 123
SUMM 124
SUMM 125
SUMM 126
SUMM 127
SUMM 128
SUMM 129
SUMM 130
SUMM 131
SUMM 132
SUMM 133
SUMM 134
SUMM 135
SUMM 136
SUMM 137
SUMM 138
SUMM 139
SUMM 140
SUMM 141
SUMM 142
SUMM 143
SUMM 144
SUMM 145
SUMM 146
SUMM 147
SUMM 148
SUMM 149
SUMM 150
SUMM 151
SUMM 152
SUMM 153
SUMM 154
SUMM 155
SUMM 156
SUMM 157
SUMM 158
SUMM 159
SUMM 160
SUMM 161
SUMM 162
SUMM 163
SUMM 164
SUMM 165
SUMM 166
SUMM 167

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

•NDOM - 5 PERCENT LEVEL /12X.72H** - INDICATES RESIDUALS ARE SIGNIF SUMM 168
 •ICANTLY NON-RANDOM - 1 PERCENT LEVEL) SUMM 169
 5300 FORMAT(6X,A3,A7,1X,A6,8X,14,F12.3) SUMM 170
 5400 FORMAT(1H+,50X,F9.3,2A1,F9.3) SUMM 171
 5500 FORMAT(1H+,80X,14,3X,F11.3) SUMM 172
 5600 FORMAT(1H+,100X,F9.3,2A1,F9.3) SUMM 173
 5700 FORMAT(1H0,76X,3HALL,15,22H WEIGHTED MEASUREMENTS,F14.3) SUMM 174
 5800 FORMAT(1H0,1H,16X,50HFOR THIS ITERATION NO MEASUREMENTS WITH RESIDUALS .12HGREATER THAN .F9.2.6H SIGMA/1H .20X,25HWERE USED IN THE SOLUTION) SSUMM 175
 5900 FORMAT(1H1,23X,3HARC,13,38H RESIDUAL SUMMARY BY MEASUREMENT TYPE .SUMM 177
 • 19HFOR INNER ITERATION,13,19H OF OUTER ITERATION,12/ SUMM 178
 2 1H0,38X,11HMEASUREMENT,9X,1CHNUMBER OF . SUMM 179
 3 8HWEIGHTED,8X,8HWEIGHTED/1H,41X,4HTYPE,16X,9HRESIDUALS. SUMM 180
 4 15X,3HRMS/) SUMM 181
 6000 FORMAT(1H,40X,A6,15X,15X,F10.3) SUMM 182
 6100 FORMAT(1H,43X,11HRMS FOR ALL,15,22H WEIGHTED MEASUREMENTS,F10.3) SUMM 184
 END SUMM 185





SUNGRV

DESCRIPTION

Subroutine SUNGRV evaluates the accelerations of the satellite due to the gravitational potentials of the Moon, the Sun, and the Planets, Venus, Mars, Jupiter, and Saturn. Intermediate data is stored in COMMON MOONGR for including the effects of these potentials in the variational equation computations in VEVAL.

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

NAME SUNGRV

PURPOSE TO COMPUTE GRAVITATIONAL ACCELERATIONS DUE TO :
1) MOON
2) SUN
3) VENUS
4) MARS
5) JUPITER
6) SATURN

CALLING SEQUENCE CALL SUNGRV(DX)

SYMBOL TYPE DESCRIPTION
DX DP INPUT & OUTPUT - ACCELERATION VECTOR
(3)

SUBROUTINES USED DOTPRD

COMMON BLOCKS CEPHEM MOONGR INTBLK XYZ

INPUT FILES NONE

OUTPUT FILES NONE

```

SUBROUTINE SUNGRV(DX)                                SUNG 29
  IMPLICIT REAL*8 (A-H,C-Z)                          SUNG 30
  DIMENSION DX(3)                                    SUNG 31
  COMMON/CEPHEM/UVBODY(4,6),EQ(644)                  SUNG 32
  COMMON/MOONGR/DPXUV(6),RHO4(3,6),PHO50(6),RHO3(6)  SUNG 33
  COMMON/INTBLK/THDOT1(9),GM3(6),IR(101),NBODY      SUNG 34
  COMMON/XYZ/X(6),R,RSQ,ISAT,IFORCE(2)              SUNG 35
  DO 20 N=1,NBCDY                                    SUNG 36
    DPXUV(N)=DOTPRC(X,UVBODY(1,N))                   SUNG 37
    RRBCDY=UVBODY(4,N)**2                             SUNG 38
    RHOSQ(N)=RRBCDY-2.0DC*DPXUV(N)*UVBODY(4,N)+RSQ   SUNG 39
    RHO3(N)=RHOSC(N)*DSQRT(RHOSQ(N))                 SUNG 40
    DO 20 I=1,3                                       SUNG 41
      RHO4(I,N)=X(I)-UVBODY(I,N)+UVBODY(4,N)        SUNG 42
  20 DX(I)=DX(I)-GM3(N)*(RHO4(I,N)/RHO3(N)+UVBODY(I,N)/RRBCDY) SUNG 43
  RETURN                                             SUNG 44
  END                                               SUNG 45

```

SURDEN

DESCRIPTION

This subroutine computes the gravitational acceleration due to surface density blocks and the partial derivatives of this acceleration with respect to adjusted surface density parameters. These partials include the constraints affecting the adjustment of surface densities as described in the GEODYN Systems Description, Volume 1.

NAME SURDEN
 ENTRY FCINT PURPOSE
 SURDN1 INITIAL IZATION
 SURDEN TO COMPUTE :
 1) THE GRADIENT OF THE POTENTIAL DUE TO SURFACE DENSITIES
 2) THE PARTIALS OF THE GRADIENTS WITH RESPECT TO THE SURFACE DENSITIES FOR THOSE DENSITIES TO BE ADJUSTED

CALLING SEQUENCE CALL SURDN1(DENSE,AREA,CENTER,PART,DENCON)

SYMBOL	TYPE	DESCRIPTION
DENSE (1)	DP	INPUT - ARRAY CONTAINING THE SURFACE DENSITIES OF THE BLOCKS
AREA	DP	INPUT - ARRAY CONTAINING THE SURFACE AREAS OF THE SUB-BLOCKS
CENTER (3,1)	DP	INPUT - ARRAY CONTAINING THE EARTH FIXED X,Y,Z COORDINATES OF THE SUB-BLOCK CENTERS
PART (3,1)	DP	OUTPUT - ARRAY CONTAINING THE PARTIALS FOR THE BLOCKS TO BE ADJUSTED
DENCON (NCONST,1)	DP	INPUT - COEFFICIENTS RELATING CONSTRAINED AND UNCONSTRAINED ADJUSTED SURFACE DENSITIES




CALLING SEQUENCE CALL SURDEN(FCT,THETG)


SYMBOL	TYPE	DESCRIPTION
FCT (3)	DP	INPUT & OUTPUT - ARRAY CONTAINING THE INERTIAL X,Y,Z COORDINATES OF THE GRADIENT TO BE UPDATED
THETG	DP	INPUT - RIGHT ASCENSION OF GREENWICH

SUBROUTINES USED XEFIX YEFIX XINERT YINERT
 COMMON BLOCKS CPARAM CSTHET INTBLK XYZ
 INPUT FILES NONE
 OUTPUT FILES NONE
 REFERENCES *GEOODYN SYSTEMS DESCRIPTION*
 VOLUME 1 - GEOODYN DOCUMENTATION

C	SUBROUTINE SURDN1(DENSE,ARFA,CENTER,PART,DENCON)	SURD	56
	IMPLICIT REAL*8(A-H,O-Z)	SURD	57
	LOGICAL CMPGPR,INNRSW	SURD	58
	DIMENSION DENSE(1),AREA(1),CENTER(3,1),GRAD(3),PART(3,1),PAR(3),	SURD	59
	• PVEC(3),DP(3),FCT(3),DENCON(NCONST,1)	SURD	60
	COMMON/CPARAM/NSTA,NMAST,NSTEST,NDIM,MBIAS,NGPC1,NGPC2,NGPCOM,	SURD	61
	• NCSEST,CMPGPR,IM1,LIM2,NBLOCK,NADJ,NTIDST,NTIDEN,INNRSW,	SURD	62
	• NCCNST,NDCONS	SURD	63
	COMMON/CSTHET/CTHETG,STHETG	SURD	64
	COMMON/INTBLK/THOOT(5),NECN(16)	SURD	65
	COMMON/XYZ/POINT(8),ISAT,IFORCE(2).	SURD	66
	DATA NSUR/4/	SURD	67
C	INITIALIZE	SURD	68
	NCON2=NADJ-NCONST	SURD	69
	RETURN	SURD	70
	ENTRY SURDEN(FCT,THETG)	SURD	71
	IF(NBLOCK.LE.0) RETURN	SURD	72
	CTHETG=DCOS(THETG)	SURD	73
	STHETG=DSIN(THETG)	SURD	74
C	COMPUTE THE EARTH FIXED COORDINATES OF SATELLITE	SURD	75
	PVEC(1)=XEFIX(POINT(1),POINT(2))	SURD	76
	PVEC(2)=YEFIX(POINT(1),POINT(2))	SURD	77
	PVEC(3)=POINT(3)	SURD	78
	GRAD(1)=0.000	SURD	79
	GRAD(2)=0.000	SURD	80
	GRAD(3)=0.000	SURD	81
	NC=1	SURD	82
	K=NECN(ISAT)-7-NCON2	SURD	83
	K1=K+1	SURD	84
	K2=K+NCON2	SURD	85
	DO 100 KK=1,NBLOCK	SURD	86
C	LOOP THROUGH 100 FOR EACH DENSITY BLOCK	SURD	87
	K=K+1	SURD	88
	DO 20 J=1,3	SURD	89
	20 PAR(J)=0.00	SURD	90
C	LOOP THROUGH 30 COMPUTES EARTH FIXED SURFACE DENSITY PARTIALS FOR	SURD	91
C	THE BLOCK	SURD	92
	DO 50 I=1,NSUB	SURD	93
	DMAG=0.00	SURD	94
	DO 30 J=1,3	SURD	95
	DP(J)=PVEC(J)-CENTER(J,NC)	SURD	96
30	DMAG=DMAG+DP(J)**2	SURD	97
	C=AREA(NC)/(DMAG*DSORT(DMAG))	SURD	98
	DO 40 J=1,3	SURD	99
40	PAR(J)=PAR(J)-C*DP(J)	SURD	100
50	NC=NC+1	SURD	101
	DO 60 J=1,3	SURD	102
C	COMPUTE EARTH FIXED GRADIENT DUE TO BLOCK AND SUM	SURD	103
60	GRAD(J)=GRAD(J)+DENSE(KK)*PAR(J)	SURD	104
	IF(KK.GT.NADJ) GO TO 100	SURD	105
	IF(.NOT.INNRSW) GO TO 100	SURD	106
	IF(KK.GT.NCON2) GO TO 70	SURD	107
C	IF BLOCK ADJUSTED COMPUTE THE INERTIAL PARTIALS	SURD	108
	PART(1,K)=XINERT(PAR(1),PAR(2))	SURD	109
	PART(2,K)=YINERT(PAR(1),PAR(2))	SURD	110
		SURD	111

PART(3,K)=PAR(3)	SURD 112
GO TO 100	SURD 113
C IF BLOCK ADJUSTED THROUGH CONSTRAINTS COMPUTE CORRECTIONS TO	SURD 114
C PARTIALS FOR ADJUSTED DENSITIES AND SUM	SURD 115
70 DP(1)=XINERT(PAR(1),PAR(2))	SURD 116
DP(2)=YINERT(PAR(1),PAR(2))	SURD 117
DP(3)=PAR(3)	SURD 118
00 80 I=K1,K2	SURD 119
I1=I-K1+1	SURD 120
PCCDA=DENCON(KK-NCON2,I1)	SURD 121
PACT(1,I)=PART(1,I)+PCCDA*DP(1)	SURD 122
PART(2,I)=PART(2,I)+PCCDA*DP(2)	SURD 123
80 PART(3,I)=PART(3,I)+PCCDA*DP(3)	SURD 124
100 CONTINUE	SURD 125
C CONVERT GRADIENT DUE TO ALL BLOCKS TO INERTIAL AND UPDATE TOTAL	SURD 126
C FORCE MODEL GRADIENT	SURD 127
FCT(1)=FCT(1)+XINERT(GRAD(1),GRAD(2))	SURD 128
FCT(2)=FCT(2)+YINERT(GRAD(1),GRAD(2))	SURD 129
FCT(3)=FCT(3)+GRAD(3)	SURD 130
RETURN	SURD 131
END	SURD 132





SYMINV

DESCRIPTION

Subroutine SYMINV is a double precision matrix inversion routine designed specifically for inverting a compressed storage symmetric matrix such as is used by ESTIM.

The technique used is the method of partitioning. The initialization consists of inverting a 1×1 . The routine then constructs successively larger $(N \times N)$ inverted partitions of the original matrix until the entire matrix has been inverted.

This routine destroys the input matrix. It also requires a double precision scratch storage vector of length equal to the row dimension of the matrix to be inverted.

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

NAME SYMINV

PURPOSE TO RECURSIVELY FIND INVERSE OF SYMMETRIC MATRIX

CALLING SEQUENCE CALL SYMINV(SUM1,NDIM,NLIM,DELTA)

SYMBOL	TYPE	DESCRIPTION
SUM1 (1)	DP	INPUT - LOWER RECTANGULAR PART OF MATRIX TO BE INVERTED OUTPUT - LOWER RECTANGULAR PART OF INVERTED MATRIX
NDIM	I	INPUT - DIMENSION OF MATRIX
NLIM	I	INPUT - DIMENSION OF PARTITION TO BE INVERTED
DELTA (1)	DP	SCRATCH

SUBROUTINES USED NONE

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES *GEODYN SYSTEMS DESCRIPTION*
VOLUME 1 - GEODYN DOCUMENTATION

SUBROUTINE SYMINV(SUM1,NDIM,NLIM,DELTA)	
... DOUBLE PRECISION SUM1(1),DELTA(1)	SYMI 34
C INITIALIZE BY FINDING INVERSE OF 1X1	SYMI 35
SUM1(1)=1.000/SUM1(1)	SYMI 36
IF(NLIM.EQ.1) RETURN	SYMI 37
N1=NDIM-1	SYMI 38
C RECURSIVELY FIND INVERSE OF NXN KNOWING INVERSE OF (N-1)X(N-1) UNTIL	SYMI 39
C THE INVERSE OF AN NLIM X NLIM SQUARE PARTITION IS FOUND	SYMI 40
DO 400 N=2,NLIM	SYMI 41
NM1=N-1	SYMI 42
L1=0	SYMI 43
DO 100 L=1,NM1	SYMI 44
J1=0	SYMI 45
DELTA(L)=0.000	SYMI 46
DO 60 J=1,L	SYMI 47
JL=J1+L	SYMI 48
JN=J1+N	SYMI 49
DELTA(L)=DELTA(L)+SUM1(JL)*SUM1(JN)	SYMI 50
60 J1=J1+NDIM-J	SYMI 51
IF(L.EQ.NM1) GO TO 100	SYMI 52
LP1=L+1	SYMI 53
DO 80 J=LP1,NM1	SYMI 54
	SYMI 55

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

JN=J1+N	SYMI	56
JL=L1+J	SYMI	57
DELTA(L)=DELTA(L)+SUM1(JL)*SUM1(JN)	SYMI	59
80 J1=J1+NDIM-J	SYMI	59
100 L1=L1+NDIM-L	SYMI	60
J1=N	SYMI	61
NN=N1+N	SYMI	62
DO 150 J=1,NM1	SYMI	63
SUM1(NN)=SUM1(NN)-DELTA(J)*SUM1(J1)	SYMI	64
150 J1=J1+NDIM-J	SYMI	65
SUM1(NN)=1.000/SUM1(NN)	SYMI	66
J1=N	SYMI	67
DO 200 J=1,NM1	SYMI	68
SUM1(J1)=-DELTA(J)*SUM1(NN)	SYMI	69
200 J1=J1+NDIM-J	SYMI	70
I1=N	SYMI	71
DO 300 I=1,NM1	SYMI	72
J1=I	SYMI	73
DO 250 J=1,I	SYMI	74
SUM1(J1)=SUM1(J1)-SUM1(I1)*DELTA(J)	SYMI	75
250 J1=J1+NDIM-J	SYMI	76
300 I1=I1+NDIM-I	SYMI	77
400 N1=N1+NDIM-N	SYMI	78
RETURN	SYMI	79
END	SYMI	80

TDIF

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

DESCRIPTION

TDIF computes the differences in seconds between the time systems

UT1, UT2, UTC, and A.1.

Tabular information relating A.1 and UTC is required by TDIF and must periodically be updated. This data is available from the U.S. Naval Observatory.

Tabular information relating A.1 and UT1 is required by TDIF. This data also must be periodically update. A full description of the method used to update this table is available in GEODYN Support Programs, Volume IV.

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

NAME TDIF
PURPOSE TO COMPUTE THE DIFFERENCE IN SECONDS BETWEEN ANY TWO OF THE FOLLOWING TIME SYSTEMS
A.1, UTC, UT1, UT2

CALLING SEQUENCE X=TDIF(BASE,IN,DAY0)

SYMBOL	TYPE	DESCRIPTION
BASC	I	INPUT - DESIRED TIME SYSTEM : (1=UT1, 2=UT2, 3=UTC, 4=A.1)
IN	I	INPUT - CURRENT TIME SYSTEM : (1=UT1, 2=UT2, 3=UTC, 4=A.1)
DAYC	DP	INPUT - CURRENT TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR
TDIF	R	OUTPUT - COMPUTED TIME DIFFERENCE IN SECONDS

COMMON BLOCKS INITBK

REFERENCES *GEDDYN SYSTEMS DESCRIPTION*
VOLUME 1 - GEDDYN DOCUMENTATION

REAL FUNCTION TDIF(BASC,IN,DAYC)	TDIF	30
DOUBLE PRECISION DAY0,REFTIM,YMDAY,DAYS,UTCT,TWOPI	TDIF	31
INTEGER BASE,ID(4)	TDIF	32
COMMON/INITBK/IG1(55),NOT1ST,IG2	TDIF	33
LOGICAL NOT1ST	TDIF	34
DIMENSION UTCT(53),UTCDF(52),UTCRT(52),AIT1(613),9IT1(152),	TDIF	35
• CIT1(152),DIT1(152)	TDIF	36
EQUIVALENCE (BIT1(1),AIT1(158)),(CIT1(1),AIT1(310)),	TDIF	37
• (DIT1(1),AIT1(452))	TDIF	38
DATA ID/2,1,4,3/	TDIF	39
DATA TWOPI/6.283185307	TDIF	40
DATA LLIMIT/528/	TDIF	41
C TABLE A.1-UT1 -- AT 10 DAY INTERVALS FROM 570917 TO 720303	TDIF	42
• DATA AIT1/	TDIF	43
• -.1649,-.1513,-.1361,-.1193,-.1013,-.0937,-.0649,-.0458,-.0270,	TDIF	44
• -.0096,0.0098,0.0285,0.0457,0.0618,0.0784,0.0948,0.1111,0.1277,	TDIF	45
• 0.1453,0.1629,0.1815,0.1993,0.2176,0.2351,0.2517,0.2665,0.2799,	TDIF	46
• 0.2879,0.2991,0.3063,0.3131,0.3169,0.3249,0.3310,0.3378,0.3458,	TDIF	47
• 0.3554,0.3652,0.3794,0.3942,0.4098,0.4261,0.4431,0.4596,0.4765,	TDIF	48
• 0.4929,0.5093,0.5247,0.5401,0.5559,0.5727,0.5866,0.6000,0.6136,	TDIF	49
• 0.6267,0.6405,0.6548,0.6694,0.6893,0.6981,0.7115,0.7243,0.7364,	TDIF	50
• 0.7471,0.7572,0.7660,0.7734,0.7806,0.7889,0.7977,0.8070,0.8094,	TDIF	51
• 0.8222,0.8319,0.8450,0.8506,0.8770,0.8945,0.9116,0.9297,0.9474,	TDIF	52
• 0.9639,0.9775,0.9942,1.0180,1.0209,1.0335,1.0455,1.0605,1.0753,	TDIF	53
• 1.0905,1.1059,1.1222,1.1390,1.1540,1.1708,1.1854,1.2007,1.2138,	TDIF	54
• 1.2250,1.2343,1.2418,1.2479,1.2526,1.2565,1.2596,1.2634,1.2689,	TDIF	55

• 1.2762	1.2858	1.2968	1.3094	1.3226	1.3364	1.3507	1.3652	1.3801	TDIF 56
• 1.3947	1.4094	1.4230	1.4363	1.4488	1.4601	1.4691	1.4778	1.4871	TDIF 57
• 1.4983	1.5139	1.5320	1.5495	1.5638	1.5758	1.5864	1.5978	1.6105	TDIF 58
• 1.6223	1.6331	1.6422	1.6499	1.6563	1.6620	1.6672	1.6725	1.6783	TDIF 59
• 1.6850	1.6930	1.7023	1.7130	1.7249	1.7380	1.7521	1.7665	1.7813	TDIF 60
• 1.7950	1.8103	1.8239	1.8365						TDIF 61
									TDIF 62
DATA BIT1/									TDIF 63
• 1.8497	1.8630	1.8758	1.8891	1.9015	1.9164	1.9314	1.9468		TDIF 64
• 1.9621	1.9781	1.9947	2.0118	2.0285	2.0446	2.0598	2.0724		TDIF 65
• 2.0820	2.0883	2.0938	2.0994	2.1046	2.1104	2.1165	2.1237		TDIF 66
• 2.1320	2.1425	2.1553	2.1696	2.1843	2.1997	2.2169	2.2346		TDIF 67
• 2.2515	2.2685	2.2864	2.3029	2.3155	2.3235	2.3399	2.3514		TDIF 68
• 2.3632	2.3746	2.3872	2.4026	2.4192	2.4352	2.4542	2.4731		TDIF 69
• 2.4915	2.5091	2.5263	2.5423	2.5567	2.5695	2.5806	2.5887		TDIF 70
• 2.5949	2.6005	2.6084	2.6199	2.6326	2.6462	2.6616	2.6789		TDIF 71
• 2.6973	2.7182	2.7409	2.7639	2.7847	2.8043	2.8253	2.8467		TDIF 72
• 2.8581	2.8893	2.9103	2.9309	2.9509	2.9710	2.9914	3.0128		TDIF 73
• 3.0346	3.0565	3.0782	3.1004	3.1227	3.1446	3.1665	3.1876		TDIF 74
• 3.2077	3.2261	3.2412	3.2526	3.2632	3.2744	3.2859	3.2981		TDIF 75
• 3.3103	3.3253	3.3417	3.3604	3.3810	3.4038	3.4268	3.4487		TDIF 76
• 3.4713	3.4954	3.5104	3.5420	3.5641	3.5850	3.6051	3.6252		TDIF 77
• 3.6457	3.6667	3.6883	3.7111	3.7352	3.7608	3.7886	3.8162		TDIF 78
• 3.8411	3.8643	3.8870	3.9097	3.9325	3.9541	3.9744	3.9928		TDIF 79
• 4.0097	4.0256	4.0414	4.0574	4.0740	4.0919	4.1108	4.1316		TDIF 80
• 4.1554	4.1800	4.2049	4.2300	4.2558	4.2825	4.3095	4.3349		TDIF 81
• 4.3592	4.3817	4.4059	4.4291	4.4536	4.4781	4.5021	4.5259		TDIF 82
DATA CIT1/									TDIF 83
• 4.5497	4.5744	4.6015	4.6295	4.6579	4.6864	4.7150	4.7428		TDIF 84
• 4.7693	4.7929	4.8144	4.8333	4.8505	4.8669	4.8834	4.9005		TDIF 85
• 4.9195	4.9398	4.9617	4.9858	5.0110	5.0372	5.0651	5.0932		TDIF 86
• 5.1211	5.1487	5.1765	5.2031	5.2291	5.2524	5.2762	5.2997		TDIF 87
• 5.3223	5.3461	5.3701	5.3951	5.4210	5.4476	5.4745	5.5015		TDIF 88
• 5.5290	5.5568	5.5855	5.6139	5.6404	5.6644	5.6855	5.7065		TDIF 89
• 5.7252	5.7422	5.7583	5.7742	5.7909	5.8089	5.8280	5.8471		TDIF 90
• 5.8673	5.8891	5.9128	5.9384	5.9652	5.9923	6.0200	6.0477		TDIF 91
• 6.0742	6.0996	6.1239	6.1487	6.1743	6.2001	6.2260	6.2523		TDIF 92
• 6.2790	6.3064	6.3324	6.3567	6.3806	6.4119	6.4421	6.4684		TDIF 93
• 6.4923	6.5175	6.5422	6.5673	6.5880	6.6052	6.6243	6.6436		TDIF 94
• 6.6633	6.6941	6.7061	6.7295	6.7539	6.7789	6.8047	6.8311		TDIF 95
• 6.8574	6.8937	6.9108	6.9392	6.9673	6.9940	7.0212	7.0464		TDIF 96
• 7.0710	7.0955	7.1207	7.1467	7.1744	7.2042	7.2350	7.2659		TDIF 97
• 7.2963	7.3270	7.3558	7.3903	7.4202	7.4481	7.4751	7.5015		TDIF 98
• 7.5261	7.5486	7.5687	7.5879	7.6074	7.6269	7.6470	7.6688		TDIF 99
• 7.6924	7.7191	7.7479	7.7758	7.8051	7.8337	7.8632	7.8924		TDIF 100
• 7.9210	7.9498	7.9790	8.0089	8.0381	8.0668	8.0953	8.1237		TDIF 101
• 8.1527	8.1828	8.2143	8.2464	8.2786	8.3104	8.3421	8.3738		TDIF 102
DATA CIT1/									TDIF 103
• 8.4050	8.4350	8.4633	8.4893	8.5136	8.5365	8.5583	8.5789		TDIF 104
• 8.5982	8.6166	8.6331	8.6547	8.6757	8.6983	8.7229	8.7493		TDIF 105
• 8.7779	8.8075	8.8379	8.8655	8.8949	8.9285	8.9575	8.9854		TDIF 106
• 9.0122	9.0380	9.0635	9.0895	9.1151	9.1403	9.1653	9.1935		TDIF 107
• 9.2234	9.2546	9.2847	9.3201	9.3517	9.3840	9.4172	9.4478		TDIF 108
• 9.4763	9.5032	9.5270	9.5499	9.5743	9.5992	9.6233	9.6465		TDIF 109
• 9.6704	9.6975	9.7223	9.7560	9.7873	9.8211	9.8554	9.8898		TDIF 110
• 9.9294	9.9648	9.9982	10.0281	10.0556	10.0834	10.1137	10.1449		TDIF 111
• 10.1763	10.2114	10.2451	10.2784						

C DAYS FROM JAN 1, 1966 FOR UTC CLOCK CORRECTIONS

DATA UTCT/

.-3586.20833333333300,-3565.20833333333300,-3445.20833333333300,
 .-3419.20833333333300,-3390.20833333333300,-3249.20833333333300,
 .-3234.20833333333300,-3264.20833333333300,-3215.20833333333300,
 .-3165.20833333333300,-3131.20833333333300,-3117.20833333333300,
 .-3103.20833333333300,-3069.20833333333300,-3061.20833333333300,
 .-2999.20833333333300,-2977.20833333333300,-2942.20833333333300,
 .-2907.20833333333300,-2985.20833333333300,-2872.20833333333300,
 .-2823.20833333333300,-2760.20833333333300,-2739.20833333333300,
 .-2725.21633333333300,-2627.20833333333300,-2592.20833333333300,
 .-2564.20833333333300,-2529.20833333333300,-2501.20833333333300,
 .-2340.20833333333300,-2319.20833333333300,-2284.20833333333300,
 .-2249.20833333333300,-2235.20833333333300,-2207.20833333333300,
 .-1826.00,-1614.00,-1460.789311750000,-792.00,-730.9436619718300,
 .-640.00,-487.00,-457.00,-365.00,-305.00,-184.00,-122.00,0.000,
 .761.000,2191.000,2373.000,9999.000/

TDIF 112
 TDIF 113
 TDIF 114
 TDIF 115
 TDIF 116
 TDIF 117
 TDIF 118
 TDIF 119
 TDIF 120
 TDIF 121
 TDIF 122
 TDIF 123
 TDIF 124
 TDIF 125
 TDIF 126
 TDIF 127
 TDIF 128

C A.1 - UTC AFTER EACH UTC CLOCK CORRECTION

DATA UTCDF/

.-7.61676E-1,-7.59279E-1,-6.61401E-1,-6.32826E-1,-6.04843E-1,
 .-5.67623E-1,-5.42543E-1,-4.74538E-1,-4.19671E-1,-3.67990E-1,
 .-3.25983E-1,-2.98414E-1,-2.70676E-1,-2.43070E-1,-2.06172E-1,
 .-1.26720E-1,-8.76561E-2,-3.49597E-2, 1.84152E-2, 5.68497E-2,
 . 8.83297E-2, 1.50575E-1, 2.24081E-1, 2.63269E-1, 2.96013E-1,
 . 4.02225E-1, 4.51382E-1, 4.94829E-1, 5.46547E-1, 5.92719E-1,
 . 7.59409E-1, 7.97601E-1, 8.48046E-1, 8.99091E-1, 9.31595E-1,
 . 9.76413E-1, 1.46119E-0, 1.69407E-0, 1.89265E-0, 2.77230E-0,
 . 2.80069E-0, 3.02863E-0, 3.31660E-0, 3.35650E-0, 3.57559E-0,
 . 3.75200E-0, 4.01010E-0, 4.19045E-0, 4.35010E-0, 6.21980E-0,
 . 10.0343917E-0,11.0343917E-0/

TDIF 129
 TDIF 130
 TDIF 131
 TDIF 132
 TDIF 133
 TDIF 134
 TDIF 135
 TDIF 136
 TDIF 137
 TDIF 138
 TDIF 139
 TDIF 140
 TDIF 141

DATA UTCRT/

C A.1 - UTC RATE OF SEPARATION AFTER EACH UTC CLOCK CORRECTION

.1.06551E-3,6.48989E-4,3.17592E-4,2.85086E-4,4.10020E-4,3.62794E-4,
 .6.65787E-4,7.11582E-4,6.46546E-4,6.28629E-4,5.40983E-4,5.52709E-4,
 .5.43303E-4,6.03510E-4,9.43680E-4,9.07322E-4,9.34465E-4,9.53572E-4,
 .8.77939E-4,8.19999E-4,9.62151E-4,8.49298E-4,9.13707E-4,9.10296E-4,
 .8.79719E-4,8.33035E-4,9.37410E-4,9.06216E-4,9.34700E-4,9.11123E-4,
 .8.66292E-4,8.69835E-4,8.97014E-4,8.93154E-4,8.86354E-4,1.25857E-3,
 .1.28717E-3,1.29643E-3,1.12091E-3,1.12000E-3,1.29750E-3,1.29392E-3,
 .1.29664E-3,1.29434E-3,1.29525E-3,1.29590E-3,1.29597E-3,1.29631E-3,
 . 2.58863E-3,2.59200E-3,0.0.0.0/

TDIF 142
 TDIF 143
 TDIF 144
 TDIF 145
 TDIF 146
 TDIF 147
 TDIF 148
 TDIF 149
 TDIF 150
 TDIF 151

DATA NUTC/52/

IF(NOT1ST) GO TO 5
 NOT1ST=.TRUE.
 REFTIM=YMDAY(660101,0,0,00)
 LIMIT=LLIMIT-1
 5 TDIF=0.
 IF(BASE.EQ.IN) RETURN
 IF(BASE.GT.4.OP.IN.GT.4) RETURN
 IF(BASE.LE.3.OP.IN.LE.0) RETURN
 DAYS=DAYJ-REFTIM

TDIF 152
 TDIF 153
 TDIF 154
 TDIF 155
 TDIF 156
 TDIF 157
 TDIF 158
 TDIF 159
 TDIF 160
 TDIF 161

C TIME CORRECTIONS IN ORDER

C UT2-UT1 TO JT1-A.1 TO A.1-UTC

I1=ID(IN)
 I2=ID(BASE)
 J1=MINC(I1,I2)

TDIF 162
 TDIF 163
 TDIF 164
 TDIF 165
 TDIF 166
 TDIF 167

J2=MAXC(I1,I2)	TDIF 168
C START CORRECTION CALCULATION WITH SMALLER TIME BASE	TDIF 169
GO TO (10,20,30,50),J1	TDIF 170
C COMPUTE UT2-UT1	TDIF 171
10 TWCPID=TWCP1*(DAYS-.201D0)/365.2422D0	TDIF 172
FORPID=2.*TWCPID	TDIF 173
TDIF=TDIF+.022*SIN(TWCPID)-.012*COS(TWCPID)	TDIF 174
-.006*SIN(FORPID)+.007*COS(FORPID)	TDIF 175
C TEST FOR OUTPUT TIME SYSTEM	TDIF 176
IF(J2.EQ.2) GO TO 50	TDIF 177
C COMPUTE UT1-A.1	TDIF 178
20 DT=(DAYS+3037.0D0)*1.0D-1	TDIF 179
I=MINO(LIMIT,MAX1(1.,DT))	TDIF 180
DT=DT-FLDAT(I)	TDIF 181
TDIF=TDIF-AIT1(I)-DT*(AIT1(I+1)-AIT1(I))	TDIF 182
IF(J2.EQ.3) GO TO 50	TDIF 183
C COMPUTE A.1-UTC	TDIF 184
30 DO 40 I=1,NUTC	TDIF 185
IF(DAYS.LT.UTCT(I+1)) GO TO 45	TDIF 186
40 CONTINUE	TDIF 187
I=NUTC	TDIF 188
45 TDIF=TDIF+UTCDF(I)+UTCRT(I)*(DAYS-UTCT(I))	TDIF 189
C SET DIRECTION OF CORRECTION	TDIF 190
50 IF(I1.EQ.J1)TDIF=-TDIF	TDIF 191
RETURN	TDIF 192
END	TDIF 193

TIDAL

DESCRIPTION

TIDAL evaluates the acceleration on a satellite caused by the Earth tidal bulge resulting from Lunar and Solar gravitational effects on the Earth.

NAME TICAL

PURPOSE TO COMPUTE ACCELERATION DUE TO SOLID EARTH TIDAL BULGES CAUSED BY LUNAR AND SOLAR GRAVITATIONAL EFFECTS ON EARTH. USES ONLY K2 IN SPHERICAL HARMONIC EXPANSION

CALLING SEQUENCE CALL TICAL(DX)

SYMBOL	TYPE	DESCRIPTION
DX (3)	DP	INPUT & OUTPUT - SATELLITE ACCELERATION VECTOR

SUBROUTINES USED NONE

COMMON BLOCKS CEPHEM INITBK INTBLK MOONGR XYZ

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES 'GEODYN SYSTEMS DESCRIPTION'
VOLUME 1 - GEODYN DOCUMENTATION

```

SUBROUTINE TICAL(DX)
  IMPLICIT REAL*8 (A-H,C-Z)
  LOGICAL NOT1ST
  DOUBLE PRECISION K2,K3,LAMBDA
  DIMENSION CONST(2),ACCEL(3),DX(3)
  COMMON/CEPHEM/UVBODY(24),EQ(644)
  COMMON/INITBK/IG1(56),NOT1ST
  COMMON/INTBLK/THCOTS(3),GM,AE,AESQ(4),RATIO(53),K2,K3,LAMBDA
  *   TOREF
  COMMON/MOONGR/DPXUV(6),RHOM(3,6),RHOSQ(12)
  COMMON/XYZ/XYZ(6),R,RSQ,ISAT,IFORCE(2)
  IF(NOT1ST) GO TO 10
  DO 5 I=1,2
  *   CONST(I)=K2*AE*RATIO(I)*0.5D0
  NOT1ST=.TRUE.
  *   TIDA 43
10  K=C
  DO 100 I1=1,8,4
  *   K=K+1
  *   DP=DPXUV(K)/R
  *   GP2=1.0D0-15.0D0*DP**2
  DO 20 I=1,3
  *   J=I1+I-1
  *   TIDA 49
20  ACCEL(I)=DP*6.0D0*UVBODY(J)+DP**3*XYZ(I)/R
  *   J1=I1+J
  *   TIDA 52
  *   RATIO4=(AE/R)**4/UVBODY(J1)**3
  DO 100 I=1,3
  *   TIDA 54
100 DX(I)=DX(I)+ACCEL(I)*CONST(K)*RATIO4
  *   TIDA 55
  RETURN
  *   TIDA 56
  END
  *   TIDA 57

```

NAME TRUEP
ENTRY POINT PURPOSE
TRUEP INITIALIZATION
TRUEP TO ROTATE TRACKING STATION COORDINATES TO ACCOUNT FOR POLAR WANDERING

CALLING SEQUENCE CALL TRUEP1(STAXYZ)

SYMBOL	TYPE	DESCRIPTION
STAXYZ	DP	INPUT - TRACKING STATION CARTESIAN COORDINATES
(3,1)		

CALLING SEQUENCE CALL TRUEP(DAY,ISTA)

SYMBOL	TYPE	DESCRIPTION
DAY	DP	INPUT - OBSERVATION TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR FOR THE ARC
ISTA	I	INPUT - STATION INDEX

SUBROUTINES USED POLE

COMMON BLOCKS TRUPOL

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES 'GEODYN SYSTEMS DESCRIPTION'
VOLUME 1 - GEODYN DOCUMENTATION

SUBROUTINE TRUEP1(STAXYZ)	TRUE	40
IMPLICIT REAL*8 (A-H, O-Z)	TRUE	41
DOUBLE PRECISION MATRX(3,3)	TRUE	42
DIMENSION STAXYZ(3,1)	TRUE	43
COMMON/TRUPOL/TRUE(3)	TRUE	44
DATA MATRX/1.00,3*0.00,1.00,3*0.00,1.00/	TRUE	45
RETURN	TRUE	46
ENTRY TRUEP(DAY,ISTA)	TRUE	47
C COMPUTE POLE COORDINATES	TRUE	48
CALL POLE (X,Y,DAY)	TRUE	49
C ROTATION MATRIX	TRUE	50
MATRX(1,3) = -X	TRUE	51
MATRX(2,1) = X*Y	TRUE	52
MATRX(2,3) = Y	TRUE	53
MATRX(3,1) = X	TRUE	54
MATRX(3,2) = -Y	TRUE	55

C ROTATE COORDINATES

DO 100 I=1,3

TRUE(I) = 0.00

DO 100 J=1,3

TRUE(I)=TRUE(I)+MATR X(I,J)*STAXY7(J,ISTA)

100 CONTINUE

RETURN

END

TRUE 56

TRUE 57

TRUE 58

TRUE 59

TRUE 60

TRUE 61

TRUE 62

TRUE 63

TWOSTA

DESCRIPTION

Subroutine TWOSTA computes the residuals and partial derivatives for observations involving two tracking stations. Subroutine UPDOWN is used by TWOSTA to compute up- and down-link signal transit times necessary in the computation of precise transmission and transponder relay times. The partial derivatives and calculated measurements are sums and/or differences of equivalent range and range rate quantities computed by subroutine PREDCT.

The observation types by program index are:

- 27) VLBI time delay, τ_g
- 28) VLBI fringe rate, v_F
- 29) Two-way average range rate, $\dot{\bar{\rho}}_2$, and
- 30) Three-way average range rate, $\dot{\bar{\rho}}_3$.

NAME TWOSTA
 ENTRY POINT PURPOSE
 TWOST1 INITIAL IZATION
 TWOSTA TO COMPUTE MEASUREMENTS & PARTIALS FOR VLBI & AVERAGE RANGE RATE DATA

CALLING SEQUENCE CALL TWOST1(PMPX0,NPARM)

SYMBOL	TYPE	DESCRIPTION
PMPX0 (NPARM,1)	DP	OUTPUT - MEASUREMENT PARTIALS WITH RESPECT TO EPOCH PARAMETERS
NPARM	I	INPUT - MAXIMUM NUMBER OF PARAMETERS PER MEASUREMENT

CALLING SEQUENCE CALL TWOSTA(ISTA, DAY2, RESID, DATASW)

SYMBOL	TYPE	DESCRIPTION
ISTA	I	INPUT - STATION INDEX
DAY2	DP	INPUT - MEASUREMENT TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR
RESID	DP	INPUT - MEASUREMENT RESIDUAL (O-C)
DATASW	L	INPUT - .TRUE. WHEN POSITION OF SATELLITE WANTED .FALSE. WHEN MEASUREMENT PARTIALS WANTED

SUBROUTINES USED GRHRAN UPDOWN ORBIT OBSDOT PROCES
 PREOCT

COMMON BLOCKS CONSTS CORBI CUVECT PREBLK

INPUT FILES NONE

OUTPUT FILES NONE

GEOCYN SYSTEMS DESCRIPTION
 VOLUME 1 - GEOCYN DOCUMENTATION

SUBROUTINE TWOST1(PMPX0,NPARM)	TWOS	40
IMPLICIT REAL*8 (A-H,C-Z)	TWOS	50
LOGICAL*1 VHFCHN,PREPR7,SAVEPR	TWOS	51
LOGICAL DATASW,LOGIC	TWOS	52
INTEGER*2 MTYPE,NMCAS,PRETYO,ISNO,ISAT	TWOS	53
INTEGER RECNO	TWOS	54
DIMENSION PMPX0(NPARM,1),RENDX(2)	TWOS	55

REAL D1D,D1U,D2D,D2U,RFNDX	TWOS	56
COMMON/CONSTS/GPI,DTWOP1,DPAD,CRSEC	TWOS	57
COMMON/CRB1/T,W,U,THETG,PERHT(2),APHT(2),PRD(2)	TWOS	58
COMMON/CUVECT/UHAT(3,2),XY7(3,2),RXYZ(3,2),RFNV(3,2),R(2),	TWOS	59
RSO(2),XYSO(2)	TWOS	60
COMMON/PREBLK/DAYSAT,CHS,DAY1,SIG,D2D,D2U,D1D,D1U,ISN,MTYPE,	TWOS	61
AMEAS,ISAT,PRETYP,ISN2,VHFCNN,PREPRO,RECND	TWOS	62
DATA C/2.99792503/	TWOS	63
EQUIVALENCE (DSTA1,DAY1),(DSTA2,D2D),(FREQ,D1D)	TWOS	64
EQUIVALENCE (RFNDX(1),REFRAC)	TWOS	65
C INITIALIZE	TWOS	66
NP3=NPARM-3	TWOS	67
NP6=NPARM-6	TWOS	68
RETURN	TWOS	69
ENTRY TWOSTA(ISTA,DAY2,PESID,DATASW)	TWOS	70
IF(.NOT.DATASW).GO TO 100	TWOS	71
C OBTAIN EPHEMERIS	TWOS	72
CALL PREDCT(ISTA,DAY2,RESID,PESID2,DATASW)	TWOS	73
RETURN	TWOS	74
C SET SWITCHES	TWOS	75
100 IF(.NOT.PREPRO) PRETYP=0	TWOS	76
IPRE1=PRETYP/10	TWOS	77
IPRE2=PRETYP-IPRE1*10	TWOS	78
SAVEPP=PREPRO	TWOS	79
ISTA2=ISN2	TWOS	80
KTYPE=MTYPE-26	TWOS	81
F=1.000	TWOS	82
IF(PREPRO) GO TO (205,200,300,300),KTYPE	TWOS	83
GO TO (250,260,400,400),KTYPE	TWOS	84
C VLBI DATA	TWOS	85
200 F=DSTA2	TWOS	86
205 IF(IPRE1.LE.0) GO TO 240	TWOS	87
DSTA1=DAY2	TWOS	88
C COMPUTE VLBI TRANSIT TIMES	TWOS	89
CALL UPDOWN(DAY2,DSTA1,DSTA2,1.0DC,ISTA,ISTA2,ISAT,.FALSE.,RCOR,	TWOS	90
RFNDX)	TWOS	91
240 PREPRO=IPRE2.GT.0	TWOS	92
IF(.NOT.PREPRO) GO TO 250	TWOS	93
C APPLY REFRACTION CORRECTION TO OBSERVATION	TWOS	94
PRETYP=IPRE2	TWOS	95
OBSAVE=OBS	TWOS	96
OBS=0.000	TWOS	97
REFRAC=FREQ	TWOS	98
FREQ=RFNDX(2)	TWOS	99
ISAVE=MTYPE	TWOS	100
MTYPE=KTYPE+1	TWOS	101
CALL CRBIT(DAY2)	TWOS	102
THETG=GRHRAN(DSTA2,ISTA2)	TWOS	103
CALL PROCES(ISTA2,DSTA2,THETG)	TWOS	104
OBS=-OBS	TWOS	105
FREQ=RFNDX(1)	TWOS	106
THETG=GRHRAN(DSTA1,ISTA)	TWOS	107
PREPRO=SAVEPP	TWOS	108
CALL PROCES(ISTA,DSTA1,THETG)	TWOS	109
OBS=OBSAVE+F*OBS/C	TWOS	110
MTYPE=ISAVE	TWOS	111


```




250 FREQ=F
C OBTAIN ORBIT
260 CALL ORBIT(DAY2)
    JTYPE=KTYPE+1
    F=FREQ
    IF(KTYPE.EQ.1) F=1.000
C CALCULATE RIGHT ASCENSION OF GREENWICH AND QUANTITIES NECESSARY TO
C COMPUTE PARTIALS FOR SECOND STATION
    THETG=GRHRAN(DSTA2,ISTA2)
    T=ATAN(RENV(3,ISAT)/COSPT(1.000-RENV(3,ISAT)**2))/DRAD
C COMPUTE PARTIALS FOR SECOND STATION
    CALL PREDCT(ISTA2,DAY2,RESID,RESID2,DATASW)
    PMPX0(NPARM-9,1)=ORSDDT(JTYPE,ISTA2,P)*F/C
    ORSC=-RESID
    DO 270 I=1,NPARM
270 PMPX0(I,2)=-PMPX0(I,1)
C CALCULATE RIGHT ASCENSION OF GREENWICH AND QUANTITIES NECESSARY TO
C COMPUTE PARTIALS FOR FIRST STATION
    THETG=GRHRAN(DSTA1,ISTA1)
C COMPUTE PARTIALS FOR FIRST STATION
    CALL PREDCT(ISTA,DAY2,RESID,RESID2,DATASW)
    PMPX0(NPARM-6,1)=ORSDDT(JTYPE,ISTA,P)*F/C
    ORSC=ORSC+RESID
    DO 280 I=1,NP6
280 PMPX0(I,1)=PMPX0(I,1)+PMPX0(I,2)
    DO 290 I=1,3
290 PMPX0(NP6+I+3,1)=PMPX0(NP6+I,2)
    PMPX0(NPARM-7,1)=1.000
    RESID=OBS-ORSC
    RETURN
C AVERAGE RANGE RATE DATA
300 RCR=0.000
    IF(IPRE1.LE.0) GO TO 400
    REFRAC=0.000
    IF(IPRE2.LE.0) GO TO 350
C SET REFRACTION INDICES FOR TRANSMITTER & RECEIVER
    REFRAC=FREQ
    IF(RFNDX(1).LE.0.) RFNDX(1)=328.5
    IF(RFNDX(2).LE.0.) RFNDX(2)=328.5
    RFNDX(1)=RFNDX(1)*0.84323360-2
    RFNDX(2)=RFNDX(2)*0.84323360-2
350 DAYR=DAY1
    PREPRO=IPRE2.GT.0
    LOGIC=PREPRO
C COMPUTE TRANSIT TIMES FOR START OF COUNTING INTERVAL
    CALL UPDOWN(DAY1,DAYR,CAYT,-1.000,ISTA,ISTA2,ISAT,LOGIC,RCR,
    RFNDX)
    DID=DAYR-DAY1
    DIU=DAY1-DAYT
    DAYR=DAY2
    CAY2=DAYR-DID
    RCR=-RCR
C COMPUTE TRANSIT TIMES FOR END OF COUNTING INTERVAL
    CALL UPDOWN(DAY2,DAYR,CAYT,-1.000,ISTA,ISTA2,ISAT,LOGIC,RCR,
    RFNDX)
    D2D=DAYR-DAY2
  
```



TWOS 112
 TWOS 113
 TWOS 114
 TWOS 115
 TWOS 116
 TWOS 117
 TWOS 118
 TWOS 119
 TWOS 120
 TWOS 121
 TWOS 122
 TWOS 123
 TWOS 124
 TWOS 125
 TWOS 126
 TWOS 127
 TWOS 128
 TWOS 129
 TWOS 130
 TWOS 131
 TWOS 132
 TWOS 133
 TWOS 134
 TWOS 135
 TWOS 136
 TWOS 137
 TWOS 138
 TWOS 139
 TWOS 140
 TWOS 141
 TWOS 142
 TWOS 143
 TWOS 144
 TWOS 145
 TWOS 146
 TWOS 147
 TWOS 148
 TWOS 149
 TWOS 150
 TWOS 151
 TWOS 152
 TWOS 153
 TWOS 154
 TWOS 155
 TWOS 156
 TWOS 157
 TWOS 158
 TWOS 159
 TWOS 160
 TWOS 161
 TWOS 162
 TWOS 163
 TWOS 164
 TWOS 165
 TWOS 166
 TWOS 167

```

    O2U=DAY2-DAYT
  400 DAYR=DAY1+D1D
    DAYT=DAY1-D1U
    CALL ORBIT(DAY1)
    THETG=GRHRAN(DAYT, ISTA2)
    T=DATAN(PENV(3, ISAT)/CSQRT(1.000-PENV(3, ISAT)**2))/DRAD
    OBSO=-R(ISAT)
  C COMPUTE UPLINK RANGE & PARTIALS AT START
    CALL PREDCT(ISTA2, DAY1, RESID, RESID2, DATASW)
    PMPX0(NPARM-8, 1)=OBSOCT(2, ISTA2, P)
    DO 410 I=1, NP6
  410 PMPX0(I, 2)=-PMPX0(I, 1)
    DO 420 I=1, 3
    PMPX0(I+NP6, 2)=0.000
  420 PMPX0(I+NP3, 2)=-PMPX0(I+NP6, 1)
    THETG=GRHRAN(DAYR, ISTA)
    OBSO=OBSO-R(ISAT)
  C COMPUTE DOWNLINK RANGE & PARTIALS AT START
    CALL PREDCT(ISTA, DAY1, RESID, RESID2, DATASW)
    PMPX0(NPARM-6, 1)=OBSOCT(2, ISTA, P)
    DO 430 I=1, NP3
  430 PMPX0(I, 2)=PMPX0(I, 2)-PMPX0(I, 1)
    CALL ORBIT(DAY2)
    DAYT=DAY2-O2U
    DAYR=DAY2+O2D
    THETG=GRHRAN(DAYT, ISTA2)
    OBSO=OBSO+R(ISAT)
  C COMPUTE UPLINK RANGE & PARTIALS AT END
    CALL PREDCT(ISTA2, DAY2, RESID, RESID2, DATASW)
    PMPX0(NPARM-8, 1)=OBSOCT(2, ISTA2, P)
    DO 440 I=1, NP6
  440 PMPX0(I, 2)=PMPX0(I, 2)+PMPX0(I, 1)
    DO 450 I=1, 3
  450 PMPX0(I+NP3, 2)=PMPX0(I+NP3, 2)+PMPX0(I+NP6, 1)
    THETG=GRHRAN(DAYR, ISTA)
    OBSO=OBSO+R(ISAT)
  C COMPUTE DOWNLINK RANGE & PARTIALS AT END
    CALL PREDCT(ISTA, DAY2, RESID, RESID2, DATASW)
    PMPX0(NPARM-6, 1)=OBSOCT(2, ISTA, P)
    DO 460 I=1, NP3
  460 PMPX0(I, 2)=PMPX0(I, 2)+PMPX0(I, 1)
    DELTAT=(DAY2+O2D-DAY1-D1D)*17.2804
    DTINV=1.000/DELTAT
    IF(PREPRO) OBS=OBS+RCCR*DTINV
    PREPRO=.FALSE.
  C CALCULATE RESIDUAL
    RESID=OBS-OBSO*DTINV
    DO 470 I=1, NPARM
  470 PMPX0(I, 1)=PMPX0(I, 2)*DTINV
    PMPX0(NPARM-7, 1)=1.000
    IF(MTYPE.EQ.30) RETURN
    PMPX0(NP6+1, 1)=PMPX0(NP6+1, 1)+PMPX0(NP3+1, 1)
    PMPX0(NP6+2, 1)=PMPX0(NP6+2, 1)+PMPX0(NP3+2, 1)
    PMPX0(NP6+3, 1)=PMPX0(NP6+3, 1)+PMPX0(NP3+3, 1)
    PMPX0(NPARM-6, 1)=PMPX0(NPARM-6, 1)+PMPX0(NPARM-8, 1)
    PMPX0(NPARM-8, 1)=0.000
    PMPX0(NP3+1, 1)=0.000
    PMPX0(NP3+2, 1)=0.000
    PMPX0(NP3+3, 1)=0.000
    RETURN
  END
  
```

TWOS 168
 TWOS 169
 TWOS 170
 TWOS 171
 TWOS 172
 TWOS 173
 TWOS 174
 TWOS 175
 TWOS 176
 TWOS 177
 TWOS 178
 TWOS 179
 TWOS 180
 TWOS 181
 TWOS 182
 TWOS 183
 TWOS 184
 TWOS 185
 TWOS 186
 TWOS 187
 TWOS 188
 TWOS 189
 TWOS 190
 TWOS 191
 TWOS 192
 TWOS 193
 TWOS 194
 TWOS 195
 TWOS 196
 TWOS 197
 TWOS 198
 TWOS 199
 TWOS 200
 TWOS 201
 TWOS 202
 TWOS 203
 TWOS 204
 TWOS 205
 TWOS 206
 TWOS 207
 TWOS 208
 TWOS 209
 TWOS 210
 TWOS 211
 TWOS 212
 TWOS 213
 TWOS 214
 TWOS 215
 TWOS 216
 TWOS 217
 TWOS 218
 TWOS 219
 TWOS 220
 TWOS 221
 TWOS 222
 TWOS 223
 TWOS 224
 TWOS 225
 TWOS 226
 TWOS 227
 TWOS 228





REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

NAME TYPORB

PURPOSE TO PRINT ARC SUMMARY PAGE

CALLING SEQUENCE CALL TYPORB(DAYO,MINDEX,ARCNO,OUTER,RMSPOS)

SYMBOL	TYPE	DESCRIPTION
DAYO	DP	INPUT - TIME OF FINAL SET OF PARTIAL DERIVATIVES FOR DRAG
MINDEX	I	INPUT - TOTAL NUMBER OF MEASUREMENT PLUS 1
ARCNO	I	INPUT - ARC NUMBER
OUTER	I	INPUT - OUTER ITERATION
RMSPOS (2,1)	DP	INPUT - ARRAY CONTAINING RMS OF POSITION AND VELOCITY FOR ADJUSTED ELEMENTS

SUBROUTINES USED DOTPRD ELEM DATES APPER

COMMON BLOCKS APARAM ALPMRC CELEM CGEOS CORB1
 CSTINF CTIME GEODYN INITRK INTBLK
 PRIORI TPEBLK XYZOUT CONSTS

INPUT FILES NONE

OUTPUT FILES NONE

'GEODYN SYSTEMS DESCRIPTION'
VOLUME 1 - GEODYN DOCUMENTATION

SUBROUTINE TYPORB(DAYO,MINDEX,ARCNO,OUTER,RMSPOS)		
IMPLICIT REAL*8 (A-H,C-Z)		TYPO 37
LOGICAL DRAGSW,ORBSW,HYPER		TYPO 38
INTEGER ADDR,OUTP,DATP,ARCNO,OUTER		TYPO 39
REAL RMSTOT,TYPRMS,BSUM		TYPO 40
DOUBLE PRECISION LHAT,MSAT		TYPO 41
DIMENSION LHAT(2),OL(2),OT(2),RMSPOS(2,1)		TYPO 42
COMMON/APARAM/INPAR(4),NSAT,NGPARC(5)		TYPO 43
COMMON/ALPMRC/ITNMS(14),ATYPE(47),HYPER		TYPO 44
COMMON/CELEM/ELEMST(6,2),ORBLA(6,2),EC,XNU,RMSTDT		TYPO 45
COMMON/CGEOS/ISATIO(2),IPREFR(453)		TYPO 46
COMMON/CONSTS/PI,TWOP1,DRAG,PSEC		TYPO 47
COMMON/CORB1/RANCOT(2),PERDOT(2),PERPHT(2),APHT(2),PRD(2)		TYPO 48
COMMON/CSTINF/MEASNO(35),TYPRMS(30),NOTYPE(2,30),BSUM(155),		TYPO 49
NWTOT,LPASE		TYPO 50
COMMON/CTIME/DATAEP(2),DSTART,DAYSTP,DAYINT(7),IYREG		TYPO 51
COMMON/GEODYN/DATE(5)		TYPO 52
COMMON/INITEK/IEPYND,IEPHM,EPSEFC,IDUM1(32),DRATSW,MISLOG(20)		TYPO 53
COMMON/INTBLK/THDOT(3),GM,AE,AESQ(47),ASAT(2),MSAT(2),VARSTP(3),		TYPO 54
		TYPO 55

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

```


    ADDR(14)
COMMON/PRIORI/ELEMIN(48),TITLE(30),DRAG(12),CD(2),CDD(2),EMISS(2)
COMMON/TPERBLK/INTP,OUTP,OUTP(10)
COMMON/XYZOUT/ELEMS(6,2),DRGPAR(6,2)
C COMPUTE ORBITAL CHARACTERISTICS FROM KEPLER ELEMENTS FOR EACH
C-SATELLITE
DO 100 I=1,NSAT
  DRAGSW=ADDR(I).GT.0
  CALL ELEM(ELEMST(1,I),ORBELA(1,I),1.,.FALSE.,ORBELA)
  IF(.NOT.DRAGSW) GO TO 75
  RV=DOTPRD(ELEMS(1,I),ELEMS(4,I))
  VSO=DOTPRD(ELEMS(4,I),ELEMS(4,I))
  RSO=DOTPRD(ELEMS(1,I),ELEMS(1,I))
  D=DSORT(VSO-RV**2/RSO)
  DO 50 J=1,3
    50 LHAT(J)=(-RV*ELEMS(J,I)/RSO+ELEMS(J+3,I))/D
    DL(I)=DOTPRD(LHAT,DRGPAR(1,I))*CD(I)/(DAYC-DSTART)**2
    75 AEA35=(AE/DAES(ORBELA(1,I))**3.5/(1.000-ORBELA(2,I)**2))**2
    COSI=DCOS(ORBELA(3,I)*ORAD)
    RANDOT(I)=-9.9700*AEA35*COSI
    PEROOT(I)=4.9800*AEA35*(5.000*COSI**2-1.000)
    100 IF (DRAGSW) DT(I)=6.000*P(ABSORT(DABS(ORBELA(1,I))**3/GM)+
      (DOTPRD(ELEMS(1,I),DRGPAR(1,I))/(RSO*DSORT(RSO))+
      DOTPRD(ELEMS(4,I),DRGPAR(4,I))/GM)*CD(I)/(DAYC-DSTART)
      NOBS=MINDEX-1
C COMPUTE DATA START & STOP TIMES
  CALL DATES(CATAEP(1),IYMD1,IHM1,SEC1)
  CALL DATES(DAYSTR,ITMD,INH,SEC)
  DO 150 K=1,2
C WRITE TITLE & EPOCH
  WRITE(OUTP,1000) DATE,ARCNO,CUTER,TITLE
  WRITE(OUTP,4000) IEPYMD,IEPHM,EPSEC,IYMD1,IHM1,SEC1,IYMD,IHM,SEC
  IF(CRSTW) GO TO 250
  IF(NOBS.LE.0) GO TO 250
C WRITE RESIDUAL SUMMARY FOR DATA REDUCTION RUN
  WRITE(OUTP,5000)
  DO 200 I=1,NS
    200 IF(NOTYPE(2,I).GT.0) WRITE(OUTP,6000) ATYPE(I),NOTYPE(2,I),
      TYPRMS(I)
  WRITE(OUTP,7000) NOWTCB,RMSTOT,NOBS
  250 CALL APPR
C WRITE ELEMENTS & ORBITAL CHARACTERISTICS FOR EACH SATELLITE
DO 150 I=1,NSAT
  WRITE(OUTP,3000) I,(ELEMST(J,I),J=1,6),(ORBELA(J,I),J=1,5),
    ISATID(I),ASAT(I),MSAT(I),CD(I),CDD(I),EMISS(I)
  IF(.NOT.CRSTW) WRITE(OUTP,9000) (RMSPOS(J,I),J=1,2)
  DRAGSW=ADDR(I).GT.0
  PERMIL=PERHT(I)/1.6093500
  APRMIL=APHT(I)/1.6093500
  PRD(I)=TWOPI*DSORT(DABS(ORBELA(1,I))**3/GM)
  PRDMIN=PRD(I)/60.000
  WRITE(OUTP,2000) RANDOT(I),PEROOT(I),PERHT(I),PERMIL,APHT(I),
    APRMIL,PRDMIN
  150 IF(DRAGSW) WRITE(OUTP,3000) DL(I),DT(I)
  RETURN
1000 FORMAT(1H1,14X,5A8,'0A2C',14)
  
```

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

```

    * SUMMARY FOR LAST INNER ITERATION OF OUTER *
    * ITERATION',I2/3(/1X,10A8))
2000 FORMAT(1H0, 9HMODE RATE,3X,12HARG PER RATE,6X,14HPERIGEE HFIGHT
    * ,12X,13HAPDCEE HEIGHT,10X,6HPERIOD/
    * 1X,2(9H(DEG/DAY),4X),4X,4H(KM),
    * 6X,7H(MILES),8X,4H(KM),7X,7H(MILES),4X,9H(MINUTES)/
    * 1X,F9.5,F12.5,3X,2F12.4,2F13.4,F12.3)
3000 FORMAT(1H0,34X,4HORAG,7X,13HPERIOD DEC/34X,10H(M/DAY**2),4X,
    * 9H(SEC/DAY)/28X,2F14.4)
4000 FORMAT('EPOCH',I7,I5,F8.4,' BEGIN TIME',I7,I5,F8.4,
    * ' END',I7,I5,F8.4)
5000 FORMAT('MEAS. TYPE',5X,'NO. OF WTD RESIDUALS',5X,'WTD RMS')
6000 FORMAT(3X,A6,7X,I12,8X,F12.3)
7000 FORMAT('C TOTAL',8X,I12,8X,F12.3,' TOTAL NO. OF OBS =',I8)
8000 FORMAT('SAT',I2,' CURRENT ELEMENTS. UNITS: LENGTH-METERS, '
    * 'TIME-SECONDS, ANGLES-DEGREES,'/ ' X =',G22.16,' Y =',G22.16
    * ' Z =',G22.16/' XDOT=',G22.16,' YDOT=',G22.16,' ZDOT=',
    * G22.16/' A =',G22.16,' E =',G22.16,' INCL=',G22.16/
    * ' NODE=',G22.16,' PERG=',G22.16,' MEAN=',G22.16/' SATID=',I8,
    * ' AREA (M**2) =',1PD12.4,' MASS (KG) =',3PD12.2/' DRAG=',
    * 1PD12.4,' CRAG DOT=',D12.4,' SOLRAD=',D12.4)
9000 FORMAT(' RMS POSITION =',F12.3,' RMS VELOCITY =',F12.6)
    END
  
```

TYPD 112
 TYPD 113
 TYPD 114
 TYPD 115
 TYPD 116
 TYPD 117
 TYPD 118
 TYPD 119
 TYPD 120
 TYPD 121
 TYPD 122
 TYPD 123
 TYPD 124
 TYPD 125
 TYPD 126
 TYPD 127
 TYPD 128
 TYPD 129
 TYPD 130
 TYPD 131
 TYPD 132
 TYPD 133
 TYPD 134



UPDATE

DESCRIPTION

This subroutine uses the matrix partitioning techniques described in the GEODYN Systems Description, Volume I, to update the estimated values of arc adjusted parameters for corrections resulting from the adjustment of common parameters in the partitioned solution.

NAME UPDATE

PURPOSE TO UPDATE ARC ADJUSTED PARAMETERS TO COMPENSATE FOR ADJUSTMENTS TO COMMON PARAMETERS

CALLING SEQUENCE CALL UPDATE(NARCS, OUTER, NSTART, SUM1, DELTA, DDELTA, TTL, BSNOS, BSVAL)

SYMBOL	TYPE	DESCRIPTION
NARCS	I	INPUT - NUMBER OF ARCS
OUTER	I	INPUT - OUTER ITERATION NUMBER
NSTART	I	INPUT - STARTING LOCATION IN NORMAL MATRIX OF COMMON PARAMETERS
SUM1 (1)	DP	INPUT - NORMAL MATRIX
DELTA (1)	DP	INPUT & OUTPUT - CORRECTION VECTORS FOR ADJUSTED PARAMETERS
DDELTA (1)	DP	SCRATCH
TTL (1)	DP	INPUT - PARAMETER TITLE ARRAY
BSNOS (1)	DP	INPUT - LOCATIONS IN NORMAL MATRIX OF BIASES, DRAG, SOLRAD, AND GEOPOTENTIAL PARAMETERS
BSVAL (1)	DP	INPUT & OUTPUT - VALUES OF BIASES, DRAG, SOLRAD, AND GEOPOTENTIAL PARAMETERS

SUBROUTINES USED CORREL DATARD

COMMON BLOCKS APARAM CELEM CONSTS CPARAM TPEBLK
 PRIORI

INPUT FILES NONE

OUTPUT FILES OUTP - PRINTER

GEDDYN SYSTEMS DESCRIPTION
VOLUME 1 - GEDDYN DOCUMENTATION

SUBROUTINE UPDATE(NARCS, OUTER, NSTART, SUM1, DELTA, DDELTA, TTL, BSNOS, BSVAL)	UPDA	51
IMPLICIT REAL*8 (A-H, C-Z)	UPDA	52
LOGICAL EMPGR	UPDA	53
INTEGER ARCN0, OUTER, OUTP	UPDA	54
	UPDA	55



INTEGER*2 BSNOS	UPDA	56
REAL BSVL	UPDA	57
DIMENSION SUM1(1), DELTA(1), DDELTA(1), TTL(1), BSNOS(1), BSVL(1)	UPDA	58
COMMON/APARAM/ INPAR, INPAR1, NBIAS, NSTSTA, NSAT, NGPARC, NPRECI, NPARAM,	UPDA	59
NEBIAS, MAXPAR	UPDA	60
COMMON/CELEM/ELEMST(12), ORRELA(12), IDUM(5)	UPDA	61
COMMON/CONSTS/DPI, DTWCPI, DRAD, DRSEC	UPDA	62
COMMON/CPARAM/NSTA, NMAST, NSTEST, NDIM, MBIAS, NGPC1, NGPC2, NGPCOM,	UPDA	63
NCEST, CMPGPR, LIM1, LIM2, NGEN, NGENST, NTIDST, NTIDEN, INNRSW,	UPDA	64
NCONST, NCONNS	UPDA	65
COMMON/PRIORI/ELEM IN(5), CD(2,3)	UPDA	66
COMMON/TPEBLK/INTP, OUTP, ITAPES(10)	UPDA	67
INDXNO(I)=NDIM*(I-1)-(I*(I-1))/2	UPDA	68
C LOOP THROUGH 1000 FOR EACH ARC	UPDA	69
DO 1000 ARCNC=1, NARCS	UPDA	70
C READ ARC INFORMATION	UPDA	71
CALL DATARD(ARCNC, .FALSE., .FALSE., .TRUE.)	UPDA	72
II=0	UPDA	73
C COMPUTE CORRECTIONS	UPDA	74
DO 100 I=1, NPARAM	UPDA	75
DELTA(I)=0.000	UPDA	76
DO 50 L=NSTART, NDIM	UPDA	77
LI=II+L	UPDA	78
50 DELTA(I)=DELTA(I)-SUM1(LI)*DELTA(L)	UPDA	79
100 II=II+NDIM-1	UPDA	80
II=5+NSAT	UPDA	81
C CORRECT ELEMENTS	UPDA	82
DO 150 I=1, II	UPDA	83
150 ELEMST(I)=ELEMST(I)+DELTA(I)	UPDA	84
NBSGP=MBIAS+NGPARC	UPDA	85
C CORRECT BIASES, GEOPOTENTIAL COEFFICIENTS, DRAG & SOLRAD	UPDA	86
DO 200 I=1, NBSGP	UPDA	87
IF(BSNOS(I).EQ.0) GO TO 200	UPDA	88
II=II+1	UPDA	89
BSVAL(I)=BSVAL(I)+DELTA(II)	UPDA	90
200 CONTINUE	UPDA	91
II=0	UPDA	92
DO 250 L=1, NSAT	UPDA	93
DO 250 I=1, 3	UPDA	94
II=II+1	UPDA	95
IF(BSNOS(II).LE.0) GO TO 250	UPDA	96
CD(L, I)=BSVAL(II)	UPDA	97
250 CONTINUE	UPDA	98
IST=INDXNO(NSTART)	UPDA	99
LI=0	UPDA	100
C LOOP THROUGH 500 CORRECTS NORMAL MMATRIX	UPDA	101
DO 600 L=1, NPARAM	UPDA	102
MST=IST	UPDA	103
DO 400 M=NSTART, NDIM	UPDA	104
DDELTA(M)=0.000	UPDA	105
NM=IST+M	UPDA	106
DO 300 N=NSTART, M	UPDA	107
NL=L1+N	UPDA	108
DDELTA(M)=DDELTA(M)+SUM1(NM)*SUM1(NL)	UPDA	109
300 NM=N+NDIM-N	UPDA	110
IF(M.EQ.NDIM) GO TO 400	UPDA	111

```

MP1=M+1
DO 350 N=MP1,NDIM
NM=MST+N
NL=L1+N
350 DDELTA(M)=DDELTA(M)+SUM1(NM)*SUM1(NL)
400 MST=MST+NDIM-M
I1=L1
DO 500 I=L,NPARAM
I2=L1+I
DO 450 M=NSTART,NDIM
IM=I1+M
450 SUM1(I2)=SUM1(I2)+SUM1(IM)*DDELTA(M)
500 I1=I1+NDIM-I
DO 550 M=NSTART,NDIM
IM=L1+M
550 SUM1(IM)=-DDELTA(M)
600 L1=L1+NDIM-L
C PRINT UPDATED ELEMENTS & CORRELATIONS
WRITE(OUTP,44446)
J1=NSAT*6
WRITE(OUTP,10304) ARCNO,OUTER,(ELEMST(J),J=1,J1)
CALL CORREL(SUM1,NPARAM,NDIM,ARCNO,TTL)
WRITE(OUTP,44460) ARCNO
C PRINT UPDATED PARAMETER SIGMAS
DO 650 I=1,NPARAM
II=INDXNO(I)+1
650 WRITE(OUTP,44461) TTL(I),SUM1(II)
C PRINT CROSS CORRELATIONS
WRITE(OUTP,44450) ARCNO
WRITE(OUTP,10214) (TTL(J),J=NSTART,NDIM)
IST=NSTART
ISTP=NDIM
DO 800 I=1,NPARAM
INDEXI=INDXNO(I)
II=INDEXI+1
DO 700 J=NSTART,NDIM
JJ=INDXNO(J)+J
IJ=INDEXI+J
700 SUM1(IJ)=SUM1(IJ)/(SUM1(II)*DSORT(SUM1(JJ)))
WRITE(OUTP,10215) TTL(I),(SUM1(J),J=IST,ISTP)
IST=IST+NDIM-I
800 ISTP=ISTP+NDIM-I
C STORE UPDATED ARC INFORMATION
1000 CALL DATARC(ARCNO,.FALSE...TRUE...FALSE.)
RETURN
10214 FORMAT(1H0,6X,18A5)
10215 FORMAT(1H0,A6,18F6.3/(7X,18F6.3))
10304 FORMAT(1H0//,40X,24HUPDATED ELEMENTS FOR ARC,13,22H AFTER OUTER ITERATION,12//
. (44X,1HX,25X,1HY,25X,1H7//,30X,3024,16//,42X,4HX00T,22X,
. 4HY00T,22X,4H700T//,30X,3024,16//)
44446 FORMAT(1H1)
44450 FORMAT(1H0/7X,'CROSS CORRELATION COEFFICIENTS BETWEEN STATION ',
I ' POSITIONS AND ARC',13,' ADJUSTED PARAMETERS')
44460 FORMAT(1H0,10X,'ADJUSTED PARAMETER SIGMAS FOR ARC',13/)
44461 FORMAT(13X,A6,G10,3)
END

```

UPDA 112
UPDA 113
UPDA 114
UPDA 115
UPDA 116
UPDA 117
UPDA 118
UPDA 119
UPDA 120
UPDA 121
UPDA 122
UPDA 123
UPDA 124
UPDA 125
UPDA 126
UPDA 127
UPDA 128
UPDA 129
UPDA 130
UPDA 131
UPDA 132
UPDA 133
UPDA 134
UPDA 135
UPDA 136
UPDA 137
UPDA 138
UPDA 139
UPDA 140
UPDA 141
UPDA 142
UPDA 143
UPDA 144
UPDA 145
UPDA 146
UPDA 147
UPDA 148
UPDA 149
UPDA 150
UPDA 151
UPDA 152
UPDA 153
UPDA 154
UPDA 155
UPDA 156
UPDA 157
UPDA 158
UPDA 159
UPDA 160
UPDA 161
UPDA 162
UPDA 163
UPDA 164
UPDA 165
UPDA 166
UPDA 167
UPDA 168



NAME UPDOWN
PURPOSE TO COMPUTE UPLINK AND DOWNLINK TRANSIT TIME FOR AVERAGE RANGE RATE DATA OF THE TWO DOWNLINK TRANSIT TIMES FOR VLBI DATA
CALLING SEQUENCE CALL UPDOWN(DAY,DSTA1,DSTA2,SIGN,ISTA1,ISTA2,ISAT,PREPRO,OBS,RFINDX)

SYMBOL	TYPE	DESCRIPTION
DAY	DP	INPUT & OUTPUT - SATELLITE TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR INPUT - NOMINAL ESTIMATE OUTPUT - CORRECTED SATELLITE TIME
DSTA1	DP	INPUT - FIRST GROUND TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR
DSTA2	DP	OUTPUT - SECOND GROUND TIME IN DAYS FROM JAN 0.0 OF THE REFERENCE YEAR
SIGN	DP	INPUT - =+1 FOR VLBI DATA =-1 FOR AVERAGING RANGE RATE DATA
ISTA1	I	INPUT - INDEX FOR FIRST STATION
ISTA2	I	INPUT - INDEX FOR SECOND STATION
ISAT	I+2	INPUT - SATELLITE INDEX
PREPRO	L	INPUT - SWITCH REQUESTING TROPOSPHERIC REFRACTION PREPROCESSING
OBS	DP	OUTPUT - VALUE OF MEASUREMENT
RFINDX (2)	R	INPUT - REFRACTION INDICES FOR MEASUREMENT PREPROCESSING

SUBROUTINES USED GRBIT GRHRAN

COMMON BLOCKS CUVECT

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES 'GEODYN SYSTEMS DESCRIPTION'
VOLUME 1 - GEODYN DOCUMENTATION

SUBROUTINE UPDOWN(DAY,DSTA1,DSTA2,SIGN,ISTA1,ISTA2,ISAT,PREPRO, OBS,RFINDX) UPDOWN SA
UPDOWN SS

IMPLICIT REAL*8 (A-H,C-Z)	UPDN	54
LOGICAL NOT1ST,PREPRO	UPDN	57
INTEGER*2 ISAT	UPDN	58
REAL RFINDX	UPDN	59
DIMENSION RFINDX(2)	UPDN	60
COMMON/CUVECT/UHAT(3,2),XYZ(3,2),RXYZ(3,2),RENV(3,2),R(2),	UPDN	61
RSQ(2),XYSQ(2)	UPDN	62
DATA C/2.99792508/,DTOL/1.00-10/,NOT1ST/,FALSE./	UPDN	63
IF(NOT1ST) GO TO 100	UPDN	64
NOT1ST=.TRUE.	UPDN	65
DAYLIT=1.000/(C*B.6404)	UPDN	66
100 N=C	UPDN	67
200 DPREV=DAY	UPDN	68
CALL ORBIT(CAY)	UPDN	69
THETG=GRHRAN(DSTA1,ISTA1)	UPDN	70
DAY=DSTA1-R(ISAT)*DAYLIT	UPDN	71
DT=CABS(DAY-DPREV)	UPDN	72
IF(DT.LE.DTOL) GO TO 300	UPDN	73
N=N+1	UPDN	74
IF(N.LE.5) GO TO 200	UPDN	75
DT=DT*B.6404	UPDN	76
PRINT 1000,DT,DTOL	UPDN	77
300 IF(PREPRO) OBS=OBS-RFINDX(1)/(C.02600+RENV(3,ISAT))	UPDN	78
DSTA2=SIGN*(DSTA1-DAY)+CAY	UPDN	79
N=0	UPDN	80
400 DPREV=DSTA2	UPDN	81
THETG=GRHRAN(DSTA2,ISTA2)	UPDN	82
DSTA2=DAY+SIGN*R(ISAT)*DAYLIT	UPDN	83
DT=CABS(DSTA2-DPREV)	UPDN	84
IF(DT.LT.DTOL) GO TO 500	UPDN	85
N=N+1	UPDN	86
IF(N.LE.5) GO TO 400	UPDN	87
DT=DT*B.6404	UPDN	88
PRINT 1000,DT,DTOL	UPDN	89
500 IF(PREPRO) OBS=OBS-RFINDX(2)/(C.02600+RENV(3,ISAT))	UPDN	90
RETURN	UPDN	91
1000 FORMAT(' ***UPDOWN**\$ ACCEPTED TRANSIT TIME ERROR AFTER SIX ',	UPDN	92
• ' ITERATIONS =',F12.5,' SECONDS. GREATER THAN',E12.5,	UPDN	93
• ' DAYS ***UPDOWN**\$')	UPDN	94
END	UPDN	95

NAME VCONV

PURPOSE TO CONVERT VARIANCE-COVARIANCE FROM ONE SYSTEM TO ANOTHER

CALLING SEQUENCE CALL VCONV(VARIN,VAROUT,PARTL)

SYMBOL	TYPE	DESCRIPTION
VARIN	R	INPUT - INPUT VARIANCE-COVARIANCE MATRIX
VAROUT	R	OUTPUT - OUTPUT VARIANCE-COVARIANCE MATRIX
PARTL	R	INPUT - PARTIALS OF -VAROUT- VARIABLES WITH RESPECT TO -VARIN- VARIABLES

SUBROUTINES USED NONE

COMMON BLOCKS NONE

INPUT FILES NONE

OUTPUT FILES NONE

	SUBROUTINE VCONV(VARIN,VAROUT,PARTL)	VCON	28
	DIMENSION VARIN(3,3),VAROUT(3,3),PARTL(3,3),T(3)	VCON	29
C	PRE-MULTIPLY INPUT MATRIX BY TRANSPOSE OF PARTIAL MATRIX	VCON	30
	DO 10 I=1,3	VCON	31
	DO 10 J=1,3	VCON	32
	VAROUT(I,J)=0.	VCON	33
	DO 10 K=1,3	VCON	34
10	VAROUT(I,J)=VAROUT(I,J)+PARTL(K,I)*VARIN(K,J)	VCON	35
C	POST-MULTIPLY ABOVE BY PARTIAL MATRIX	VCON	36
	DO 30 I=1,3	VCON	37
	DO 20 J=1,3	VCON	38
	T(J)=0.	VCON	39
	DO 20 K=1,3	VCON	40
20	T(J)=T(J)+VAROUT(I,K)*PARTL(K,J)	VCON	41
	DO 30 J=1,3	VCON	42
30	VAROUT(I,J)=T(J)	VCON	43
	RETURN	VCON	44
	END	VCON	45

VEVAL

DESCRIPTION

VEVAL is a major subroutine in GEODYN and is closely linked with the force model subroutines. Its purpose is to evaluate the variational equations.

Various intermediate data is computed in other routines, especially the direct partial derivatives of the accelerations with respect to the force model coefficients being determined. SUNGRV and EGRAV also supply information for computing the partial derivatives of the gradient of the gravitational potential with respect to the position of the satellite at the current time, i.e.

$$\frac{\partial}{\partial \bar{r}} (\nabla U)$$

Subroutine DENSTY supplies the partial derivative of the atmospheric density with respect to spheroid height.

The order of computation is:

- compute $U_{2c} = \frac{\partial}{\partial \bar{r}} (\nabla U)$;
- compute $D_r = \frac{\partial}{\partial \bar{r}} (D)$, where D is the acceleration due to drag;
- evaluate the variational equations.

NAME VEVAL
ENTRY POINT PURPOSE
VEVAL1 INITIALIZATION
VEVAL TO COMPUTE PARTIALS OF ACCELERATION WITH RESPECT TO INSTANTANEOUS ORBITAL ELEMENTS. TO MULTIPLY THESE PARTIALS BY THE VARIATIONAL PARTIALS TO OBTAIN THE VARIATIONAL ACCELERATION FOR INTEGRATION OF THE VARIATIONAL EQUATIONS

CALLING SEQUENCE CALL VEVAL1(GRPAR)

SYMBOL	TYPE	DESCRIPTION
GRPAR	DP	INPUT - PARTIALS OF FORCE MODEL PARAMETERS
(3,1)		

CALLING SEQUENCE CALL VEVAL(XI,FCT,MDIM,FEVAL,M2)

SYMBOL	TYPE	DESCRIPTION
XI	DP	INPUT - ORBITAL ELEMENTS AND VARIATIONAL PARTIALS
(MDIM,1)		
FCT	DP	INPUT - ACCELERATIONS AND VARIATIONAL PARTIALS
(3,1)		
MDIM	I	INPUT - DIMENSION OF XI IN THE CALLING PROGRAM
FEVAL	L	INPUT - FLAG TO DETERMINE WHICH ACCELERATION PARTIALS ARE REQUIRED: TRUE : WITH RESPECT TO EPOCH ELEMENTS AND PARAMETERS FALSE : WITH RESPECT TO INSTANTANEOUS ELEMENTS
M2	I	INPUT - VARIATIONAL EQUATION DISPLACEMENT

SUBROUTINES USED RESPAR CLEAR

COMMON BLOCKS CPARAM DRGBLK FMODEL INTBLK MOOVGR
VMAT VRBLOK XYZ

INPUT FILES NONE

OUTPUT FILES NONE

REFERENCES 'GECODYN SYSTEMS DESCRIPTION'
VOLUME 1 - GECODYN DOCUMENTATION

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

SUBROUTINE VEVAL(GRPAR)	VEVA	57
IMPLICIT REAL*8 (A-H,C-Z)	VEVA	58
LOGICAL FEVAL,CMGPR	VEVA	59
DOUBLE PRECISION MODEL	VEVA	60
DIMENSION GRPAR(3,1),XI(MDIM,1),FCT(3,1),RID(3),PID(3),AL1D(3),	VEVA	61
R2D(3,3),P2D(3,3),AL2D(3,3)	VEVA	62
COMMON/CPARAM/NSTA,NMAST,NSTEST,NDIM,NBIAS,NGPC1,NGPC2,	VEVA	63
NGPCM,NCSEST,CMGPR,LIM1,LIM2,NDEN,NDENST,NTIDST,NTIDEN,	VEVA	64
INNRSW,NCONST,NCCNS	VEVA	65
COMMON/DRGBLK/HT,SP,SI,SO,EXPT(3),CO,SBRHO,EBRHOV,VEL,VELR(3)	VEVA	66
COMMON/FMODEL/INDEX1,INDEX2,INDEX3,INDEX4,CS(30,33),MODEL(8)	VEVA	67
COMMON/INTELK/THOOT1(2),THOT25,GM,AE,AESC(2),FSQ32,FFSQ32,	VEVA	68
GM3(6),B(2),BDOT(41),NEQNS(15),NEODY	VEVA	69
COMMON/POBNGR/DPXUV(6),RHO(3,6),RHOC(6),RHO3(6)	VEVA	70
COMMON/VMAT/VMATRX(3,6),U2D(3,3),C1DER(3,3),C2DER(3,3,3)	VEVA	71
COMMON/VRBLCK/A1,CSLM(31),SNLM(31),UID(3),P(33,30),AORN(30),	VEVA	72
TPM(39)	VEVA	73
COMMON/XYZ/X(6),R1,R2,ISAT,IFORCE(2)	VEVA	74
EQUIVALENCE (PID(1),C1DER(1,1)),(PID(1),C1DER(1,2)),	VEVA	75
(AL1D(1),C1DER(1,3)),(SP,P(1,1)),(CP,P(2,1)),(R2D(1,1),	VEVA	76
C2DER(1,1,1)),(P2D(1,1),C2DER(1,1,2)),(AL2D(1,1),C2DER(1,1,3))	VEVA	77
EQUIVALENCE (PDPHDD,EXPT(1))	VEVA	78
RETURN	VEVA	79
ENTRY VEVAL(XI,FCT,MDIM,FEVAL,M2)	VEVA	80
C-DATA IN COEFFICIENT PARTIALS IF GEOPOTENTIAL IS ADJUSTED	VEVA	81
IF (CMGPR) CALL RESPAR	VEVA	82
CALL CLEAR(VMATRX,63,2)	VEVA	83
RINV=1.000/R1	VEVA	84
R3=R1*R2	VEVA	85
AL1D(1)=X(2)/A1	VEVA	86
AL1D(2)=X(1)/A1	VEVA	87
AL2D(1,1)=-2.000*AL1D(1)*AL1D(2)	VEVA	88
AL2D(2,2)=-AL2D(1,1)	VEVA	89
AL2D(1,2)=AL1D(1)**2-AL1D(2)**2	VEVA	90
UID(2)=UID(2)/CP	VEVA	91
C7=1.000/CP**2	VEVA	92
NEQN=NEQNS(ISAT)	VEVA	93
C COMPUTE PARTIALS OF GEOPOTENTIAL WITH RESPECT TO R, PHI, LAMBDA	VEVA	94
DO 110 NC=2,INDEX4	VEVA	95
NS=31-NC	VEVA	96
FNI=NC+1	VEVA	97
FM=1.00	VEVA	98
C0=P(2,NC)	VEVA	99
MMX=NC+1	VEVA	100
DO 110 MC=1,MMX	VEVA	101
MS=34-MC	VEVA	102
FM=FM+1.00	VEVA	103
P1=P(MC,NC)	VEVA	104
C1=(CS(MC,MC)*CSLM(MC)+CS(NS,MS)*SNLM(MC))*AORN(MC)	VEVA	105
C2=(-CS(MC,MC)*SNLM(MC)+CS(NS,MS)*CSLM(MC))*ACRN(MC)*FM	VEVA	106
C3=C0	VEVA	107
CC=P(MC+2,NC)-TFM(MC+1)*P(MC+1,NC)	VEVA	108
U2D(1,1)=L2D(1,1)+FNI*C1*(FNI+1.00)*P1	VEVA	109
U2D(2,1)=L2D(2,1)-FNI*C1*C3	VEVA	110
L2D(3,1)=L2D(3,1)-FNI*C2*P1	VEVA	111

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

```

U2D(3,2)=L2D(3,2)+C2*C3
U2D(3,3)=U2D(3,3)-FM**2*C1*P1
110 U2D(2,2)=L2D(2,2)+C1*(C0-TPM(MC)*C3-FM*P1*C7)
L2D(1,1)=2.0D0*GM/R3+U2D(1,1)/R2
U2D(2,1)=L2D(2,1)/(R1*CPI)
U2D(3,1)=L2D(3,1)/R1
U2D(2,2)=L2D(2,2)*C7+UID(2)*TPM(2)/CP
L2D(3,2)=L2D(3,2)/CP
L2D(1,2)=U2D(2,1)
L2D(1,3)=L2D(3,1)
U2D(2,3)=L2D(3,2)
C=X(3)/R2
DO 205 I=1,3
RID(I)=X(I)/R1
205 PID(I)=-C*RID(I)
PID(3)=PID(3)+RINV
DO 210 I=1,3
P2D(I,I)=-X(3)/R3
IF(I.EQ.3) P2D(3,3)=P2D(3,3)+P2C(3,3)
C1=3.0D0*X(3)*X(I)/R2
DO 210 J=I,3
IF(J.EQ.3) P2D(I,J)=P2D(I,J)-X(I)/R3
210 P2D(I,J)=P2D(I,J)+C1*X(J)/R3
DO 215 I=1,3
DO 216 J=I,3
214 R2D(I,J)=-X(I)*X(J)/P3
215 R2D(I,I)=R2D(I,I)+RINV
C COMPUTE ACCELERATION PARTIAL COMPONENTS FROM GEOPOTENTIAL, MOON & SUN
DO 175 I=1,3
DO 178 K=1,3
C=0.0D0
DO 177 J=1,3
177 C=C+CIDER(I,J)*U2D(J,K)
DO 178 L=1,3
178 VMATRX(I,L)=VMATRX(I,L)+C*CIDER(L,K)
DO 180 N=1,NSOY
IF(GM3(N).LE.0.0D0) GO TO 180
C7=GM3(N)/RHOS(N)
C8=3.0D0*C7/RHOSQ(N)
DO 179 I=1,3
VMATRX(I,I)=VMATRX(I,I)-C7
DO 179 J=I,3
179 VMATRX(I,J)=VMATRX(I,J)+C8*RHOM(I,N)*RHOM(J,N)
180 CONTINUE
DO 190 I=1,3
DO 190 J=1,3
DO 185 K=1,3
185 VMATRX(I,J)=VMATRX(I,J)+UID(K)*C2DER(I,J,K)
190 IF(I.NE.J) VMATRX(J,I)=VMATRX(I,J)
NN=3
IF(.NOT.B(I SAT).GT.0.0D0) GO TO 300
C COMPUTE ACCELERATION PARTIAL COMPONENT FROM DRAG
NN=5
BBRHQ=BBRHQV/VEL
BRHOTH=BBRHC*THOT2S
BRHCV=BBRHQ/VEL

```

VEVA 112
VEVA 113
VEVA 114
VEVA 115
VEVA 116
VEVA 117
VEVA 118
VEVA 119
VEVA 120
VEVA 121
VEVA 122
VEVA 123
VEVA 124
VEVA 125
VEVA 126
VEVA 127
VEVA 128
VEVA 129
VEVA 130
VEVA 131
VEVA 132
VEVA 133
VEVA 134
VEVA 135
VEVA 136
VEVA 137
VEVA 138
VEVA 139
VEVA 140
VEVA 141
VEVA 142
VEVA 143
VEVA 144
VEVA 145
VEVA 146
VEVA 147
VEVA 148
VEVA 149
VEVA 150
VEVA 151
VEVA 152
VEVA 153
VEVA 154
VEVA 155
VEVA 156
VEVA 157
VEVA 158
VEVA 159
VEVA 160
VEVA 161
VEVA 162
VEVA 163
VEVA 164
VEVA 165
VEVA 166
VEVA 167

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS POOR

BRHTDV=BRHDV*THD T2S	VEVA 169
C1=2.00*F SQ32*SPSISO-FFSQ32	VEVA 169
C2=2.00*C1/R1	VEVA 170
C3=(SPSISO-1.00)*C2	VEVA 171
C2=SPSISO*C2	VEVA 172
C0=BRHCV*PDPHOD/R1	VEVA 173
C1=(1.00+C2)*C0	VEVA 174
VELR(3)=X(6)	VEVA 175
EXPT(1)=X(1)*C1	VEVA 176
EXPT(2)=X(2)*C1	VEVA 177
EXPT(3)=X(3)*(C3+1.00)*C0	VEVA 178
C1=SRHTDV*VELR(1)*VELR(2)	VEVA 179
VMATRIX(1,2)=VMATRIX(1,2)-BRHOTH*(VEL+VELR(1)**2/VEL)	VEVA 180
VMATRIX(2,1)=VMATRIX(2,1)+BRHOTH*(VEL+VELR(2)**2/VEL)	VEVA 181
VMATRIX(1,1)=VMATRIX(1,1)+C1	VEVA 182
VMATRIX(2,2)=VMATRIX(2,2)-C1	VEVA 183
VMATRIX(3,1)=VMATRIX(3,1)+BRHTDV*VELR(3)*VELR(2)	VEVA 184
VMATRIX(3,2)=VMATRIX(3,2)-BRHTDV*VELR(3)*VELR(1)	VEVA 185
DO 200 I=1,3	VEVA 186
DO 200 J=1,3	VEVA 187
J1=J+3	VEVA 188
IF(I.EQ.J) VMATRIX(I,J1)=VMATRIX(I,J1)-EBRHCV	VEVA 189
VMATRIX(I,J1)=VMATRIX(I,J1)-VELR(I)*EXPT(J)	VEVA 190
200 VMATRIX(I,J1)=VMATRIX(I,J1)-BRHDV*VELR(I)*VELR(J)	VEVA 191
300 DO 301 J=1,6	VEVA 192
K0=(J-1)*M2+1	VEVA 193
DO 301 I=1,3	VEVA 194
301 FCT(I,K0)=0.00	VEVA 195
IF(NEON.LT.8) GO TO 304	VEVA 196
C COMPUTE ACCELERATION PARTIALS WITH RESPECT TO EPOCH PARAMETERS	VEVA 197
DO 302 L1=8,NEON	VEVA 198
K0=(L1-2)*M2+1	VEVA 199
K0C=L1-7	VEVA 200
DO 302 I=1,3	VEVA 201
302 FCT(I,K0)=GRPAR(I,K00)	VEVA 202
304 IF(.NOT.FEVAL) RETURN	VEVA 203
DO 305 L1=2,NEON	VEVA 204
K0=(L1-2)*M2+1	VEVA 205
DO 305 I=1,3	VEVA 206
SUM=0.00	VEVA 207
DO 306 J=1,NN	VEVA 208
306 SUM=SUM+VMATRIX(I,J)*XI(J,L1)	VEVA 209
305 FCT(I,K0)=FCT(I,K0)+SUM	VEVA 210
RETURN	VEVA 211
END	VEVA 212

NAME YMDAY
PURPOSE TO COMPUTE FOR A GIVEN DATE THE NUMBER OF DAYS FROM
JAN 0.0 OF THE REFERENCE YEAR FOR THE ARC
CALLING SEQUENCE X=YMDAY(IYMD,IHM,SEC)

SYMBOL	TYPE	DESCRIPTION
IYMD	I	INPUT - DATE IN THE FORM OF YYMMDD
IHM	I	INPUT - TIME IN THE FORM OF HHMM
SEC	R	INPUT - SECONDS
YMDAY	DP	OUTPUT - NUMBER OF DAYS FROM JAN 0.0 OF THE REFERENCE YEAR FOR THE ARC

SUBROUTINES USED DIFF

COMMON BLOCKS CTIME

INPUT FILES NONE

OUTPUT FILES NONE

DOUBLE PRECISION FUNCTION YMDAY(IYMD,IHM,SEC)	YMDA	20
IMPLICIT REAL*8 (A-H,C-Z)	YMDA	31
COMMON/CTIME/DAYREF(11),IYREG	YMDA	32
IY=IYREG*10000+101	YMDA	33
IHMS=IHM*100	YMDA	34
CALL DIFF(IY,0,IYMD,IHMS,ID,IS)	YMDA	35
YMDAY=85400*(ID+1)+IS	YMDA	36
YMDAY=(YMDAY+SEC)/8.6404	YMDA	37
RETURN	YMDA	38
END	YMDA	39

SECTION 9.0
COMMON BLOCK DESCRIPTIONS

The GEODYN program contains 39 common blocks. Each common block is fully described on the following pages. Some common blocks have more than one version. Each version is described.

/ALPMRC/

COMMON/ALPMRC/ITNMS(5),TIMING,BLANK,ATYPE(31),
UNITS(15),ELCUT,HYPER

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
ITNMS (5)	R*8	000000	Alphanumeric information for printout.	BLOCK DATA	INOIPT
TIMING	R*8	000028	Alphanumeric information for printout.	BLOCK DATA	NONAME INOIPT
BLANK	R*8	000030	Alphanumeric information for printout.	BLOCK DATA	NONAME INOIPT SUMMARY
ATYPE (31)	R*8	000038	Alphanumeric information for printout.	BLOCK DATA	NONAME INOIPT SUMMARY TYPORB
UNITS (15)	R*8	000130	Alphanumeric information for printout.	BLOCK DATA	NONAME INOIPT
ELCUT	R*8	0001A8	Elevation cutoff angle.	INOIPT NEWARC	NONAME DATARD INOIPT
HYPER	L*4	0001B0	Hyperbolic element switch.	MAIN	MAIN DATARD ELEM

/APARAM/
 COMMON/APARAM/ INPAR, INPARI, NBIAS,
 ESTSTA, NSAT, NGPARC, NOREC1, NPARAM

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Program Where Used</u>
INPAR	I*4	000000	Number of force model parameters to be integrated.	NONAME NEWARC	NONAME NEWARC PREDCT STORE
INPARI	I*4	000004	Number of force model parameters in arc.	INOIPT ARCPAR	MAIN NONAME ARCPAR ESTIM INOIPT NEWARC STORE
NBIAS	I*4	000008	Number of biases in arc.	INOIPT BIAS	MAIN NONAME ARCPAR BIAS DODSRD ESTIM GEOSRD INOIPT NEWARC SIMRD STORE
ESTSTA	I*4	00000C	Number of adjusted stations.	NEWARC	NONAME NEWARC STORE
NSAT	I*4	000010	Number of satellites in arc.	MAIN	MAIN NONAME APPER ARCPAR BMTWRT BSCOMP ESTIM GEOSRD GRHRAN INOIPT NEWARC ORBIT PREDCT SIMRD STORE

/APARAM/ (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Program Where Used</u>
NGPARC	I*4	000014	Number of adjusted arc geopotential coefficients	INOIPT	MAIN NONAME ARCPAR BIAS DATARD ESTIM INOIPT NEWARC STORE UPDATE
RECNO1	I*4	000018	Number of record of first observation in arc.	INOIPT	NONAME INOIPT NEWARC STORE
NPARAM	I*4	00001C	Number of parameters in arc.	NONAME	NONAME BSCOMP DATARD ESTIM NEWARC STORE SUMMRY UPDATE
NEBIAS	I*4	0003DC	Number of electronic biases in arc.	INOIPT	MAIN NONAME ARCPAR BSCOMP CBROWN DATARD INOIPT NEWARC STORE
MAXPAR	I*4	00000C	Number of parameters per measurement for iteration.	ESTIM	BSCOMP ESTIM NEWARC STORE

/CELEM/

COMMON/CELEM/ELEMST(6,2),ORBELA(6,2),XNU,EC,RMSTOT

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
ELEMST (6,2)	R*8	000000	Epoch cartesian elements.	MAIN NONAME DODELM ESTIM ORBIT UPDATE	MAIN NONAME COWELL DATARD DODELM ESTIM INOIPT ORB1 ORBIT STORE TYPORB UPDATE
ORBELA (6,2)	R*8	000060	Epoch Kepler elements.	MAIN NONAME DODELM ORBIT	MAIN NONAME APPER DATARD DODELM INOIPT ORB1 ORBIT STORE TYPORB
XNU	R*8	0000C0	True anomaly.	ELEM	DATARD ELEM ORB1 STORE
EC	R*8	0000C8	Eccentric anomaly.	ELEM	DATARD ELEM ORB1 STORE
RMSTOT	R*4	0000D0	Total RMS	NONAME INOIPT	NONAME DATARD INOIPT NEWARC STORE SUMMRY TYPORB

/CEPHEM/ (Version 1)
COMMON/CEPHEM/AO,PMOON,SUN,ANUT,DUMMY

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Program Where Use</u>
A0 (25)	R*8	000000	Unit vector and center of mass distance from earth to MOON SUN VENUS MARS JUPITER SATURN plus A0(25) = nutation in right ascension (equation of the equinoxes).	EPHEM	DENSTY EPHEM F GRHRAN PROCES SUNGRV TIDAL
PMOON (306)	R*8	0000C8	Buffer for lunar position and inter- polation informa- tion	EPHEM	EPHEM
SUN (270)	R*8	000A58	Buffer for solar and planetary position and inter- polation information	EPHEM	EPHEM
ANUT (102)	R*4	0012C8	Buffer for nuta- tion information	EPHEM	EPHEM
DUMMY	R*8		-Dummy		

/CEPHEM/ (Version 2)

COMMON/CEPEHM/JNAME, ISTAR, ESTANO, ISTAR

Variable	Type	Hex Location	Description	Program Where Defined	Program Where Used
JNAME (381)	R*8	000000	Names of stations read from input	INOUP	DODSRD INOUP
ISTAR (381)	I*2	00BE8	Numbers of stations read from input	INOUP	DODSRD, GEOSRD INOUP SIMRD
ESTANO (381)	I*2	000EE2	Master station array for station adjustment	INOUP	INOUP STAINP
ESTANO (386)	I*2	0011DC	Number of stations to be used	INOUP DODSRD GEOSRD SIMRD	BIAS DODSRD GEOSRD INOUP SIMRD STAINP

/CGEOS/ (Version 1)

COMMON/CGEOS/ISATID(2),THETGO(15),IG6(423)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
ISATID(2)	R*4	000000	---	---	Not Used
THETGO (15)	R*8	000008	Right ascension of Greenwich in degrees on Jan.0.0 from 1958-1975.	JANTHG	JANTHG
IG6(423)	R*4	000080	---	---	Not Used

/CGEOS/ (Version 2)

COMMON/CGEOS/ISATID(2), IPREPR(4,50), RFINDX(2,50),
 INDPRE(2,50), NOPRPR, NSIG, NCULL, SIGCHG(50),
 IMTYPE(50), ISTNO(50), CULL(2,100)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
ISATID (2)	I*4	000000	Satellite ID's.	INOUP	NONAME DODSRD GEOSRD INOUP ORBI PROCES SIMRD TYPORB
IPREPR (4,50)	I*2	000008	Preprocessing indicators.	INOUP	DODSRD GEOSRD INOUP PRNTPR
RFINDX (2,50)	R84	000198	Tropospheric refraction indices and constant timing corrections	INOUP	DODSRD GEOSRD INOUP PRNTPR
INDPRE (2,50)	I*2	000328	Station numbers and measurement types for preprocessing.	INOUP	DODSRD GEOSRD INOUP PRNTPR
NOPRPR	I*4	0003F0	Number of PREPO cards input.	INOUP	DODSRD GEOSRD INOUP NEWARC PRNTPR
NSIG	I*4	0003F4	Number of SIGMA cards input.	INOUP	DODSRD GEOSRD INOUP NEWARC PCERD SIMRD
NCULL	I*4	0003F8	Number of CULL sets.	INOUP	DODSRD GEOSRD INOUP NEWARC PCERD SIMRD

/CGEOS/ (version 2) (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
SIGCHG (50)	R*4	0003FC	Input sigma changes.	INOIPT	DODSRD GEOSRD INOIPT NEWARC PCERD SIMRD
IMTYPE (50)	I*2	0004C4	Sigma change types.	INOIPT	DODSRD GEOSRD INOIPT NEWARC PCERD SIMRD
ISTNO (50)	I*2	000528	Sigma change stations.	INOIPT	DODSRD GEOSRD INOIPT NEWARC PCERD SIMRD
CULL (2,100)	I*2	0058C	Cull sets.	INOIPT	DODSRD GEOSRD INOIPT NEWARC PCERD SIMRD

/CONOUT/

COMMON/CONOUT/RMSALL, OUTCON, MINOUT, MAXOUT, LITRES,
MAXSAT, MAX2IN, NSTART, NEQNMX, IVAR, IORDER, NARCS,
NSTARD, LSTART (6)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
RMSALL	R*4	000000	RMS for all arcs.	MAIN	MAIN NONAME
OUTCON	R*4	000004	Outer iteration convergence criterion.	INOIPT	NONAME INOIPT
MINOUT	I*4	000008	Minimum number of outer iterations.	INOIPT	NONAME INOIPT
MAXOUT	I*4	00000C	Maximum number of outer iterations.	INOIPT	NONAME INOIPT
LITRES	L*4	000010	Indicates that adjustment not requested on last inner iteration.	INOIPT	NONAME INOIPT
MAXSAT	I*4	000014	Maximum number of satellites per arc.	MAIN	MAIN CBROWN DATARD
MAX2IN	I*4	000018	Maximum number of inner iterations on outer iterations after first outer iteration.	NONAME	NONAME
NSTART	I*4	00001C	Parameter number of first common parameter.	COMPAR ESTIM	NONAME CBROWN COMPAR ESTIM
NEQNMX	I*4	000020	Maximum number of force model equations to be integrated for one arc.	MAIN	MAIN CBROWN ESTIM ORBIT

/CONOUT/ (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
IVAR	I*4	000024	=1 for fixed step integration =2 for variable step integration	MAIN	MAIN
IORDER	I*4	000028	Not Used.		
NARCS	I*4	00002C	Number of arcs.	MAIN	MAIN NONAME ARCPAR
NSTARD	I*4	000030	Number of tracking stations read from input.	INOIPT	COMPAR INOIPT
STARTR	L*4	000034	.TRUE.- Restart tape input.	INOIPT	NONAME INOIPT
STARTW	L*4	000038	.TRUE.- Restart tape output.	INOIPT	NONAME DATARD INOIPT
STARTA	I*4	00003C	Number of arc where restart will begin.	INOIPT	NONAME INOIPT
STARTO	I*4	000040	Number of outer iteration where restart will begin. (not used)	INOIPT	NONAME INOIPT
INSTRT	I*4	000044	Input restart tape number.	INOIPT	NONAME INOIPT
OUTSTR	I*4	000048	Output restart tape number.	INOIPT	NONAME DATARD INOIPT

/CONSTS/

COMMON/CONSTS/DPI,DTWOPI,DRAD,DRSEC

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Defined</u>
DPI	R*8	000000	π	BLOCK DATA	NONAME AVGPOT DENSTY PREDCT PROCES TYPORB
DTWOPI	R*8	000008	2π	BLOCK DATA	NONAME DENSTY DPFCT ELEM GEOSRD GRHRAN NEWARC ORBI PREDCT PROCES TYPORB
DRAD	R*8	000010	$2\pi/360$ Conversion factor for con- verting degrees to radians.	BLOCK DATA	NONAME APPER ARCPAR AREAS AVGPOT DELTAZ DODELM ELEM GEOSRD INDENT JANTHG NEWARC ORBI PCERD POSVEL PREDCT STAINP TWPSTA TYPORB

/CONSTS/ (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
DRSEC	R*8	000018	$2\pi/360/3600$	BLOCK DATA	NONAME COMADJ COMPAR DODSRD GEOSRD INOIPT PCERD PROCES SIMRD SUMMRY

/CORB1/

COMMON/CORB1/RANDOT(2), PERDOT(2), PERHT(2), APHT(2),
PRD(2)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
RANDOT(2)	R*8	000000	Time derivative of the right ascension of the ascending node.	NONAME	NONAME INOIPT ORB1 PREDCT TWOSTA TYPORB
PERDOT(2)	R*8	000010	Time derivative of the argument of perigee.	NONAME	NONAME ORB1 PREDCT TWOSTA TYPORB
PERHT(2)	R*8	000020	Perigee height.	APPER	APPER INOIPT ORB1 TYPORB
APHT(2)	R*8	000030	Apogee height.	APPER	APPER ORB1 TYPORB
PRD(2)	R*8	000040	Orbital period.	NONAME	NONAME INOIPT ORB1 PREDCT TYPORB

/CPARAM/

COMMON/CPARAM/ NSTA,NMAST,NSTEST,NDIM,MBIAS
 NGPC1,NGPC2,NGPCOM,NCSEST,CMPGPR,
 LIM1,LIM2,NDEN,NDENST,NTIDST,
 NTIDEN,INNRSW,NCONST,NDCONS

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Program Where Used</u>
NSTA	I*4	000000	Number of tracking stations.	DODSRD GEOSRD SIMRD INOIPT CBROWN	MAIN ARCPAR CBROWN COMPAR CONSTS DODSRD GEOSRD INOIPT SIMRD SUMMRY
NMAST	I*4	000004	Number of adjusted master stations.	INOIPT	MAIN NONAME BMTWRT CBROWN COMADJ COMPAR CONSTS ESTIM INOIPT PDEN
NSTEST	I*4	000008	Number of estimated stations.	INOIPT	MAIN NONAME BMTWRT CBROWN COMADJ COMPAR CONSTS ESTIM INOIPT
NDIM	I*4	00000C	Maximum dimension of normal matrix.	MAIN	MAIN NONAME BMTWRT BSCOMP CBROWN COMADJ COMPAR CONSTS DATARD ESTIM PDEN UPDATE

/CPARAM/ (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Program Where Used</u>
MBIAS	I*4	000010	Maximum number of biases plus drag and solar radiation pressure parameters estimated in any one arc.	MAIN	MAIN ARCPAR CBROWN CONSTS DATARD ESTIM UPDATE
NGPC1	I*4	000014	Relative location of first common adjusted geopotential coefficient.	MAIN	MAIN CONSTS DATARD ESTIM
NGPC2	I*4	000018	Relative location of last common adjusted geopotential coefficient.	MAIN	MAIN CONSTS ESTIM
NGPCOM	I*4	00001C	Number of common adjusted geopotential coefficients.	MAIN	MAIN NONAME ARCPAR BMTWRT BSCOMP COMADJ COMPAR CONSTS DATARD ESTIM
NCSEST	I*4	000020	Number of adjusted geopotential coefficients in this iteration.	INOIPT NONAME	MAIN NONAME ARCPAR BMTWRT CBROWN COMADJ COMPAR CONSTS DATARD INOIPT RESPAR

/CPARAM/ (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Program Where Used</u>
CMPGPR	L*4	000024	Logical switch for geopotential partial computations. True-partial are computed.	NONAME	MAIN NONAME ARCPAR CONSTS VEVAL
LIM1	I*4	000028	Size of normal arrays to be cleared.	NONAME	NONAME CONSTS ESTIM
LIM2	I*4	00002C	.GT.0 indicates last inner iteration.	NONAME	NONAME CONSTS ESTIM
NDEN	I*4	000030	Number of surface densities.	INOIPT	MAIN NONAME CBROWN COMPAR CONSTS GEOIDH INDENT INOIPT PDEN SURDEN
NDENST	I*4	000034	Number of adjusted surface densities.	INOIPT	MAIN CBROWN COMADJ CONSTS GEOIDH INDENT INOIPT PDEN SURDEN
NTIDST	I*4	000038	Number of adjusted tidal parameters.	CONSTS	MAIN CBROWN CONSTS

/CPARAM/ (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Program Where Used</u>
NTIDEN	I*4	00003C	Number of adjusted tidal and density parameters.	MAIN	MAIN NONAME BSCOMP CBROWN COMADJ COMPAR CONSTS ESTIM RESPAR
INNRSW	L*4	000040	Logical switch. True - last inner iteration and surface densities adjusted.	NONAME	NONAME CONSTS SURDEN
NCONST	I*4	000044	Number of surface density constraint equations.	INOIPT	MAIN CBROWN CONSTS GEOIDH INOIPT PDEN SURDEN
NDCONS	I*4	000048	Maximum degree and order of surface density constraint equations.	INOIPT	CONSTS GEOIDH INOIPT

/CSLIM/

COMMON/CSLIM/LLIMIT,ULIMIT

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
LLIMIT	I*4	000000	LLIMIT(N) is the minimum order plus one of terms of degree N which are used in the geopotential expansion.	EGRAV	EGRAV
ULIMIT	I*4	00007C	ULIMIT(N) is the maximum order plus one of terms of degree N which are used in the geopotential expansion.	EGRAV	EGRAV GEOIDH

/CSTAT/

COMMON/CSTAT/RESID,SIG,NMTOT,WTSUMT

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
RESID	R*8	000000	Measurement residual.	STAINF	STAINF SMSTAT
SIG	R*8	000004	Measurement sigma.	STAINF	STAINF SMSTAT
NMTOT	I*4	000008	Total number of weighted measurements in arc.	STAINF SMSTAT	STAINF SMSTAT
WTSUMT	R*8	00000C	Sum of squares of weighted residuals for the arc.	STAINF SMSTAT	BMTWRT STAINF SMSTAT

/CSTHET/

COMMON/CSTHET/CTHETG,STHETG

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
CTHETG	R*8	000000	Cosine of right ascension of Greenwich.	GRHRAN SURDEN	DPFCT GRHRAN SURDEN
STHETG	R*8	000008	Sine of right ascension of Greenwich.	GRHRAN SURDEN	DPFCT GRHRAN SURDEN

/CTIME/

COMMON/CTIME/DATAEP, DAYREF, DSTART, DAYSTP, DAYINT,
DORBIT, DAYEND, DRATE, DORBI, DORBIE, ORBRT, IYBEG

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
DATAEP	R*8	000000	Epoch of data in days from Jan 0.0 of the reference year.	MAIN	MAIN NONAME DATARD DENSTY DODELM DODSRD GEOSRD INOIPT ORBIT PCERD SIMRD STORE TYPORB
DAYREF	R*8	000008	Reference date in days from Jan 0.0 of the reference year.	MAIN	MAIN DATARD ORBI REFCOR STORE TYPORB
DSTART	R*8	000010	Epoch in days from Jan 0.0 of the reference year.	MAIN DODELM ORBIT	MAIN NONAME COWELL DATARD DODELM F INOIPT ORBI ORBIT STORE TYPORB

/CTIME/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
DAYSTP	R*8	000018	Data stop time in days from Jan 0.0 of the reference year.	MAIN	MAIN DATARD DODSRD GEOSRD INOIPT NEWARC PCERD SIMRD STORE TYPORB
DAYINT	R*8	000020	Current integration time in days from Jan 0.0 of the reference year.	F	DATARD DENSTY F STORE
DORBIT	R*8	000028	Start time for orbit generator in days from Jan 0.0 of the reference year.	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC STORE
DAYEND	R*8	000030	Stop time for orbit generator in days from Jan 0.0 of the reference year.	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC STORE
DRATE	R*8	000038	Output interval for orbit-generator in days.	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC STORE
DORB1	R*8	000040	Start time for ORB1 tape in days from Jan 0.0 of the reference year.	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC ORB1 STORE

/CTIME/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
DORB1E	R*8	000048	Stop time for ORB1 tape in days from Jan 0.0 of the reference year.	INOIPT NEWARC	DATARD INOIPT ORB1 STORE
ORBRT	R*8	000050	Output interval for ORB1 tape in integral seconds.	INOIPT NEWARC	NONAME DATARD INOIPT STORE
IYBEG	I*4	000058	Reference year.	MAIN	MAIN DATARD DATES STORE YMDAY

/CSTINF/ (Version 1)

COMMON/CSTINF/MEASO(4),NOBS(4),RDMEAN(4),RMSO(4)
 RND(4),MEASWT(4),WTMEAN(4),RMSWTO(4),WTRND(4),
 TYPRMS(4),NOTYPE(2,30)BSUM(8,12),RMSALL(30),
 NOALL(30),NOWTOB,JBASE

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
MEASNO	R*8	000000	Measurement type numbers.	RMSCMP	RMSCMP SUMMARY
NOBS(4)	I*4	000010	Number of measurements.	RMSCMP	RMSCMP STAINF SUMMARY
RDMEAN(4)	R*4	000020	Residual means.	RMSCMP	RMSCMP SUMMARY
RMSO(4)	R*4	000030	RMS's of residuals.	RMSCMP	RMSCMP SUMMARY
RND(4)	R*4	000040	RND's of residuals.	RMSCMP	RMSCMP SUMMARY
MEASWT(4)	I*4	000050	Number of weighted measurements.	RMSCMP	RMSCMP SUMMARY
WTMEAN(4)	R*4	000060	Weighted residual measurements.	RMSCMP	RMSCMP SUMMARY
RMSWTO(4)	R*4	000070	RMS's of weighted residuals.	RMSCMP	RMSCMP SUMMARY
WTRND(4)	R*4	000080	RND's of weighted measurements.	RMSCMP	RMSCMP SUMMARY
TYPRMS(4)	R*4	000090	Measurement type weighted RMS's.	RMSCMP STAINF	NONAME RMSCMP STAINF SUMMARY TYPORB
NOTYPE (2,30)	I*4	000108	Number of measurements by type.	RMSCMP STAINF	RMSCMP STAINF SUMMARY TYPORB

/CSTINF/ (Version 1) (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
BSUM (8,12)	R*4	0001F8	Summing arrays for PCE measurement types.	STAINF	STAINF
RMSALL (300)	R*4	000378	Measurement type weighted RMS's for all arc.	STAINF	STAINF
NOALL (300)	I*4	0003F0	Number of measurements by type for all arcs.	STAINF	NONAME STAINF
NOWTOB	I*4	000468	Total number of weighted observations.	SUMMARY	NONAME BMTWRT SUMMARY TYPORB
JBASE	I*4	00046C	Number of station measurement base lines	GEOSRD SIMRD CBROWN	CBROWN GEOSRD SIMRD SUMMARY

/CSTINF (Version 2)

COMMON/CSTINF/JBASE(283),KBASE(283),LBASE

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
JBASE (283)	I*2	000000	First station in a measurement baseline.	GEOSRD SIMRD	MAIN, GEOSRD SIMRD
KBASE (283)	I*2	000236	Second station in a measurement baseline.	GEOSRD SIMRD	MAIN GEOSRD SIMRD
LBASE	I*4	00046C	Number of station measurement baselines.	GEOSRD SIMRD MAIN	MAIN GEOSRD SIMRD

/CUVECT/

COMMON/CUVECT/UHAT(3,2),XYZ(3,2),RXYZ(3,2),
RENV(3,2),R(2),RSQ(2),XYZSQ(2)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
UHAT(3,2)	R*8	000000	Earth fixed unit vector from station to satellite.	GRHRAN	GRHRAN OBSDOT PREDCT
XYZ(3,2)	R*8	000030	Earth centered fixed satellite vector.	GRHRAN	GRHRAN OBSDOT PREDCT PROCES
RXYZ(3,2)	R*8	000060	Earth fixed vector from station to satellite.	GRHRAN	GRHRAN PREDCT PROCES
RENV(3,2)	R*8	000090	Station-satellite unit local vector (direction cosines).	GRHRAN	GRHRAN OBSDOT PREDCT PROCES TWOSTA UPDOWN
R(2)	R*8	0000C0	Slant range from station to satellite.	GRHRAN	GRHRAN OBSDOT PREDCT PROCES TWOSTA UPDOWN
RSQ(2)	R*8	0000D0	R*R	GRHRAN	GRHRAN PREDCT
XYZSQ(2)	R*8	0000E0	RXYZ(1,ISAT)**2 + RXYZ(2,ISAT)**2 where ISAT = satellite index.	GRHRAN	GRHRAN PREDCT

/DODDAT/
 COMMON/DODDAT/TIME1,STNAM1,OBD1,DG(2),OBSCOR,
 SATNO,IOBNO1,IWT(6),TCOR,IG2(2),IT,IG1,TTAG,PBIT1,PBIT2,IG

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
TIME1	R*8	000000	Time of DODS observation	DATBSE	DATBSE DODSRD
STNAM1	R*8	000008	Station ID	DATBSE	DATBSE DODSRD
OBD1	R*8	000010	DODS observation	DATBSE	DATBSE DODSRD
DG(2)	R*8	000018	DG(1)-not used DG(2)-DODS Range ambiguity information (bits 56-59)	DATBSE	DATBSE DODSRD
OBSCOR	R*8	000028	DODS observation correction	DATBSE	DATBSE DODSRD
SATNO	I*4	000030	Satellite number	DATBSE	DATBSE DODSRD
IOBNO1	I*4	000034	DODS observation number	DATBSE	DATBSE
IWT(6)	R*4	000038	Not Used	DATBSE	
TCOR	R*4	000050	Time correction	DATBSE	DATBSE DODSRD
IG2 (2)	I*2	000054	Not Used	DATBSE	
IT	I*2	000058	DODS observation type number	DATBSE	DATBSE DODSRD
IG1	I*2	00005A	Not Used	DATBSE	
TTAG	I*2	00005C	Time type and station source indicator	DATBSE	DATBSE DODSRD
PBIT1	I*2	00005E	Preprocessing indicators for corrections added prior to DODS (bits 10-15)	DATBSE	DATBSE DODSRD

/DODDAT/ (CONT.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
PBIT2	I*2	000060	Preprocessing indicators for corrections added by DODS (bits 10-15)	DATBSE	DATBSE DODSRD
IG	I*2	000062	Not Used	DATBSE	

/DRGBLK/
COMMON/DRGBLK/HT,SPSISQ, C(4),C3,
C1,VEL,XDOTR,YDOTR,RHO

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
HT.	R*8	000000	DRAG-satellite spheroid height in meters DENSTY-IN- same as in DRAG -OUT- same as in drag multiplied by 10^{-5}	DRAG DENSTY	DENSTY DRAG VEVAL
SPSISQ	R*8	000008	\sin^2 (geocentric latitude)	DRAG	DRAG VEVAL
C(4)	R*8	000010	C(1)=partial of the density with respect to the spheroid height divided by the density C(I), I=2,4 scratch	DENSTY	DENSTY VEVAL
C3	R*8	000030	The density times the relative air speed	DRAG	DENSTY DRAG VEVAL
C1	R*8	000038	$C_3(1/2)C_D(A/m)$	DRAG	DENSTY DRAG VEVAL
VEL	R*8	000040	Relative air speed of the satellite	DRAG	DENSTY DRAG VEVAL
XDOTR	R*8	000048	x-component of the relative air speed	DRAG	DENSTY DRAG VEVAL
YDOTR	R*8	000050	y-component of the relative air speed	DRAG	DENSTY DRAG VEVAL
RHO	R*8	000058	Atmospheric density in kg/m^3	DRAG	DRAG VEVAL

/FERMSG/
COMMON/FERMSG/IMES(26)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
IMES (26)	I*4	000000	See IBM SYSTEM/360 GENERAL I/O PACKAGE, Alan Thompson, IBM July 14, 1970 pages 13 and 14.	Defined in system routine DREAD or DWRITE when they encounter I/O error.	ERROR d.

/FLXBLK/ (Version 1)

COMMON/FLXBLK/AVFLX(675),DFLX(675),KP(675)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
AVFLX (675)	R*8	000000	Three solar rotation midpoint average 10.7cm flux values. Average provided daily beginning with 12 hours GMT two days prior to day of data epoch.	JANTHG	NONAME ADFLUX BMTWRT DENSTY INDENT JANTHG NEWARC
DFLX (675)	R*8	001518	Daily 10.7cm flux values beginning 12 hours GMT two days prior to day of data epoch.	JANTHG	NONAME ADFLUX BMTWRT DENSTY INDENT JANTHG NEWARC
KP (675)	R*8	002A30	Daily mean of geomagnetic indices K_p beginning 12 hours GMT two days prior to day of data epoch	JANTHG	NONAME ADFLUX BMTWRT DENSTY INDENT JANTHG NEWARC

/FLXBLK/ (Version 2)

COMMON/FLXBLK/INDXCS, (960,3) PLHSIG, PLHSW

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
INDXCS (960,3)	I*2	000000	Indexes for common adjusted geopotential coefficients. INDXCS(I,1)=degree of Ith adjusted coefficient. INDXCS(I,2)=Order of Ith adjusted coefficient. INDXCS(I,3) = 1 for C's = 2 for S's	INOIPT	COMPAR INOIPT STORE
PLHSIG	R*4	001680	Sigmas and correlations on adjusted station positions.	INOIPT	COMPAR INOIPT STORE
PLHSW	L*1	003A38	Switches telling whether adjusted station sigmas and correlations refer to Cartesian or geodetic coordinates	INOIPT	COMPAR INOIPT STORE

/FLXBLK/ (VERSION 3)
 COMMON/FLXBLK/BSTRT(900),BSEND(900),BTYPE(900)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Program Where Used</u>
BSTRT (900)	R*8	000000	Start time for bias adjustment coverage	INOIPT BIAS	ARCPAR BIAS INOIPT STORE
BSEND (900)	R*8	001C20	Stop time for bias adjustment coverage	INOIPT BIAS	ARCPAR BIAS INOIPT STORE
BTYPE (900)	I*2	003840	Types for bias adjustment	INOIPT BIAS	ARCPAR BIAS INOIPT STORE

/FMODEL/

COMMON/FMODEL/INDEX1, INDEX2, INDEX3, INDEX4, CS(30,33),
MODEL(8)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
INDEX1	I*4	000000	Maximum degree of geopotential plus 1.	BLOCK DATA INOIPT	COEFL DATARD F GEOIDH INOIPT NEWARC STORE
INDEX2	I*4	000004	Maximum degree and order of Legendre polynomials to be computed.	F	AVGPOT DATARD EGRAV F GEOIDH STORE
INDEX3	I*4	000008	Maximum order of geopotential plus 1.	BLOCK DATA INOIPT	COEFL DATARD EGRAV INOIPT NEWARC STORE
INDEX4	I*4	00000C	Maximum degree and order of geopotential for variational equations.	BLOCK DATA INOIPT	DATARD F INOIPT STORE VEVAL
CS (30,33)	R*8	000010	Geopotential coefficients.	BLOCK DATA INOIPT NONAME	NONAME ARCPAR AVGPOT COEFL COMADJ COMPAR EGRAV GEOIDH INOIPT ORB1 VEVAL
MODEL (8)	R*8	001F00	Name of geopotential,	BLOCK DATA	COEFL

/GEODYN/

COMMON/GEODYN/DATE

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
DATE	R*8	000000	GEODYN system date and source tape number.	BLOCK DATA	TYPORB

/GNDTRK/

COMMON/GNDTRK/SATLAT(2),SATLON(2),SATH(2),ELEV(2),SATSW

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
SATLAT (2)	R*8	000000	Satellite latitude.	PREDCT	NONAME PREDCT
SATLON (2)	R*8	000010	Satellite longi- tude.	PREDCT	NONAME PREDCT
SATH (2)	R*8	000020	Satellite height.	PREDCT	NONAME PREDCT
ELEV (2)	R*8	000030	Satellite elevation angle.	PREDCT	NONAME PREDCT PROCES
SATSW	L*4	000040	Switch requesting —computation of SATLAT,SATLON, SATH.	NONAME	NONAME

/INITBK/

COMMON/INITBK, IEPYMD, IEPHM, EPSEC, IYREF, INNMAX, INNMIN,
 CONVRG, ORBEL(6,2), EDITN, INSUPR, IDSAT(2), ORBTSW,
 XYZFSW, SYZLSW, PLTLSW, GRDFSW, KEPLER, SUBSAT, PARTGP,
 PBMAT, BMATNO, SIMDAT, PCESIM, MISLOG(9)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
IEPYMD	I*4	000000	Epoch YYMMDD	MAIN DODELM ORBIT	MAIN NONAME DATARD DODELM INOIPT ORBIT STORE TYPORB
IEPHM	I*4	000004	Epoch HHMM	MAIN DODELM ORBIT	MAIN NONAME DATARD DODELM INOIPT ORBIT STORE TYPORB
EPSEC	R*8	000008	Epoch seconds	MAIN DODELM ORBIT	MAIN NONAME DATARD DODELM INOIPT ORBIT STORE TYPORB
IYREF	I*4	000010	Reference date	MAIN	MAIN NONAME DATARD INOIPT NEWARC STORE
INNMAX	I*4	000014	Maximum number of inner iterations	INOIPT	MAIN NONAME DATARD INOIPT NEWARC STORE

/INITBK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
INNMIN	I*4	000018	Minimum number of inner iterations	INOIPT	MAIN NONAME DATARD INOIPT NEWARC STORE
CONVRG	R*4	00001C	Inner iteration convergence criterion	INOIPT	MAIN NONAME DATARD INOIPT NEWARC STORE
ORBEL (6,2)	R*8	000020	Nominal epoch elements	MAIN	MAIN NONAME DATARD INOIPT ORBIT STORE
EDITN	R*4	000080	Editing multiplier	INOIPT	NONAME DATARD INOIPT STORE SUMMARY
INSUPR	I*4	000084	Residual printing indicator	MAIN INOIPT NEWARC	MAIN NONAME DATARD INOIPT NEWARC STORE
IDSAT (2)	I*4	000088	Satellite ID's for ORB1 tapes	INOIPT	NONAME DATARD INOIPT STORE
ORBTSW	L*4	000090	Orbit generator switch	INOIPT NEWARC	MAIN NONAME DATARD INOIPT STORE TYPORB

/INITBK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
XYZFSW	L*4	000094	Switch for satellite ephemeris on first inner of first outer	INOIPT NEWARC	NONAME DATARD INOIPT STORE
XYZLSW	L*4	000098	Switch for satellite ephemeris on last inner of last outer	INOIPT NEWARC	NONAME DATARD INOIPT STORE
PLTLSW	L*4	00009C	Binary residual tape switch	INOIPT NEWARC	NONAME DATARD INOIPT STORE
GRDFSW	L*4	0000A0	Groundtrack tape switch	INOIPT NEWARC	NONAME DATARD INOIPT STORE
KEPLER	L*4	0000A4	Kepler ephemeris switch	INOIPT NEWARC	NONAME DATARD INOIPT STORE
SUBSAT	L*4	0000A8	Satellite ground-track switch	NONAME NEWARC	NONAME DATARD PREDCT STORE
PARTGP	L*4	0000AC	Partial derivative print switch	INOIPT NEWARC	NONAME DATARD INOIPT STORE
PBMAT	L*4	0000B0	B-matrix print switch	INOIPT NEWARC	BMTWRT DATARD INOIPT STORE
BMATNO	I*4	0000B4	B-matrix number	INOIPT NEWARC	NONAME BMTWRT DATARD ESTIM INOIPT STORE

/INITBK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
SIMDAT	L*4	0000B8	Simulated data switch	INOIPT NEWARC	NONAME DATARD INOIPT STORE
PCESIM	L*4	0000BC	Simulated element data type switch	INOIPT	NONAME DATARD INOIPT STORE
MISLOG (9)	I*4	0000C0	Initialization switches for several subroutines 1 - DENSTY 2 - DPFCT 3 - EGRAV 4 - EPHEM 5 - NUTATE 6 - PRECES 7 - REFCOR 8 - TDIF 9 - TIDAL	DATARD GEOIDH	NONAME DATARD DENSTY DPFCT EGRAV EPHEM GEOIDH NUTATE PRECES REFCOR TDIF TIDAL

/INTBLK/

COMMON/INTBLK/THDOT1,THDOT2,THDT2S,GM,AE,AESQ,FLAT,
 FSQ32,FFSQ32,GM3(6),B(2),BDOT(2),BO(2),APGM(2),APLM(2),
 RPRESS,INITAL,NORRAT,THETGO,MBODY(6),STEPSZ(2,2),
 HLVERB(2),DBLERB(2),CTOL(2),RTOL(2),STPLOW(2),STEPUP(2),
 ORDER(2,2),ASAT(2),MSAT(2),VARSTP(2),HLVDSW(2),NEQN(2),
 ADDR(2),ADDRD(2),SRAD(2),LOVE(3),TOREFT,NBODY

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
THDOT1	R*8	000000	Mean advance in right ascension of Greenwich in radians per mean solar day.	BLOCK DATA NEWARC	F GRHRAN NEWARC ORBI
THDOT2	R*8	000008	Rotation rate of earth in radians per mean solar day.	NEWARC	F GRHRAN NEWARC
THDT2S	R*8	000010	Rotation rate of earth in radians per second.	NEWARC	AVGPOT DRAG EPHEM GEOIDH NEWARC OBSDOT PREDCT VEVAL
GM	R*8	000018	Universal gravitational constant times mass of earth.	BLOCK DATA INOIPT	AVGPOT EGRAV ELEM GEOIDH INOIPT OBSDOT ORBI ORBIT POSVEL START TYPORB VEVAL

/INTBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
AE	R*8	000020	Semimajor axis of earth ellipsoid.	BLOCK DATA INOIPT	APPER AREAS AVGPOT DELTAZ DRAG EGRAV GEOIDH INDENT INOIPT ORB1 ORBIT PLHOUT PREDCT SQUANT STAINP TIDAL TYPORB
AESQ	R*8	000028	AE * AE	INOIPT	AREAS F INOIPT ORBIT
FLAT	R*8	000030	Flattening of earth ellipsoid.	BLOCK DATA INOIPT	AREAS AVGPOT DELTAZ INDENT INOIPT PLHOUT PREDCT SQUANT STAINP
FSQ32	R*8	000038	$3/2 AE * FLAT^{**2}$	INOIPT	APPER DRAG INOIPT PREDCT VEVAL
FFSQ32	R*8	000040	AE * FLAT + FSQ32	INOIPT	APPER DRAG INOIPT PREDCT VEVAL

/INTBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
GM3 (6)	R*8	000048	Universal gravitational constant. times mass of perturbing celestial bodies ordered as follows: 1 - MOON 2 - SUN 3 - VENUS 4 - MARS 5 - JUPITER 6 - SATURN	ORBIT	ORBIT SUNGRV TIDAL VEVAL
B(2)	R*8	000078	$1/2 C_D A/m$	ORBIT	COWELL DRAG F ORBIT TIDAL VEVAL
BDOT (2)	R*8	000088	$1/2 \dot{C}_D A/m$	ORBIT	DRAG F ORBIT TIDAL
B0(2)	R*8	000098	$1/2 A/m$	ORBIT	NONAME DRAG F ORBIT TIDAL
-APGM (2)	R*8	0000B8	$P_R A/m$ (P_R = solar radiation pressure constant)	ORBIT	F ORBIT TIDAL
APLM (2)	R*8	0000A8	$C_R P_R A/m$	ORBIT	F ORBIT TIDAL

/INTBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
RPRESS	R*8	0000C8	Solar radiation pressure constant.	BLOCK DATA	F INOIPT ORBIT TIDAL
INITAL	L*4	0000D0	Integration initialization switch.	NONAME	NONAME COWELL F ORBIT TIDAL
NORRAT	L*4	0000D4	No range rate switch.	GEOSRD DODSRD	NONAME DATARD DODSRD F GEOSRD PCERD SIMRD TIDAL
THETGO	R*2	0000D8	Right ascension of Greenwich on Jan 0.0 of reference year.	JANTHG	NONAME ADFLUX DATARD F GRHRAN ORBI STORE TIDAL
MBODY (6)	R*8	0000E0	Ratios of masses of perturbing bodies to mass of earth ordered as follows: <ol style="list-style-type: none"> 1 - MOON 2 - SUN 3 - VENUS 4 - MARS 5 - JUPITER 6 - SATURN 	BLOCK DATA INOIPT	DATARD INOIPT NEWARC ORBI ORBIT STORE TIDAL

/INTBLK/ (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
STEPSZ (2,2)	R*8	000110	Integration step size.	BLOCK DATA INOIPT NEWARC ORBIT	NONAME DATARD INOIPT NEWARC ORBIT STORE TIDAL
HLVERB (2)	R*8	000130	Step size reduction criterion.	BLOCK DATA INOIPT NEWARC	COWELL DATARD INOIPT NEWARC STORE TIDAL
DBLERB (2)	R*8	000140	Step size increase criterion.	BLOCK DATA INOIPT NEWARC	COWELL DATARD INOIPT NEWARC STORE TIDAL
CTOL (2)	R*8	000150	Critical tolerance for determining number of integrator correction iterations.	BLOCK DATA INOIPT NEWARC	COWELL DATARD INOIPT NEWARC STORE TIDAL
RTOL (2)	R*8	000160	Tolerance used to determine new step size in vary-step integrator.	BLOCK DATA INOIPT NEWARC	COWELL DATARD INOIPT NEWARC STORE TIDAL
STPLOW (2)	R*8	000170	Minimum integrator step size.	BLOCK DATA INOIPT NEWARC	COWELL DATARD INOIPT NEWARC STORE TIDAL

/INTBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
STEPUP (2)	R*8	000180	Maximum integrator step size.	BLOCK DATA INOIPT NEWARC	COWEIL DATARD INOIPT NEWARC STORE TIDAL
ORDER (2,2)	I*4	000190	Integration orders.	BLOCK DATA INOIPT NEWARC	MAIN DATARD INOIPT NEWARC ORBIT STORE TIDAL
ASAT (2)	R*8	0001A0	Satellite areas.	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC ORB1 ORBIT STORE TIDAL TYPORB
MSAT (2)	R*8	0001B0	Satellite masses.	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC ORB1 ORBIT STORE TIDAL TYPORB
VARSTP (2)	L*4	0001C0	Vary-step switch.	INOIPT NEWARC	MAIN DATARD ERROR INOIPT NEWARC ORBIT STORE TIDAL

/INTBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
HLVDSW (2)	L*4	0001C8	Halving-doubling switch.	INOIPT NEWARC	COWELL DATARD INOIPT NEWARC STORE TIDAL
NEQN (2)	I*4	0001D0	Number of integrator equations.	NONAME	MAIN NONAME DATARD INOIPT NEWARC ORBIT PREDCT RESPAR STORE SURDEN TIDAL VEVAL
ADDR (2)	I*4	0001D8	Drag parameter numbers.	INOIPT	NONAME ARCPAR BMTERT DATARD DRAG INOIPT NEWARC ORBIT PREDCT STORE SURDEN TIDAL TYPORB VEVAL

/INTBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
ADDRD (2)	I*4	0001E0	Drag rate parameter numbers.	INOIPT	NONAME ARCPAR BMTWRT DATARD DRAG INOIPT NEWARC ORBIT PREDCT STORE SURDEN TIDAL TYPORB VEVAL
SRAD (2)	I*4	0001E8	Solar radiation parameter numbers.	INOIPT	NONAME ARCPAR BMTWRT DATARD F INOIPT NEWARC ORBIT PREDCT STORE SURDEN TIDAL TYPORB VEVAL
LOVE (3)	R*8	0001F0	Tidal parameter values.	BLOCK DATA INOIPT	MAIN ARCPAR DATARD INOIPT STORE SURDEN TIDAL TYPORB VEVAL

/INTBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
TOREFT	L*4	000208	Switch for output in true equator and equinox of reference time.	INOIPT NEWARC	MAIN NONAME ARCPAR DATARD INOIPT NEWARC ORBIT STORE SURDEN TYPORB VEVAL
NBODY	I*4	00020C	Number of perturbing celestial bodies.	BLOCK DATA INOIPT NEWARC	MAIN ARCPAR DATARD EPHEM INOIPT NEWARC STORE SUNGRV SURDEN TYPORB VEVAL

/INTERP/

COMMON/INTERP/COMB(21,21),M12(4)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
COMB (21, 21)	R*8	000000	Binomial coefficient array.	COM	COEF COM
M12 (4)	I*4	000DC8	Displacement array used by integrator and interpolator.	COWELL ORBIT	ORBIT

/MONTHS/

COMMON/MONTHS/MONTH(26)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
MONTH (26)	I*4	000000	Number of elapsed days at the beginning of each month for leap years and for non-leap years.	BLOCK DATA	ADDYMD DIFF

/MOONGR/

COMMON/MOONGR/DPXUV(6), RHOM(3,6), RHOSQ(6), RHO3(6)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
DPXUV (6)	R*8	000000	Dot products of satellite position vector with unit vectors to perturbing celestial bodies.	SUNGRV	F SUNGRV TIDAL
RHOM (3,6)	R*8	000030	Satellite position vectors in coordinates centered at perturbing bodies.	SUNGRV	SUNGRV VEVAL
RHOSQ (6)	R*8	0000C0	Square of the difference of the distance of the satellite from the earth and the distance of the satellite from the perturbing body.	SUNGRV	SUNGRV VEVAL
RHO3 (6)	R*8	0000F0	RHOSQ**1.5	SUNGRV	SUNGRV VEVAL

/PREBLK/

COMMON/PREBLK/DAYSTA,OBS1,OBS2,SIG1,SIG2,SRFNDX,
 ISTA,MTYPE,NMEAS,ISAT,PRETYP,CHANEL,VHFCHN,
 PREPRO,RECNO

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
DAYSTA	R*8	000000	Time of observation pair	GEOSRD DODSRD PCERD SIMRD NONAME	NONAME BIAS DATARD DODSRD GEOSRD NEWARC PCERD RANDOM SIMRD STORE
OBS1	R*8	000008	First observation	GEOSRD DODSRD PCERD SIMRD NONAME PROCES TWOSTA	NONAME DODSRD GEOSRD PCERD PREDCT PROCES RANDOM SIMRD TWOSTA
OBS2	R*8	000010	Second observation	PROCES TWOSTA	NONAME DODSRD GEOSRD PCERD PREDCT PROCES RANDOM SIMRD TWOSTA
SIG1	R*8	000018	Sigma for first observation	GEOSRD DODSRD PCERD SIMRD NONAME	NONAME DODSRD GEOSRD PCERD PREDCT RANDOM SIMRD

/PREBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
SIG2	R*8	000020	Sigma for second observation	PROCES TWOSTA	NONAME DODSRD GEOSRD PCERD PREDCT RANDOM SIMRD TWOSTA
SRFNDX	R*8	000028	Tropospheric re-fraction index	PROCES TWOSTA	NONAME DODSRD GEOSRD PCERD PREDCT PROCES RANDOM SIMRD TWOSTA
ISTA	I*4	000030	Station index	PROCES TWOSTA	NONAME BIAS DODSRD GEOSRD INOIPT PCERD RANDOM SIMRD
MTYPE	I*2	000034	Measurement type	PROCES TWOSTA	MAIN NONAME BIAS DODSRD GEOSRD INOIPT PCERD PREDCT PROCES RANDOM SIMRD TWOSTA

/PREBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
NMEAS	I*2	000036	Number of measurements (1 or 2)	PROCES TWOSTA	NONAME DODSRD GEOSRD PCERD PREDCT RANDOM SIMRD
ISAT	I*2	000038	Satellite index	PROCES TWOSTA	NONAME DODSRD INOIPT OBSDOT PCERD PREDCT PROCES RANDOM SIMRD TWOSTA
PRETYP	I*2	00003A	Preprocessing indicators	PROCES TWOSTA	DODSRD GEOSRD PCERD PREDCT PROCES RANDOM SIMRD TWOSTA
CHANEL	I*2	00003C	Transponder channel	PROCES TWOSTA	NONAME BIAS DODSRD GEOSRD PCERD PROCES RANDOM SIMRD TWOSTA
VHFCHN	L*1	00003E	Switch to indicate VHF transponder	PROCES TWOSTA	DODSRD GEOSRD PCERD PROCES RANDOM SIMRD

/PREBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
PREPRO	L*1	00003F	Preprocessing switch	PROCES TWOSTA	NONAME DODSRD GEOSRD PCERD PREDCT PROCES RANDOM SIMRD TWOSTA
RECNO	I*4	000040	Measurement record number	PROCES TWOSTA	NONAME DODSRD GEOSRD INOPT PCERD RANDOM SIMRD

/PRIORI/

COMMON/PRIORI/ELEMIN(6,2),VARCOV(6,6,2),TITLE(60),
DRAGSG(2,3),DRAGO(2,3),CD(2,3)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
ELEMIN (6,2)	R*8	000000	A priori epoch elements.	MAIN ORBIT	MAIN NONAME BMTWRT DATARD ESTIM INOIPT ORBIT STORE
VARCOV (6,6,2)	R*4	000060	A priori variance/ covariance matrix of the epoch elements.	INOIPT NEWARC	DATARD ESTIM INOIPT NEWARC STORE
TITLE (60)	R*8	000180	Title of arc.	MAIN	MAIN NONAME DATARD INOIPT STORE TYPORB
DRAGSG (2)	R*8	000270	A priori drag sigmas.	INOIPT	NONAME ARCPAR DATARD INOIPT NEWARC STORE
DRGDSG (2)	R*8	000280	A priori drag rate sigmas.	INOIPT	NONAME ARCPAR DATARD INOIPT NEWARC STORE

/PRIORI/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
EMISSG (2)	R*8	000290	A priori solar radiation pressure sigmas.	INOIPT	NONAME ARCFAR DATARD INOIPT NEWARC STORE
DRAGO (2)	R*8	0002A0	A priori drag coefficients, C_D .	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC STORE
DRGDO (2)	R*8	0002B0	A priori drag rate coefficients, C_D .	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC STORE
EMISS0 (2)	R*8	0002C0	A priori solar radiation pressure coefficients, C_R .	INOIPT NEWARC	NONAME DATARD INOIPT NEWARC STORE
CD (2)	R*8	0002D0	Adjusted drag coefficients C_D .	NONAME INOIPT NEWARC UPDATE	NONAME ARCPAR BMTWRT DATARD INOIPT NEWARC ORBI ORBIT STORE TYPORB UPDATE

/PRIORI/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
CDD (2)	R*8	0002E0	Adjusted drag rate coefficients, C_D .	NONAME INOPT NEWARC UPDATE	NONAME ARCPAR BMTWRT DATARD INOPT NEWARC ORB1 ORBIT STORE TYPORB UPDATE
EMISS (2)	R*8	0002F0	Adjusted solar radiation pressure coefficients, C_R .	NONAME INOPT NEWARC UPDATE	NONAME ARCPAR BMTWRT DATARD INOPT NEWARC ORB1 ORBIT STORE TYPORB UPDATE

/SIGBLK/

COMMON/SIGBLK/SIGSTD(30),SGPRNT(30),IARRAY(4)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
SIGSTD (30)	R*4	000000	Default measurements sigmas.	BLOCK DATA	DODSRD GEOSRD INOIPT PCERD
SGPRNT (30)	R*4	000078	Measurement type sigmas to be printed.	INOIPT	INOIPT DODSRD
IARRAY (4)	I*4	0000F0	Input data tape numbers.	INOIPT	GEOSRD INOIPT PCERD SIMRD

/SRFBLK/ (Version 1)

COMMON/SRFBLK/DUMMY(5400),BESINO(675),BETYPE(675)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
Single Words 1-5400	R*4	000000	Scratch		GEOSRD
BESTNO (675)	I*2	005460	Station number for which electronic biases will be extracted	INOUP	INOUP
BETYPE (675)	I*2	0059A6	Measurement types for electronic bias extraction	INOUP	INOUP

/SRFBLK/ (Version 2)

COMMON/SRFBLK/PHI(675),XLAM(675),DP(675),SD(675),
SSD(675),NP(675),NL(675),NSD

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
PHI (675)	R*8	000000	Latitudes of centers of master surface density blocks	INOUP	INOUP PDEN 1
XLAM (675)	R*8	001518	Longitude of cen- ters of master surface density blocks	INOUP	INOUP PDEN 1
DP (675)	R*4	002A30	Surface density block latitude increments	INOUP	INOUP PDEN 1
DL (675)	R*4	0034BC	Surface density block longitude increments	INOUP	INOUP PDEN 1
SD (675)	R*4	003F48	Surface density values	INOUP	INOUP PDEN 1
SSD (675)	R*4	0049D4	Surface density <u>a priori</u> sigmas	INOUP	INOUP PDEN 1
NP (675)	I*2	005460	Numbers of latitude divisions for sur- face density master blocks	INOUP	INOUP PDEN 1
NL (675)	I*2	0059A6	Numbers of longitude divisions for sur- face density master blocks	INOUP	INOUP PDEN 1
NSD	I*4	005EEC	Number of surface density master blocks	INOUP	INOUP PDEN 1

/STANUM/

COMMON/STANUM/NAME, STANOS, NOSTOR

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
NAME	R*8	000000	Station names.	BLOCK DATA	DODSRD STAINP
STANOS	I*2	0008C0	Station numbers.	BLOCK DATA	DODSRD GEOSRD SIMRD STAINP
NOSTOR	I*4	000AF0	Number of stored stations.	BLOCK DATA	DODSRD GEOSRD SIMRD STAINP

/STAPOS/

COMMON/STAPOS/LAT(280),LON(280),HT(280)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
LAT (280)	R*8	000000	Station geodetic latitude in radians.	BLOCK DATA	STAINP
LON (280)	R*8	0008C0	Station geodetic longitude in radians.	BLOCK DATA	STAINP
HT (280)	R*8	001180	Station height in radians.	BLOCK DATA	STAINP

/TPEBLK/

COMMON/TPEBLK/INTP, OUTP, DATP, XYZTP, KEPTAP, RVTP,
PLOTP, IOBS, SCRA, SCRC, FLTP, GRDTP

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
INTP	I*4	000000	Card input unit-5.	BLOCK DATA	MAIN ADFLUX ALIST GEOSRD INOIPT STAINP
OUTP	I*4	000004	Printer output unit-6.	BLOCK DATA	NONAME ALIST BMTWRT COMADJ COMPAR CORREL INOIPT PDEN PDEN1 SUMMRY TYPORB UPDATE
DATP	I*4	000008	Direct access data storage disk unit-12.	BLOCK DATA	not used
XYZTP	I*4	00000C	Printer output unit for XYZ ephemeris 6, 8 or 9.	BLOCK DATA INOIPT	NONAME DATARD NEWARC STORE
KEPTAP	I*4	000010	Printer output unit for Kepler ephemeris 6,8, or 9.	BLOCK DATA INOIPT	NONAME DATARD INOIPT NEWARC STORE
RVTP	I*4	000014	RV tape unit.	BLOCK DATA INOIPT	NONAME DATARD INOIPT NEWARC STORE

/TPEBLK/(Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
PLOTP	I*4	000018	Binary residual tape unit.	BLOCK DATA	NONAME
IOBS	I*4	00001C	Data selection indicator.	INOUP	DODSRD GEOSRD INOUP NEWARC
SCRA	I*4	000020	Scratch disk for a priori information-14.	BLOCK DATA	NONAME ARCPAR DATARD
SCRC	I*4	000024	Scratch disk for normal matrix-16.	BLOCK DATA	NONAME ARCPAR COMPAR DATARD INOUP STORE
FLTP	I*4	000028	Scratch disk for flux data-13.	BLOCK DATA	NONAME ADFLUX
GRDTP	I*4	00002C	Groundtrack tape unit-18.	BLOCK DATA	NONAME

/TRUPOL/

COMMON/TRUPOL/TRUE

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
TRUE	R*8	000000	Coordinates of tracking station corrected from the position of the true pole.	TRUEP	GRHRAN TRUEP

/XYZ/

COMMON/XYZ/ELEM(6),R,RSQ,ISAT,IFORCE(2)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
ELEM (6)	R*8	000000	Satellite Cartesian elements.	F	NONAME AVGPOT DENSTY DRAG EGRAV F GEOIDH RESPAR SUNGRV SURDEN TIDAL VEVAL
R	R*8	000030	Satellite earth centered position radius.	EGRAV	NONAME AVGPOT DENSTY DRAG EGRAV GEOIDH RESPAR SURDEN TIDAL VEVAL
RSQ	R*8	000038	R*R	EGRAV	NONAME DRAG EGRAV RESPAR SUNGRV SURDEN VEVAL
ISAT	I*4	000040	Index of satellite orbit being integrated.	ORBIT	DRAG F ORBIT RESPAR SURDEN VEVAL
IFORCE (2)	I*4	000044	Displacements in partial array of drag and solrad partials.	NONAME	NONAME DRAG F

/XYZOUT/

COMMON/XYZOUT/XYZEND(6,2),DRGPAR(6,2)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Program Where Defined</u>	<u>Programs Where Used</u>
XYZEND (6,2)	R*8	000000	Cartesian satellite coordinates corresponding to time of integration call.	ORBIT	NONAME GRHRAN OBSDOT ORB1 ORBIT PREDCT TYPORB
DRGPAR (6,2)	R*8	000060	Cartesian drag partials corresponding to time of integration call.	ORBIT	GRHRAN OBSDOT ORBIT PREDCT TYPORB

/VMAT/

Common block VMAT is all scratch and used by subroutines
COWELL, INOUPPT and VEVAL.

/VRBLOK/ (Version 1)

COMMON/VRBLOK/GPSIG(960)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
GPSIG (960)	R*4	000000	Sigmas on common adjusted geopo- tential coef- ficients.	INOIPT	COMPAR INOIPT STORE

/VRBLOK/ (Version 2)

COMMON/VRBLOK/BIASSO(900),BIASSG(900),BSTANO(900)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
BIASSO (900)	R*4	000000	<u>A priori value</u> of adjusted biases.	INOUP T BIAS	ARCPAR BIAS INOUP T STORE
BIASSG (900)	R*4	000E10	<u>A priori sigmas</u> of adjusted biases.	INOUP T BIAS	ARCPAR BIAS INOUP T STORE
BSTANO	I*2	001C20	Station numbers for adjusted biases.	INOUP T BIAS	ARCPAR BIAS INOUP T STORE

/VRBLOK/ (Version 3)

COMMON/VRBLOK/A1,IBUF,AORN

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Defined</u>
A1(66)	R*8	000000	Dummy	---	---
IBUF (1980)	I*4	000210	Scratch	DATARD	DATARD
AORN (69)	R*8	002100	Dummy	---	---

/VRBLOK/ (Version 4)

COMMON/VRBLOK/XYSQ,COSLAM(31),SINLAM(31,PR,PPSI,
PLAMDA,P(33,30),AORN(30),TPSIM(39)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
XY SQ	R*8	000000	X**2+Y**2 where X and Y are the earth centered X&Y coordinate of the satellite.	E GRAV	CBROWN E GRAV INDENT NEWARC RESPAR VEVAL
COSLAM (31)	R*8	000008	The cosines of m*longitude of the satellite for m=0,30.	CBROWN E GRAV	CBROWN E GRAV GEOIDH INDENT NEWARC RESPAR VEVAL
SINLAM (31)	R*8	000100	The sines of m*longitude of the satellite for m=0,30.	E GRAV	CBROWN E GRAV GEOIDH INDENT NEWARC RESPAR VEVAL
PR	R*8	0001F8	Partial of the earth potential with respect to the radial direction	E GRAV	CBROWN E GRAV INDENT NEWARC VEVAL
PPSI	R*8	000200	Partial of the earth potential with respect to geocentric latitude.	E GRAV	CBROWN E GRAV INDENT NEWARC VEVAL
PLAMDA	R*8	000208	Partial of the earth potential with respect to east longitude.	E GRAV	CBROWN E GRAV INDENT NEWARC VEVAL

/VRBLOK/ (Version 4) (Cont.)

<u>Variable</u>	<u>Type</u>	<u>Hex Location</u>	<u>Description</u>	<u>Programs Where Defined</u>	<u>Programs Where Used</u>
P(33,30)	R*8	000210	The Legendre and associated Legendre polynomials of the spherical harmonic expansion. P(m+1,n) is the polynomial associated with degree n and order m.	EGRAV	AVGPOT BMTWRT EGRAV GEOIDH INDENT NEWARC RESPAR VEVAL
AORN(30)	R*8	002100	$AORN(n) = \left(\frac{a_e}{R}\right)^n$ where R is the distance from the earth's center to the point where the potential is being evaluated. A_e is the semi-major axis of the earth, and n is the degree.	EGRAV	AVGPOT EGRAV GEOIDH NEWARC RESPAR VEVAL
TPSIM (30)	R*8	0021F0	The tangents of m times the geocentric latitude for m=0,30.	EGRAV	EGRAV NEWARC RESPAR VEVAL